


Assignment 5 - due Tue Sept. 26 - Mainly Chapters 9 - 12. "Families of orbits and their contacting envelopes."



The atoms of NIST or volcanoes of Io

1. Suppose one of the volcanoes on Jupiter's moon Io detonates in a constant gravity- $g(m \cdot s^{-2})$ vacuum sending equi-velocity $\pm v_0(m \cdot s^{-1})$ fragments off at initial elevation angles $\alpha=0^\circ, 15^\circ, 30^\circ, \dots, 75^\circ,$ and 90° with the latter one going straight up to an altitude of $y=h_0=1$ -unit in the attached graph and falling straight down.

- That one distance unit has what *mks*-value in terms of $g(m \cdot s^{-2})$ and $v_0(m \cdot s^{-1})$? $h_0 = \underline{\hspace{2cm}}$ ().
- Derive the parabolic time-coordinates $x(t) = \underline{\hspace{2cm}}, y(t) = \underline{\hspace{2cm}}$ in terms of $g(m \cdot s^{-2})$ and $v_0(m \cdot s^{-1})$ and elevation angle α .
- Derive the parabolic focus-locus coordinates $x_{foc} = \underline{\hspace{2cm}}, y_{foc} = \underline{\hspace{2cm}}$ in terms of $g(m \cdot s^{-2})$ and $v_0(m \cdot s^{-1})$ and elevation angle α for $h_0=1$ and construct its curve on graph. (This curve has aspects of Thales geometry (subtended angle of circle diameter) that relate to trajectories. If you can show these below.)
- Derive the parabolic directrix coordinate $y_{dir} = \underline{\hspace{2cm}}$ in terms of $h_0=1$ and elevation angle α and construct this directrix line on graph for the cases $\alpha=0^\circ-90^\circ$ listed above.
- Give general parabolic trajectory curve function $y(x) = \underline{\hspace{2cm}}$ in terms of $g(m \cdot s^{-2})$ and $v_0(m \cdot s^{-1})$ and α for $h_0=1$. For the cases $\alpha=0^\circ, 30^\circ, 45^\circ,$ and 90° construct enough of their curve points and tangents to accurately represent them on the graph.
- Locate the envelope contact points for the cases $\alpha=0^\circ, 30^\circ, 45^\circ,$ and 90° and construct enough of the envelope points and tangents to accurately represent the envelope on the graph. If a contact point lies off the graph indicate where. Deduce $y_{envelope}(x) = \underline{\hspace{2cm}}$ in terms of $h_0=1$.
- Each parabola trajectory has kite-like structure (Recall Fig. 9.4.) as does envelope. Draw and relate them.
- Do any of the α -trajectories have the same shape as the envelope? If so, tell which one.

2. Now consider time behavior implicit in problem 1. In a "snapshot" of each moment, volcano fragments lie on "blast-front" curve. A geometric time unit T_1 is the time for the $\alpha=90^\circ$ fragment to reach its peak.

- That one time unit has what *mks*-value in terms of $g(m \cdot s^{-2})$ and $v_0(m \cdot s^{-1})$? $T_1 = \underline{\hspace{2cm}}$ ().
- Give a brief explanation addressing why this "snapshot" curve or locus has to be (whichever): a parabola? ___ straight line? ___ circle? ___ ellipse? ___ (Check one and explain choice on graph.)
- Derive and/or construct the "blast-front" curve for the case $\alpha=90^\circ$ at the moment when that fragment first contacts volcano envelope. Give time in T_1 units. $T_{90^\circ} = \underline{\hspace{2cm}}$ Find polar angle of contact normal.
- Derive and/or construct the "blast-front" curve for the case $\alpha=45^\circ$ at the moment when that fragment first contacts volcano envelope. Give time in T_1 units. $T_{45^\circ} = \underline{\hspace{2cm}}$ Find polar angle of contact normal.
- Derive and/or construct the "blast-front" curve for the case $\alpha=30^\circ$ at the moment when that fragment first contacts volcano envelope. Give time in T_1 units. $T_{30^\circ} = \underline{\hspace{2cm}}$ Find polar angle of contact normal.

3. Suppose fragments continue falling into a tunnel through moon-Io that has radius $R_{Io} = 0.5 \cdot 10^6 h_0$. Estimate radius of tunnel at widest point if it just big enough to let all fragments orbit without hitting its walls. $R_{tunnel} = \underline{\hspace{2cm}}$ (h_0)

Note: For this problem the gravity is not uniform constant $g=9.8ms^{-2}$ except near surface. (Ellipse geometry.)

