

Lecture 9
Thur. 9.19.2017

Equations of Lagrange and Hamilton mechanics in Generalized Curvilinear Coordinates (GCC)

(Ch. 12 of Unit 1 and Ch. 1-5 of Unit 2 and Ch. 1-5 of Unit 3)

Quick Review of Lagrange Relations in Lectures 7-8

Using differential chain-rules for coordinate transformations

Polar coordinate example of Generalized Curvilinear Coordinates (GCC)

Getting the GCC ready for mechanics: Generalized **velocity** and **Jacobian Lemma 1**

Getting the GCC ready for mechanics: Generalized **acceleration** and **Lemma 2**

How to say Newton's "F=ma" in Generalized Curvilinear Coords.

Use Cartesian KE quadratic form $KE=T=1/2\mathbf{v}\cdot\mathbf{M}\cdot\mathbf{v}$ and $\mathbf{F}=\mathbf{M}\cdot\mathbf{a}$ to get GCC force

Lagrange GCC trickery gives Lagrange force equations

Lagrange GCC trickery gives Lagrange potential equations (Lagrange 1 and 2)

GCC Cells, base vectors, and metric tensors

Polar coordinate examples: Covariant \mathbf{E}_m vs. Contravariant \mathbf{E}^m

Covariant g_{mn} vs. Invariant δ_m^n vs. Contravariant g^{mn}

Lagrange prefers Covariant g_{mn} with Contravariant **velocity**

GCC Lagrangian definition

GCC "canonical" momentum p_m definition

GCC "canonical" force F_m definition

Coriolis "fictitious" forces (... and weather effects)

Quick Review of Lagrange Relations in Lectures 7-8

 *0th and 1st equations of Lagrange and Hamilton*

Quick Review of Lagrange Relations in Lectures 7-8

0th and 1st equations of Lagrange and Hamilton

Starts out with simple demands for explicit-dependence, “loyalty” or “fealty to the colors”

*Lagrangian and Estrangian have no explicit dependence on **momentum** \mathbf{p}*

$$\frac{\partial L}{\partial \mathbf{p}_k} \equiv 0 \equiv \frac{\partial E}{\partial \mathbf{p}_k}$$

*Hamiltonian and Estrangian have no explicit dependence on **velocity** \mathbf{v}*

$$\frac{\partial H}{\partial \mathbf{v}_k} \equiv 0 \equiv \frac{\partial E}{\partial \mathbf{v}_k}$$

*Lagrangian and Hamiltonian have no explicit dependence on **speedium** \mathbf{V}*

$$\frac{\partial L}{\partial \mathbf{V}_k} \equiv 0 \equiv \frac{\partial H}{\partial \mathbf{V}_k}$$

Such non-dependencies hold in spite of “under-the-table” matrix and partial-differential connections

$$\begin{aligned} \nabla_{\mathbf{v}} L &= \frac{\partial L}{\partial \mathbf{v}} = \frac{\partial}{\partial \mathbf{v}} \frac{\mathbf{v} \cdot \mathbf{M} \cdot \mathbf{v}}{2} \\ &= \mathbf{M} \cdot \mathbf{v} = \mathbf{p} \end{aligned}$$

$$\begin{aligned} \nabla_{\mathbf{p}} H &= \mathbf{v} = \frac{\partial H}{\partial \mathbf{p}} = \frac{\partial}{\partial \mathbf{p}} \frac{\mathbf{p} \cdot \mathbf{M}^{-1} \cdot \mathbf{p}}{2} \\ &= \mathbf{M}^{-1} \cdot \mathbf{p} = \mathbf{v} \end{aligned}$$

(Forget Estrangian for now)

$$\begin{pmatrix} \frac{\partial L}{\partial v_1} \\ \frac{\partial L}{\partial v_2} \end{pmatrix} = \begin{pmatrix} m_1 & 0 \\ 0 & m_2 \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = \begin{pmatrix} p_1 \\ p_2 \end{pmatrix}$$

Lagrange’s 1st equation(s)

$$\frac{\partial L}{\partial v_k} = p_k \quad \text{or:} \quad \frac{\partial L}{\partial \mathbf{v}} = \mathbf{p}$$


$$\begin{pmatrix} \frac{\partial H}{\partial p_1} \\ \frac{\partial H}{\partial p_2} \end{pmatrix} = \begin{pmatrix} m_1^{-1} & 0 \\ 0 & m_2^{-1} \end{pmatrix} \begin{pmatrix} p_1 \\ p_2 \end{pmatrix} = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

Hamilton’s 1st equation(s)


$$\frac{\partial H}{\partial p_k} = v_k \quad \text{or:} \quad \frac{\partial H}{\partial \mathbf{p}} = \mathbf{v}$$

*p. 25 of
Lecture 8*


(a) *Lagrangian plot*
 $L(\mathbf{v}) = \text{const.} = \mathbf{v} \cdot \mathbf{M} \cdot \mathbf{v} / 2$




(b) *Hamiltonian plot*
 $H(\mathbf{p}) = \text{const.} = \mathbf{p} \cdot \mathbf{M}^{-1} \cdot \mathbf{p} / 2$




(c) *Overlapping plots*




(d) *Less mass*




(e) *More mass*



(a) *Lagrangian plot*
 $L(\mathbf{v}) = \text{const.} = \mathbf{v} \cdot \mathbf{M} \cdot \mathbf{v} / 2$



(b) *Hamiltonian plot*
 $H(\mathbf{p}) = \text{const.} = \mathbf{p} \cdot \mathbf{M}^{-1} \cdot \mathbf{p} / 2$




(c) *Overlapping plots*

1st equation of Lagrange

$$L = \text{const.} = E$$

1st equation of Hamilton

$$H = \text{const.} = E$$



(d) *Less mass*

(e) *More mass*

Using differential chain-rules for coordinate transformations

→ *Polar coordinate example of Generalized Curvilinear Coordinates (GCC)*

*Getting the GCC ready for mechanics: Generalized **velocity and Jacobian Lemma 1***

*Getting the GCC ready for mechanics: Generalized **acceleration and Lemma 2***

Using differential chain-rules[†] for coordinate transformations

A pair of 2-variable functions $f(x,y)$ and $g(x,y)$ can define a coordinate system on (x,y) -space

$$df(x,y) = \frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy$$

$$dg(x,y) = \frac{\partial g}{\partial x} dx + \frac{\partial g}{\partial y} dy$$

for example: polar coordinates

$$r^2(x,y) = x^2 + y^2 \quad \text{and} \quad \theta(x,y) = \text{atan2}(y,x)$$

(Not in text. Recall Lecture 8 p. 15-19)[†]

$$dr(x,y) = \frac{\partial r}{\partial x} dx + \frac{\partial r}{\partial y} dy$$

$$d\theta(x,y) = \frac{\partial \theta}{\partial x} dx + \frac{\partial \theta}{\partial y} dy$$

Using differential chain-rules[†] for coordinate transformations

A pair of 2-variable functions $f(x,y)$ and $g(x,y)$ can define a coordinate system on (x,y) -space

for example: polar coordinates

$$df(x,y) = \frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy$$
$$dg(x,y) = \frac{\partial g}{\partial x} dx + \frac{\partial g}{\partial y} dy$$
$$r^2(x,y) = x^2 + y^2 \quad \text{and} \quad \theta(x,y) = \text{atan2}(y,x)$$
$$dr(x,y) = \frac{\partial r}{\partial x} dx + \frac{\partial r}{\partial y} dy$$
$$d\theta(x,y) = \frac{\partial \theta}{\partial x} dx + \frac{\partial \theta}{\partial y} dy$$

(Not in text. Recall Lecture 8 p. 15-19)[†]

Easy to invert differential chain relations (even if functions are not easily inverted)

$$dx = \frac{\partial x}{\partial f} df + \frac{\partial x}{\partial g} dg$$
$$dy = \frac{\partial y}{\partial f} df + \frac{\partial y}{\partial g} dg$$
$$x = r \cos \theta$$
$$y = r \sin \theta$$
$$dx = \frac{\partial x}{\partial r} dr + \frac{\partial x}{\partial \theta} d\theta$$
$$dy = \frac{\partial y}{\partial r} dr + \frac{\partial y}{\partial \theta} d\theta$$
$$\begin{pmatrix} dx \\ dy \end{pmatrix} = \begin{pmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{pmatrix} \begin{pmatrix} dr \\ d\theta \end{pmatrix} = \begin{pmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{pmatrix} \begin{pmatrix} dr \\ d\theta \end{pmatrix}$$

Using differential chain-rules† for coordinate transformations

A pair of 2-variable functions $f(x,y)$ and $g(x,y)$ can define a coordinate system on (x,y) -space

$$df(x,y) = \frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy$$

for example: polar coordinates
 $r^2(x,y) = x^2 + y^2$ and $\theta(x,y) = \text{atan2}(y,x)$

$$dr(x,y) = \frac{\partial r}{\partial x} dx + \frac{\partial r}{\partial y} dy$$

$$dg(x,y) = \frac{\partial g}{\partial x} dx + \frac{\partial g}{\partial y} dy$$

(Not in text. Recall Lecture 8 p. 15-19)†

$$d\theta(x,y) = \frac{\partial \theta}{\partial x} dx + \frac{\partial \theta}{\partial y} dy$$

Easy to invert differential chain relations (even if functions are not easily inverted)

$$dx = \frac{\partial x}{\partial f} df + \frac{\partial x}{\partial g} dg$$

$$x = r \cos \theta$$

$$dx = \frac{\partial x}{\partial r} dr + \frac{\partial x}{\partial \theta} d\theta$$

$$dy = \frac{\partial y}{\partial f} df + \frac{\partial y}{\partial g} dg$$

$$y = r \sin \theta$$

$$dy = \frac{\partial y}{\partial r} dr + \frac{\partial y}{\partial \theta} d\theta$$

$$\begin{pmatrix} dx \\ dy \end{pmatrix} = \begin{pmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{pmatrix} \begin{pmatrix} dr \\ d\theta \end{pmatrix} = \begin{pmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{pmatrix} \begin{pmatrix} dr \\ d\theta \end{pmatrix}$$

Notation for differential GCC (Generalized Curvilinear Coordinates $\{q^1, q^2, q^3, \dots\}$)

$$dx^j = \frac{\partial x^j}{\partial q^m} dq^m \left(\equiv \sum_{m=1}^N \frac{\partial x^j}{\partial q^m} dq^m \quad \left\{ \text{Defining a shorthand dummy-index } m\text{-sum} \right\} \right)$$

What does "q" stand for?
 One guess: "Queer"
 And they do get pretty queer!

These x^j are plain old CC (Cartesian Coordinates $\{dx^1=dx, dx^2=dy, dx^3=dz, dx^4=dt\}$)

Using differential chain-rules† for coordinate transformations

A pair of 2-variable functions $f(x,y)$ and $g(x,y)$ can define a coordinate system on (x,y) -space

$$df(x,y) = \frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy$$

for example: polar coordinates
 $r^2(x,y) = x^2 + y^2$ and $\theta(x,y) = \text{atan2}(y,x)$

$$dr(x,y) = \frac{\partial r}{\partial x} dx + \frac{\partial r}{\partial y} dy$$

$$dg(x,y) = \frac{\partial g}{\partial x} dx + \frac{\partial g}{\partial y} dy$$

(Not in text. Recall Lecture 8 p. 15-19)†

$$d\theta(x,y) = \frac{\partial \theta}{\partial x} dx + \frac{\partial \theta}{\partial y} dy$$

Easy to invert differential chain relations (even if functions are not easily inverted)

$$dx = \frac{\partial x}{\partial f} df + \frac{\partial x}{\partial g} dg$$

$$x = r \cos \theta$$

$$dx = \frac{\partial x}{\partial r} dr + \frac{\partial x}{\partial \theta} d\theta$$

$$dy = \frac{\partial y}{\partial f} df + \frac{\partial y}{\partial g} dg$$

$$y = r \sin \theta$$

$$dy = \frac{\partial y}{\partial r} dr + \frac{\partial y}{\partial \theta} d\theta$$

$$\begin{pmatrix} dx \\ dy \end{pmatrix} = \begin{pmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{pmatrix} \begin{pmatrix} dr \\ d\theta \end{pmatrix} = \begin{pmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{pmatrix} \begin{pmatrix} dr \\ d\theta \end{pmatrix}$$

Notation for differential GCC (Generalized Curvilinear Coordinates $\{q^1, q^2, q^3, \dots\}$)

$$dx^j = \frac{\partial x^j}{\partial q^m} dq^m \left(\equiv \sum_{m=1}^N \frac{\partial x^j}{\partial q^m} dq^m \quad \left\{ \begin{array}{l} \text{Defining a shorthand} \\ \text{dummy-index } m\text{-sum} \end{array} \right\} \right)$$

What does "q" stand for?
 One guess: "Queer"
 And they do get pretty queer!

Connection lines may help to indicate summation (OK on scratch paper...Difficult in text)

These x^j are plain old CC (Cartesian Coordinates $\{dx^1=dx, dx^2=dy, dx^3=dz, dx^4=dt\}$)

Using differential chain-rules for coordinate transformations

Polar coordinate example of Generalized Curvilinear Coordinates (GCC)

- *Getting the GCC ready for mechanics: Generalized **velocity and Jacobian Lemma 1***
- Getting the GCC ready for mechanics: Generalized **acceleration and Lemma 2***

Getting the GCC ready for mechanics:

Generalized *velocity* relation follows from GCC chain rule

$$dx^j = \frac{\partial x^j}{\partial q^m} dq^m$$

Same kind of linear relation exists between CC velocity $v^j \equiv \dot{x}^j \equiv \frac{dx^j}{dt}$ and GCC velocity $v^m \equiv \dot{q}^m \equiv \frac{dq^m}{dt}$

$$\dot{x}^j = \frac{\partial x^j}{\partial q^m} \dot{q}^m$$

Getting the GCC ready for mechanics:

Generalized *velocity* relation follows from GCC chain rule

$$dx^j = \frac{\partial x^j}{\partial q^m} dq^m$$

Same kind of linear relation exists between CC velocity $v^j \equiv \dot{x}^j \equiv \frac{dx^j}{dt}$ and GCC velocity $v^m \equiv \dot{q}^m \equiv \frac{dq^m}{dt}$

This is a key “*lemma-1*” for setting up mechanics:

$$\dot{x}^j = \frac{\partial x^j}{\partial q^m} \dot{q}^m$$

or:

$$\frac{\partial \dot{x}^j}{\partial \dot{q}^m} = \frac{\partial x^j}{\partial q^m} \text{ lemma-1}$$

Getting the GCC ready for mechanics:

Generalized *velocity* relation follows from GCC chain rule

$$dx^j = \frac{\partial x^j}{\partial q^m} dq^m$$

Same kind of linear relation exists between CC velocity $v^j \equiv \dot{x}^j \equiv \frac{dx^j}{dt}$ and GCC velocity $v^m \equiv \dot{q}^m \equiv \frac{dq^m}{dt}$

$$\dot{x}^j = \frac{\partial x^j}{\partial q^m} \dot{q}^m$$

$$\frac{\partial \dot{x}^j}{\partial \dot{q}^m} = \frac{\partial x^j}{\partial q^m} \quad \text{lemma-1}$$

This is a key “*lemma-1*” for setting up mechanics:

Jacobian J_m^j matrix gives each CCC differential dx^j or velocity \dot{x}^j in terms of GCC dq^m or \dot{q}^m .

$$J_m^j \equiv \frac{\partial x^j}{\partial q^m} = \frac{\partial \dot{x}^j}{\partial \dot{q}^m} \quad \left\{ \begin{array}{l} \text{Defining } \textit{Jacobian} \\ \text{matrix component} \end{array} \right\}$$

Recall polar coordinate transformation matrix: $\begin{pmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{pmatrix} = \begin{pmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{pmatrix}$

Getting the GCC ready for mechanics:

Generalized *velocity* relation follows from GCC chain rule

$$dx^j = \frac{\partial x^j}{\partial q^m} dq^m$$

Same kind of linear relation exists between CC velocity $v^j \equiv \dot{x}^j \equiv \frac{dx^j}{dt}$ and GCC velocity $v^m \equiv \dot{q}^m \equiv \frac{dq^m}{dt}$

$$\dot{x}^j = \frac{\partial x^j}{\partial q^m} \dot{q}^m$$

$$\frac{\partial \dot{x}^j}{\partial \dot{q}^m} = \frac{\partial x^j}{\partial q^m} \quad \text{lemma-1}$$

This is a key “*lemma-1*” for setting up mechanics:

Jacobian J_m^j matrix gives each CCC differential dx^j or velocity \dot{x}^j in terms of GCC dq^m or \dot{q}^m .

$$J_m^j \equiv \frac{\partial x^j}{\partial q^m} = \frac{\partial \dot{x}^j}{\partial \dot{q}^m} \quad \left\{ \begin{array}{l} \text{Defining } \textit{Jacobian} \\ \text{matrix component} \end{array} \right\}$$

Recall polar coordinate transformation matrix: $\begin{pmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{pmatrix} = \begin{pmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{pmatrix}$

Inverse (so-called) *Kajobian* K_j^m matrix is flipped partial derivatives of J_m^j .

$$K_j^m \equiv \frac{\partial q^m}{\partial x^j} = \frac{\partial \dot{q}^m}{\partial \dot{x}^j} \quad \left\{ \begin{array}{l} \text{Defining } \textit{Kajobian} \\ \text{(inverse to Jacobian)} \end{array} \right\}$$

Polar coordinate inverse transformation matrix: $\begin{pmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{pmatrix}^{-1} = \begin{pmatrix} \frac{\partial r}{\partial x} & \frac{\partial r}{\partial y} \\ \frac{\partial \theta}{\partial x} & \frac{\partial \theta}{\partial y} \end{pmatrix} = \frac{\begin{pmatrix} r \cos \theta & r \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}}{(\det J = r)} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\frac{\sin \theta}{r} & \frac{\cos \theta}{r} \end{pmatrix}$

Defining 2x2 matrix inverse: (always test inverse matrices!)

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix}^{-1} = \frac{\begin{pmatrix} D & -B \\ -C & A \end{pmatrix}}{AD - BC}$$

Getting the GCC ready for mechanics:

Generalized *velocity* relation follows from GCC chain rule

$$dx^j = \frac{\partial x^j}{\partial q^m} dq^m$$

Same kind of linear relation exists between CC velocity $v^j \equiv \dot{x}^j \equiv \frac{dx^j}{dt}$ and GCC velocity $v^m \equiv \dot{q}^m \equiv \frac{dq^m}{dt}$

$$\dot{x}^j = \frac{\partial x^j}{\partial q^m} \dot{q}^m$$

$$\frac{\partial \dot{x}^j}{\partial \dot{q}^m} = \frac{\partial x^j}{\partial q^m} \quad \text{lemma-1}$$

This is a key “*lemma-1*” for setting up mechanics:

Jacobian J_m^j matrix gives each CCC differential dx^j or velocity \dot{x}^j in terms of GCC dq^m or \dot{q}^m .

$$J_m^j \equiv \frac{\partial x^j}{\partial q^m} = \frac{\partial \dot{x}^j}{\partial \dot{q}^m} \quad \left\{ \begin{array}{l} \text{Defining } \textit{Jacobian} \\ \text{matrix component} \end{array} \right\}$$

Recall polar coordinate transformation matrix: $\begin{pmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{pmatrix} = \begin{pmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{pmatrix}$

Inverse (so-called) *Kajobian* K_j^m matrix is flipped partial derivatives of J_m^j .

$$K_j^m \equiv \frac{\partial q^m}{\partial x^j} = \frac{\partial \dot{q}^m}{\partial \dot{x}^j} \quad \left\{ \begin{array}{l} \text{Defining } \textit{Kajobian} \\ \text{(inverse to Jacobian)} \end{array} \right\}$$

Polar coordinate inverse transformation matrix: $\begin{pmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{pmatrix}^{-1} = \begin{pmatrix} \frac{\partial r}{\partial x} & \frac{\partial r}{\partial y} \\ \frac{\partial \theta}{\partial x} & \frac{\partial \theta}{\partial y} \end{pmatrix} = \frac{\begin{pmatrix} r \cos \theta & r \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}}{(\det J = r)} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\frac{\sin \theta}{r} & \frac{\cos \theta}{r} \end{pmatrix}$

Defining 2x2 matrix inverse: (always test inverse matrices!)

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix}^{-1} = \frac{\begin{pmatrix} D & -B \\ -C & A \end{pmatrix}}{AD - BC} = \begin{pmatrix} \frac{D}{AD - BC} & \frac{-B}{AD - BC} \\ \frac{-C}{AD - BC} & \frac{A}{AD - BC} \end{pmatrix}$$

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} D & -B \\ -C & A \end{pmatrix} = \begin{pmatrix} AD - BC & 0 \\ 0 & AD - BC \end{pmatrix}$$

Getting the GCC ready for mechanics:

Generalized *velocity* relation follows from GCC chain rule

$$dx^j = \frac{\partial x^j}{\partial q^m} dq^m$$

Same kind of linear relation exists between CC velocity $v^j \equiv \dot{x}^j \equiv \frac{dx^j}{dt}$ and GCC velocity $v^m \equiv \dot{q}^m \equiv \frac{dq^m}{dt}$

$$\dot{x}^j = \frac{\partial x^j}{\partial q^m} \dot{q}^m$$

$$\frac{\partial \dot{x}^j}{\partial \dot{q}^m} = \frac{\partial x^j}{\partial q^m} \quad \text{lemma-1}$$

This is a key “*lemma-1*” for setting up mechanics:

Jacobian J_m^j matrix gives each CCC differential dx^j or velocity \dot{x}^j in terms of GCC dq^m or \dot{q}^m .

$$J_m^j \equiv \frac{\partial x^j}{\partial q^m} = \frac{\partial \dot{x}^j}{\partial \dot{q}^m} \quad \left\{ \begin{array}{l} \text{Defining } \textit{Jacobian} \\ \text{matrix component} \end{array} \right\}$$

Recall polar coordinate transformation matrix: $\begin{pmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{pmatrix} = \begin{pmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{pmatrix}$

Inverse (so-called) *Kajobian* K_j^m matrix is flipped partial derivatives of J_m^j .

$$K_j^m \equiv \frac{\partial q^m}{\partial x^j} = \frac{\partial \dot{q}^m}{\partial \dot{x}^j} \quad \left\{ \begin{array}{l} \text{Defining } \textit{Kajobian} \\ \text{(inverse to Jacobian)} \end{array} \right\}$$

$$\begin{pmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{pmatrix}^{-1} = \begin{pmatrix} \frac{\partial r}{\partial x} & \frac{\partial r}{\partial y} \\ \frac{\partial \theta}{\partial x} & \frac{\partial \theta}{\partial y} \end{pmatrix} = \frac{\begin{pmatrix} r \cos \theta & r \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}}{(\det J = r)} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\frac{\sin \theta}{r} & \frac{\cos \theta}{r} \end{pmatrix}$$

Product of matrix J_m^j and K_j^m is a unit matrix by definition of partial derivatives. (*always test inverse matrices!*)

$$K_j^m \cdot J_n^j \equiv \frac{\partial q^m}{\partial x^j} \cdot \frac{\partial x^j}{\partial q^n} = \frac{\partial q^m}{\partial q^n} = \delta_n^m = \begin{cases} 1 & \text{if } m = n \\ 0 & \text{if } m \neq n \end{cases}$$

$$\begin{pmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{pmatrix} \begin{pmatrix} \cos \theta & \sin \theta \\ -\frac{\sin \theta}{r} & \frac{\cos \theta}{r} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

Using differential chain-rules for coordinate transformations

Polar coordinate example of Generalized Curvilinear Coordinates (GCC)

*Getting the GCC ready for mechanics: Generalized **velocity and Jacobian Lemma 1***

 *Getting the GCC ready for mechanics: Generalized **acceleration and Lemma 2***

Getting the GCC ready for mechanics (2nd part)

Generalized *acceleration* relations are a little more complicated (It's curved coords, after all!)

First apply $\frac{d}{dt}$ to velocity \dot{x}^j and use product rule: $\frac{d}{dt}(u \cdot v) = \frac{du}{dt} \cdot v + u \cdot \frac{dv}{dt}$

$$\ddot{x}^j \equiv \frac{d}{dt} \dot{x}^j = \frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \dot{q}^m \right) = \frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \right) \dot{q}^m + \frac{\partial x^j}{\partial q^m} \ddot{q}^m$$

Getting the GCC ready for mechanics (2nd part)

Generalized *acceleration* relations are a little more complicated (It's curved coords, after all!)

First apply $\frac{d}{dt}$ to velocity \dot{x}^j and use product rule: $\frac{d}{dt}(u \cdot v) = \frac{du}{dt} \cdot v + u \cdot \frac{dv}{dt}$

$$\ddot{x}^j \equiv \frac{d}{dt} \dot{x}^j = \frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \dot{q}^m \right) = \frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \right) \dot{q}^m + \frac{\partial x^j}{\partial q^m} \ddot{q}^m$$

Apply derivative chain sum to Jacobian.

$$\frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \right) = \frac{\partial}{\partial q^n} \left(\frac{\partial x^j}{\partial q^m} \right) \frac{dq^n}{dt} = \left(\frac{\partial^2 x^j}{\partial q^n \partial q^m} \right) \frac{dq^n}{dt}$$

Getting the GCC ready for mechanics (2nd part)

Generalized *acceleration* relations are a little more complicated (It's curved coords, after all!)

First apply $\frac{d}{dt}$ to velocity \dot{x}^j and use product rule: $\frac{d}{dt}(u \cdot v) = \frac{du}{dt} \cdot v + u \cdot \frac{dv}{dt}$

$$\ddot{x}^j \equiv \frac{d}{dt} \dot{x}^j = \frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \dot{q}^m \right) = \frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \right) \dot{q}^m + \frac{\partial x^j}{\partial q^m} \ddot{q}^m$$

(Not in text. Recall Lecture 9 p. 15-19)[†]

Apply derivative chain sum to Jacobian. Partial derivatives are reversible. $\partial_m \partial_n = \partial_n \partial_m$

$$\frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \right) = \frac{\partial}{\partial q^n} \left(\frac{\partial x^j}{\partial q^m} \right) \frac{dq^n}{dt} = \left(\frac{\partial^2 x^j}{\partial q^n \partial q^m} \right) \frac{dq^n}{dt} = \left(\frac{\partial^2 x^j}{\partial q^m \partial q^n} \right) \frac{dq^n}{dt} = \frac{\partial}{\partial q^m} \left(\frac{\partial x^j}{\partial q^n} \frac{dq^n}{dt} \right)$$

Getting the GCC ready for mechanics (2nd part)

Generalized *acceleration* relations are a little more complicated (It's curved coords, after all!)

First apply $\frac{d}{dt}$ to velocity \dot{x}^j and use product rule: $\frac{d}{dt}(u \cdot v) = \frac{du}{dt} \cdot v + u \cdot \frac{dv}{dt}$

$$\ddot{x}^j \equiv \frac{d}{dt} \dot{x}^j = \frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \dot{q}^m \right) = \frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \right) \dot{q}^m + \frac{\partial x^j}{\partial q^m} \ddot{q}^m$$

(Not in text. Recall Lecture 9 p. 15-19)[†]

Apply derivative chain sum to Jacobian. Partial derivatives are reversible. $\partial_m \partial_n = \partial_n \partial_m$

$$\begin{aligned} \frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \right) &= \frac{\partial}{\partial q^n} \left(\frac{\partial x^j}{\partial q^m} \right) \frac{dq^n}{dt} = \left(\frac{\partial^2 x^j}{\partial q^n \partial q^m} \right) \frac{dq^n}{dt} = \left(\frac{\partial^2 x^j}{\partial q^m \partial q^n} \right) \frac{dq^n}{dt} = \frac{\partial}{\partial q^m} \left(\frac{\partial x^j}{\partial q^n} \frac{dq^n}{dt} \right) \\ & \text{By chain-rule def. of CC velocity:} \qquad \qquad \qquad = \frac{\partial}{\partial q^m} (\dot{x}^j) \end{aligned}$$

Getting the GCC ready for mechanics (2nd part)

Generalized *acceleration* relations are a little more complicated (It's curved coords, after all!)

First apply $\frac{d}{dt}$ to velocity \dot{x}^j and use product rule: $\frac{d}{dt}(u \cdot v) = \frac{du}{dt} \cdot v + u \cdot \frac{dv}{dt}$

$$\ddot{x}^j \equiv \frac{d}{dt} \dot{x}^j = \frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \dot{q}^m \right) = \frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \right) \dot{q}^m + \frac{\partial x^j}{\partial q^m} \ddot{q}^m$$

(Not in text. Recall Lecture 9 p. 15-19)[†]

Apply derivative chain sum to Jacobian. Partial derivatives are reversible. $\partial_m \partial_n = \partial_n \partial_m$

$$\begin{aligned} \frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \right) &= \frac{\partial}{\partial q^n} \left(\frac{\partial x^j}{\partial q^m} \right) \frac{dq^n}{dt} = \left(\frac{\partial^2 x^j}{\partial q^n \partial q^m} \right) \frac{dq^n}{dt} = \left(\frac{\partial^2 x^j}{\partial q^m \partial q^n} \right) \frac{dq^n}{dt} = \frac{\partial}{\partial q^m} \left(\frac{\partial x^j}{\partial q^n} \frac{dq^n}{dt} \right) \\ &= \frac{\partial}{\partial q^m} (\dot{x}^j) \end{aligned}$$

By chain-rule def. of CC velocity:

This is the key “*lemma-2*” for setting up Lagrangian mechanics .

$$\frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \right) = \frac{\partial \dot{x}^j}{\partial q^m} \text{ lemma 2}$$

Getting the GCC ready for mechanics (2nd part)

Generalized *acceleration* relations are a little more complicated (It's curved coords, after all!)

First apply $\frac{d}{dt}$ to velocity \dot{x}^j and use product rule: $\frac{d}{dt}(u \cdot v) = \frac{du}{dt} \cdot v + u \cdot \frac{dv}{dt}$

$$\ddot{x}^j \equiv \frac{d}{dt} \dot{x}^j = \frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \dot{q}^m \right) = \frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \right) \dot{q}^m + \frac{\partial x^j}{\partial q^m} \ddot{q}^m$$

(Not in text. Recall Lecture 9 p. 15-19)[†]

Apply derivative chain sum to Jacobian. Partial derivatives are reversible. $\partial_m \partial_n = \partial_n \partial_m$

$$\begin{aligned} \frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \right) &= \frac{\partial}{\partial q^n} \left(\frac{\partial x^j}{\partial q^m} \right) \frac{dq^n}{dt} = \left(\frac{\partial^2 x^j}{\partial q^n \partial q^m} \right) \frac{dq^n}{dt} = \left(\frac{\partial^2 x^j}{\partial q^m \partial q^n} \right) \frac{dq^n}{dt} = \frac{\partial}{\partial q^m} \left(\frac{\partial x^j}{\partial q^n} \frac{dq^n}{dt} \right) \\ &= \frac{\partial}{\partial q^m} (\dot{x}^j) \end{aligned}$$

By chain-rule def. of CC velocity:

The “*lemma-1*” was in the GCC velocity analysis just before this one for acceleration.

This is the key “*lemma-2*” for setting up Lagrangian mechanics .

$$\frac{\partial \dot{x}^j}{\partial \dot{q}^m} = \frac{\partial x^j}{\partial q^m} \quad \text{lemma 1}$$

$$\frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \right) = \frac{\partial \dot{x}^j}{\partial q^m} \quad \text{lemma 2}$$

How to say Newton's "F=ma" in Generalized Curvilinear Coords.

- *Use Cartesian KE quadratic form $KE=T=1/2\mathbf{v}\cdot\mathbf{M}\cdot\mathbf{v}$ and $\mathbf{F}=\mathbf{M}\cdot\mathbf{a}$ to get GCC force*
- Lagrange GCC trickery gives Lagrange force equations*
- Lagrange GCC trickery gives Lagrange potential equations (Lagrange 1 and 2)*

Deriving GCC mechanics from Cartesian Coord. (CC) Newton I-II

Start with stuff we know...(sort of)

Multidimensional CC version of kinetic energy $\frac{1}{2} \mathbf{v} \cdot \mathbf{M} \cdot \mathbf{v}$

$$T = \frac{1}{2} M_{jk} v^j v^k = \frac{1}{2} M_{jk} \dot{x}^j \dot{x}^k \quad \text{where: } M_{jk} \text{ are CC inertia constants}$$

Multidimensional CC version of Newt-II ($\mathbf{F} = \mathbf{M} \cdot \mathbf{a}$) using M_{jk} *constants*

$$f_j = M_{jk} a^k = M_{jk} \ddot{x}^k$$

Deriving GCC mechanics from Cartesian Coord. (CC) Newton I-II

Start with stuff we know...(sort of)

Multidimensional CC version of kinetic energy $\frac{1}{2} \mathbf{v} \cdot \mathbf{M} \cdot \mathbf{v}$

$$T = \frac{1}{2} M_{jk} v^j v^k = \frac{1}{2} M_{jk} \dot{x}^j \dot{x}^k \quad \text{where: } M_{jk} \text{ are inertia constants that are symmetric: } M_{jk} = M_{kj}$$

Multidimensional CC version of Newt-II ($\mathbf{F} = \mathbf{M} \cdot \mathbf{a}$) using M_{jk} constants

$$f_j = M_{jk} a^k = M_{jk} \ddot{x}^k$$

Multidimensional CC version of work-energy differential ($dW = \mathbf{F} \cdot d\mathbf{x}$). *Insert GCC differentials dq^m*

$$dW = f_j dx^j = f_j \left(\frac{\partial x^j}{\partial q^m} dq^m \right) = M_{jk} \ddot{x}^k \left(\frac{\partial x^j}{\partial q^m} dq^m \right)$$

(It's time to bring in the queer q^m !)

Deriving GCC mechanics from Cartesian Coord. (CC) Newton I-II

Start with stuff we know...(sort of)

Multidimensional CC version of kinetic energy $\frac{1}{2} \mathbf{v} \cdot \mathbf{M} \cdot \mathbf{v}$

$$T = \frac{1}{2} M_{jk} v^j v^k = \frac{1}{2} M_{jk} \dot{x}^j \dot{x}^k \quad \text{where: } M_{jk} \text{ are inertia constants that are symmetric: } M_{jk} = M_{kj}$$

Multidimensional CC version of Newt-II ($\mathbf{F} = \mathbf{M} \cdot \mathbf{a}$) using M_{jk} constants

$$f_j = M_{jk} a^k = M_{jk} \ddot{x}^k$$

Multidimensional CC version of work-energy differential ($dW = \mathbf{F} \cdot d\mathbf{x}$). Insert GCC differentials dq^m

$$dW = f_j dx^j = f_j \left(\frac{\partial x^j}{\partial q^m} dq^m \right) = M_{jk} \ddot{x}^k \left(\frac{\partial x^j}{\partial q^m} dq^m \right) \quad \text{(It's time to bring in the queer } q^m \text{ !)}$$

dq^m are independent so dq^m -sum is true term-by-term.

$$dW = f_j dx^j = F_m dq^m = f_j \frac{\partial x^j}{\partial q^m} dq^m = M_{jk} \ddot{x}^k \frac{\partial x^j}{\partial q^m} dq^m$$

Deriving GCC mechanics from Cartesian Coord. (CC) Newton I-II

Start with stuff we know...(sort of)

Multidimensional CC version of kinetic energy $\frac{1}{2} \mathbf{v} \cdot \mathbf{M} \cdot \mathbf{v}$

$$T = \frac{1}{2} M_{jk} v^j v^k = \frac{1}{2} M_{jk} \dot{x}^j \dot{x}^k \quad \text{where: } M_{jk} \text{ are inertia constants}$$

Multidimensional CC version of Newt-II ($\mathbf{F} = \mathbf{M} \cdot \mathbf{a}$) using M_{jk} constants

$$f_j = M_{jk} a^k = M_{jk} \ddot{x}^k$$

Multidimensional CC version of work-energy differential ($dW = \mathbf{F} \cdot d\mathbf{x}$). *Insert GCC differentials dq^m*

$$dW = f_j dx^j = f_j \left(\frac{\partial x^j}{\partial q^m} dq^m \right) = M_{jk} \ddot{x}^k \left(\frac{\partial x^j}{\partial q^m} dq^m \right) \quad (\text{It's time to bring in the queer } q^m !)$$

dq^m are independent so dq^m -sum is true term-by-term. (Still holds if all dq^m are zero but one.)

$$dW = f_j dx^j = F_m dq^m = f_j \frac{\partial x^j}{\partial q^m} dq^m = M_{jk} \ddot{x}^k \frac{\partial x^j}{\partial q^m} dq^m \Rightarrow F_m = f_j \frac{\partial x^j}{\partial q^m} = M_{jk} \ddot{x}^k \frac{\partial x^j}{\partial q^m}$$

Deriving GCC mechanics from Cartesian Coord. (CC) Newton I-II

Start with stuff we know...(sort of)

Multidimensional CC version of kinetic energy $\frac{1}{2} \mathbf{v} \cdot \mathbf{M} \cdot \mathbf{v}$

$$T = \frac{1}{2} M_{jk} v^j v^k = \frac{1}{2} M_{jk} \dot{x}^j \dot{x}^k \quad \text{where: } M_{jk} \text{ are inertia constants}$$

Multidimensional CC version of Newt-II ($\mathbf{F} = \mathbf{M} \cdot \mathbf{a}$) using M_{jk} constants

$$f_j = M_{jk} a^k = M_{jk} \ddot{x}^k$$

Multidimensional CC version of work-energy differential ($dW = \mathbf{F} \cdot d\mathbf{x}$). *Insert GCC differentials dq^m*

$$dW = f_j dx^j = f_j \left(\frac{\partial x^j}{\partial q^m} dq^m \right) = M_{jk} \ddot{x}^k \left(\frac{\partial x^j}{\partial q^m} dq^m \right) \quad (\text{It's time to bring in the queer } q^m !)$$

dq^m are independent so dq^m -sum is true term-by-term. (Still holds if all dq^m are zero but one.)

$$dW = f_j dx^j = F_m dq^m = f_j \frac{\partial x^j}{\partial q^m} dq^m = M_{jk} \ddot{x}^k \frac{\partial x^j}{\partial q^m} dq^m \Rightarrow F_m = f_j \frac{\partial x^j}{\partial q^m} = M_{jk} \ddot{x}^k \frac{\partial x^j}{\partial q^m}$$

Here *generalized GCC force component F_m* is defined:

$$\text{where: } F_m = f_j \frac{\partial x^j}{\partial q^m} = M_{jk} \ddot{x}^k \frac{\partial x^j}{\partial q^m}$$

How to say Newton's "F=ma" in Generalized Curvilinear Coords.

Use Cartesian KE quadratic form $KE=T=1/2\mathbf{v}\cdot\mathbf{M}\cdot\mathbf{v}$ and $\mathbf{F}=\mathbf{M}\cdot\mathbf{a}$ to get GCC force

 *Lagrange GCC trickery gives Lagrange force equations*

Lagrange GCC trickery gives Lagrange potential equations (Lagrange 1 and 2)

Now Lagrange GCC trickery begins

Obvious stuff...(sort of, if you've looked at it for a century!)

Lagrange's clever end game: First set $A = M_{jk} \dot{x}^k$ and $B = \frac{\partial x^j}{\partial q^m}$ with calc. formula: $\ddot{A}B = \frac{d}{dt}(\dot{A}B) - \dot{A}\dot{B}$

$$F_m = f_j \frac{\partial x^j}{\partial q^m} = M_{jk} \ddot{x}^k \frac{\partial x^j}{\partial q^m} = \frac{d}{dt} \left(M_{jk} \dot{x}^k \frac{\partial x^j}{\partial q^m} \right) - M_{jk} \dot{x}^k \frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \right)$$

Now Lagrange GCC trickery begins

Obvious stuff...(sort of, if you've looked at it for a century!)

Lagrange's clever end game: First set $A = M_{jk} \ddot{x}^k$ and $B = \frac{\partial x^j}{\partial q^m}$ with calc. formula: $\left[\ddot{A}B = \frac{d}{dt}(\dot{A}B) - \dot{A}\dot{B} \right]$

$$F_m = f_j \frac{\partial x^j}{\partial q^m} = M_{jk} \ddot{x}^k \frac{\partial x^j}{\partial q^m} = \frac{d}{dt} \left(M_{jk} \dot{x}^k \frac{\partial x^j}{\partial q^m} \right) - M_{jk} \dot{x}^k \frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \right)$$

Cartesian M_{jk}
must be constant
for this to work

(Bye, Bye relativistic mechanics or QM!)

Now Lagrange GCC trickery begins

Obvious stuff...(sort of, if you've looked at it for a century!)

Lagrange's clever end game: First set $A = M_{jk} \ddot{x}^k$ and $B = \frac{\partial x^j}{\partial q^m}$ with calc. formula: $\left[\ddot{A}B = \frac{d}{dt}(\dot{A}B) - \dot{A}\dot{B} \right]$

$$F_m = f_j \frac{\partial x^j}{\partial q^m} = M_{jk} \ddot{x}^k \frac{\partial x^j}{\partial q^m} = \frac{d}{dt} \left(M_{jk} \dot{x}^k \frac{\partial x^j}{\partial q^m} \right) - M_{jk} \dot{x}^k \frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \right)$$

$\ddot{A}B$ (red arrows) points to $M_{jk} \ddot{x}^k \frac{\partial x^j}{\partial q^m}$
 $(\dot{A}B)$ (red arrows) points to $M_{jk} \dot{x}^k \frac{\partial x^j}{\partial q^m}$
 $\dot{A}\dot{B}$ (red arrows) points to $M_{jk} \dot{x}^k \frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \right)$

Then convert ∂x^j to $\partial \dot{x}^j$ by *Lemma 1* and *Lemma 2* on 2nd term.

$$F_m = \frac{d}{dt} \left(M_{jk} \dot{x}^k \frac{\partial \dot{x}^j}{\partial \dot{q}^m} \right) - M_{jk} \dot{x}^k \left(\frac{\partial \dot{x}^j}{\partial q^m} \right)$$

(Red arrow) points from $\frac{\partial x^j}{\partial q^m}$ to $\frac{\partial \dot{x}^j}{\partial \dot{q}^m}$
 (Green arrow) points from $\frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \right)$ to $\left(\frac{\partial \dot{x}^j}{\partial q^m} \right)$

Cartesian M_{jk}
must be constant
for this to work

(Bye, Bye relativistic mechanics or QM!)

Now Lagrange GCC trickery begins

Obvious stuff...(sort of, if you've looked at it for a century!)

Lagrange's clever end game: First set $A = M_{jk} \ddot{x}^k$ and $B = \frac{\partial x^j}{\partial q^m}$ with calc. formula: $\left[\ddot{A}B = \frac{d}{dt}(\dot{A}B) - \dot{A}\dot{B} \right]$

$$F_m = f_j \frac{\partial x^j}{\partial q^m} = M_{jk} \ddot{x}^k \frac{\partial x^j}{\partial q^m} = \frac{d}{dt} \left(M_{jk} \dot{x}^k \frac{\partial x^j}{\partial q^m} \right) - M_{jk} \dot{x}^k \frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \right)$$

Then convert ∂x^j to $\partial \dot{x}^j$ by *Lemma 1* and *Lemma 2* on 2nd term.

$$F_m = \frac{d}{dt} \left(M_{jk} \dot{x}^k \frac{\partial \dot{x}^j}{\partial \dot{q}^m} \right) - M_{jk} \dot{x}^k \left(\frac{\partial \dot{x}^j}{\partial q^m} \right)$$

Simplify using: $\left[M_{ij} v^i \frac{\partial v^j}{\partial q} = M_{ij} \frac{\partial}{\partial q} \frac{v^i v^j}{2} \right]$ where q may be \dot{q}^m or q^m

$$F_m = \frac{d}{dt} \frac{\partial}{\partial \dot{q}^m} \left(\frac{M_{jk} \dot{x}^k \dot{x}^j}{2} \right) - \frac{\partial}{\partial q^m} \left(\frac{M_{jk} \dot{x}^k \dot{x}^j}{2} \right)$$

Cartesian M_{jk}
must be constant
for this to work

(Bye, Bye relativistic mechanics or QM!)

Now Lagrange GCC trickery begins

Obvious stuff...(sort of, if you've looked at it for a century!)

Lagrange's clever end game: First set $A = M_{jk} \ddot{x}^k$ and $B = \frac{\partial x^j}{\partial q^m}$ with calc. formula: $\left[\ddot{A}B = \frac{d}{dt}(\dot{A}B) - \dot{A}\dot{B} \right]$

$$F_m = f_j \frac{\partial x^j}{\partial q^m} = M_{jk} \ddot{x}^k \frac{\partial x^j}{\partial q^m} = \frac{d}{dt} \left(M_{jk} \dot{x}^k \frac{\partial x^j}{\partial q^m} \right) - M_{jk} \dot{x}^k \frac{d}{dt} \left(\frac{\partial x^j}{\partial q^m} \right)$$

Then convert ∂x^j to $\partial \dot{x}^j$ by *Lemma 1* and *Lemma 2* on 2nd term.

$$F_m = \frac{d}{dt} \left(M_{jk} \dot{x}^k \frac{\partial \dot{x}^j}{\partial \dot{q}^m} \right) - M_{jk} \dot{x}^k \left(\frac{\partial \dot{x}^j}{\partial q^m} \right)$$

Simplify using: $\left[M_{ij} v^i \frac{\partial v^j}{\partial q} = M_{ij} \frac{\partial}{\partial q} \frac{v^i v^j}{2} \right]$ where q may be \dot{q}^m or q^m

$$F_m = \frac{d}{dt} \frac{\partial}{\partial \dot{q}^m} \left(\frac{M_{jk} \dot{x}^k \dot{x}^j}{2} \right) - \frac{\partial}{\partial q^m} \left(\frac{M_{jk} \dot{x}^k \dot{x}^j}{2} \right)$$

The result is *Lagrange's GCC force equation* in terms of *kinetic energy* $T = \frac{1}{2} M_{jk} \dot{x}^j \dot{x}^k$

$$F_m = \frac{d}{dt} \frac{\partial T}{\partial \dot{q}^m} - \frac{\partial T}{\partial q^m} \quad \text{or:} \quad \mathbf{F} = \frac{d}{dt} \frac{\partial T}{\partial \mathbf{v}} - \frac{\partial T}{\partial \mathbf{r}}$$

How to say Newton's "F=ma" in Generalized Curvilinear Coords.

Use Cartesian KE quadratic form $KE=T=1/2\mathbf{v}\cdot\mathbf{M}\cdot\mathbf{v}$ and $\mathbf{F}=\mathbf{M}\cdot\mathbf{a}$ to get GCC force

Lagrange GCC trickery gives Lagrange force equations

 *Lagrange GCC trickery gives Lagrange potential equations (Lagrange 1 and 2)*

But, Lagrange GCC trickery is not yet done...

(Still another trick-up-the-sleeve!)

If the force is conservative it's a gradient $\mathbf{F} = -\nabla U$

In GCC: $F_m = -\frac{\partial U}{\partial q^m}$

$$F_m = -\frac{\partial U}{\partial q^m} = \frac{d}{dt} \frac{\partial T}{\partial \dot{q}^m} - \frac{\partial T}{\partial q^m}$$

But, Lagrange GCC trickery is not yet done...

(Still another trick-up-the-sleeve!)

If the force is conservative it's a gradient $\mathbf{F} = -\nabla U$

In GCC: $F_m = -\frac{\partial U}{\partial q^m}$

$$F_m = -\frac{\partial U}{\partial q^m} = \frac{d}{dt} \frac{\partial T}{\partial \dot{q}^m} - \frac{\partial T}{\partial q^m}$$

Becomes *Lagrange's GCC potential equation* with a new definition for the *Lagrangian: $L=T-U$* .

$$0 = \frac{d}{dt} \frac{\partial L}{\partial \dot{q}^m} - \frac{\partial L}{\partial q^m}$$

$$L(\dot{q}^m, q^m) = T(\dot{q}^m, q^m) - U(q^m)$$

This trick requires: $\frac{\partial U}{\partial \dot{q}^m} \equiv 0$ *U(r) has
NO explicit
velocity
dependence!*

But, Lagrange GCC trickery is not yet done...

(Still another trick-up-the-sleeve!)

If the force is conservative it's a gradient $\mathbf{F} = -\nabla U$

In GCC: $F_m = -\frac{\partial U}{\partial q^m}$

$$F_m = -\frac{\partial U}{\partial q^m} = \frac{d}{dt} \frac{\partial T}{\partial \dot{q}^m} - \frac{\partial T}{\partial q^m}$$

Becomes *Lagrange's GCC potential equation* with a new definition for the *Lagrangian: $L=T-U$* .

$$0 = \frac{d}{dt} \frac{\partial L}{\partial \dot{q}^m} - \frac{\partial L}{\partial q^m}$$

$$L(\dot{q}^m, q^m) = T(\dot{q}^m, q^m) - U(q^m)$$

This trick requires: $\frac{\partial U}{\partial \dot{q}^m} \equiv 0$

*U(r) has
NO explicit
velocity
dependence!*

*Lagrange's 1st GCC equation
(Defining GCC momentum)*

$$p_m = \frac{\partial L}{\partial \dot{q}^m}$$

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{q}^m} = \frac{\partial L}{\partial q^m}$$

Recall:

$$\mathbf{p} = \frac{\partial L}{\partial \mathbf{v}}$$

*Lagrange's 2nd GCC equation
(Change of GCC momentum)*

$$\frac{dp_m}{dt} \equiv \dot{p}_m = \frac{\partial L}{\partial q^m}$$

But, Lagrange GCC trickery is not yet done...

(Still another trick-up-the-sleeve!)

If the force is conservative it's a gradient $\mathbf{F} = -\nabla U$

In GCC: $F_m = -\frac{\partial U}{\partial q^m}$

$$F_m = -\frac{\partial U}{\partial q^m} = \frac{d}{dt} \frac{\partial T}{\partial \dot{q}^m} - \frac{\partial T}{\partial q^m}$$

Becomes *Lagrange's GCC potential equation* with a new definition for the *Lagrangian*: $L=T-U$.

$$0 = \frac{d}{dt} \frac{\partial L}{\partial \dot{q}^m} - \frac{\partial L}{\partial q^m}$$

$$L(\dot{q}^m, q^m) = T(\dot{q}^m, q^m) - U(q^m)$$

This trick requires: $\frac{\partial U}{\partial \dot{q}^m} \equiv 0$

*U(r) has
NO explicit
velocity
dependence!*

*If L has no
explicit q^m
dependence
then:*

$$\dot{p}_m = 0$$

or :

$$p_m = \text{const.}$$

*Lagrange's 1st GCC equation
(Defining GCC momentum)*

$$p_m = \frac{\partial L}{\partial \dot{q}^m}$$

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{q}^m} = \frac{\partial L}{\partial q^m}$$

Recall :

$$\mathbf{p} = \frac{\partial L}{\partial \mathbf{v}}$$

*Lagrange's 2nd GCC equation
(Change of GCC momentum)*

$$\frac{dp_m}{dt} \equiv \dot{p}_m = \frac{\partial L}{\partial q^m}$$

GCC Cells, base vectors, and metric tensors

→ *Polar coordinate examples: Covariant \mathbf{E}_m vs. Contravariant \mathbf{E}^m
Covariant g_{mn} vs. Invariant δ_m^n vs. Contravariant g^{mn}*

A dual set of *quasi-unit vectors* show up in Jacobian J and Kajobian K.

J-Columns are *covariant vectors* $\{\mathbf{E}_1 = \mathbf{E}_r \quad \mathbf{E}_2 = \mathbf{E}_\phi\}$

K-Rows are *contravariant vectors* $\{\mathbf{E}^1 = \mathbf{E}^r \quad \mathbf{E}^2 = \mathbf{E}^\phi\}$

$$\langle J \rangle = \begin{pmatrix} \frac{\partial x^1}{\partial q^1} & \frac{\partial x^1}{\partial q^2} \\ \frac{\partial x^2}{\partial q^1} & \frac{\partial x^2}{\partial q^2} \end{pmatrix} = \begin{pmatrix} \frac{\partial x}{\partial r} = \cos \phi & \frac{\partial x}{\partial \phi} = -r \sin \phi \\ \frac{\partial y}{\partial r} = \sin \phi & \frac{\partial y}{\partial \phi} = r \cos \phi \end{pmatrix}$$

$\uparrow \mathbf{E}_1 \quad \uparrow \mathbf{E}_2 \qquad \uparrow \mathbf{E}_r \qquad \uparrow \mathbf{E}_\phi$


$$\langle K \rangle = \langle J^{-1} \rangle = \begin{pmatrix} \frac{\partial r}{\partial x} = \cos \phi & \frac{\partial r}{\partial y} = \sin \phi \\ \frac{\partial \phi}{\partial x} = \frac{-\sin \phi}{r} & \frac{\partial \phi}{\partial y} = \frac{\cos \phi}{r} \end{pmatrix} \leftarrow \begin{matrix} \mathbf{E}^r = \mathbf{E}^1 \\ \mathbf{E}^\phi = \mathbf{E}^2 \end{matrix}$$

Inverse polar definition:

$r^2 = x^2 + y^2$ and $\phi = \text{atan2}(y, x)$

Derived from polar definition: $x = r \cos \phi$ and $y = r \sin \phi$

(a) Polar coordinate bases



A dual set of *quasi-unit vectors* show up in Jacobian J and Kajobian K.

J-Columns are *covariant vectors* $\{\mathbf{E}_1=\mathbf{E}_r \ \mathbf{E}_2=\mathbf{E}_\phi\}$

K-Rows are *contravariant vectors* $\{\mathbf{E}^1=\mathbf{E}^r \ \mathbf{E}^2=\mathbf{E}^\phi\}$

$$\langle J \rangle = \begin{pmatrix} \frac{\partial x^1}{\partial q^1} & \frac{\partial x^1}{\partial q^2} \\ \frac{\partial x^2}{\partial q^1} & \frac{\partial x^2}{\partial q^2} \end{pmatrix} = \begin{pmatrix} \frac{\partial x}{\partial r} = \cos \phi & \frac{\partial x}{\partial \phi} = -r \sin \phi \\ \frac{\partial y}{\partial r} = \sin \phi & \frac{\partial y}{\partial \phi} = r \cos \phi \end{pmatrix}$$

$\uparrow \mathbf{E}_1 \quad \uparrow \mathbf{E}_2 \qquad \uparrow \mathbf{E}_r \quad \uparrow \mathbf{E}_\phi$

$$\langle K \rangle = \langle J^{-1} \rangle = \begin{pmatrix} \frac{\partial r}{\partial x} = \cos \phi & \frac{\partial r}{\partial y} = \sin \phi \\ \frac{\partial \phi}{\partial x} = \frac{-\sin \phi}{r} & \frac{\partial \phi}{\partial y} = \frac{\cos \phi}{r} \end{pmatrix}$$


$\leftarrow \mathbf{E}^r = \mathbf{E}^1$
 $\leftarrow \mathbf{E}^\phi = \mathbf{E}^2$

Inverse polar definition:


$r^2=x^2+y^2$ and $\phi = \text{atan2}(y,x)$

Derived from polar definition: $x=r \cos \phi$ and $y=r \sin \phi$

(a) Polar coordinate bases




(b) Covariant bases $\{\mathbf{E}_1 \ \mathbf{E}_2\}$
(Tangent)



NOTE: These are 2D drawings!
No 3D perspective

(c) Contravariant bases $\{\mathbf{E}^1 \ \mathbf{E}^2\}$
(Normal)




Comparison: Covariant $\mathbf{E}_m = \frac{\partial \mathbf{r}}{\partial q^m}$ vs. Contravariant $\mathbf{E}^m = \frac{\partial q^m}{\partial \mathbf{r}} = \nabla_{q^m}$

Covariant bases $\{\mathbf{E}_1, \mathbf{E}_2\}$ match ^{geometric unit} cell walls
 (Tangent)

$$\Delta \mathbf{r} = \mathbf{E}_1 \Delta q^1 + \mathbf{E}_2 \Delta q^2$$


is based on chain rule: $d\mathbf{r} = \frac{\partial \mathbf{r}}{\partial q^1} dq^1 + \frac{\partial \mathbf{r}}{\partial q^2} dq^2 = \mathbf{E}_1 dq^1 + \mathbf{E}_2 dq^2$



NOTE: These
 are 2D drawings!
No 3D perspective

Comparison: Covariant $\mathbf{E}_m = \frac{\partial \mathbf{r}}{\partial q^m}$ vs. Contravariant $\mathbf{E}^m = \frac{\partial q^m}{\partial \mathbf{r}} = \nabla_{q^m}$

Covariant bases $\{\mathbf{E}_1, \mathbf{E}_2\}$ match ^{geometric unit} cell walls
 (Tangent)



$$\Delta \mathbf{r} = \mathbf{E}_1 \Delta q^1 + \mathbf{E}_2 \Delta q^2$$


is based on chain rule: $d\mathbf{r} = \frac{\partial \mathbf{r}}{\partial q^1} dq^1 + \frac{\partial \mathbf{r}}{\partial q^2} dq^2 = \mathbf{E}_1 dq^1 + \mathbf{E}_2 dq^2$

\mathbf{E}_1 follows *tangent* to $q^2 = \text{const.}$...
 since only q^1 varies in $\frac{\partial \mathbf{r}}{\partial q^1}$
 while q^2, q^3, \dots remain constant

NOTE: These
 are 2D drawings!
No 3D perspective

Comparison: Covariant $\mathbf{E}_m = \frac{\partial \mathbf{r}}{\partial q^m}$ vs. Contravariant $\mathbf{E}^m = \frac{\partial q^m}{\partial \mathbf{r}} = \nabla_{q^m}$

Covariant bases $\{\mathbf{E}_1, \mathbf{E}_2\}$ match ^{geometric unit} cell walls
 (Tangent)



$$\Delta \mathbf{r} = \mathbf{E}_1 \Delta q^1 + \mathbf{E}_2 \Delta q^2 \quad \text{is based on chain rule: } d\mathbf{r} = \frac{\partial \mathbf{r}}{\partial q^1} dq^1 + \frac{\partial \mathbf{r}}{\partial q^2} dq^2 = \mathbf{E}_1 dq^1 + \mathbf{E}_2 dq^2$$

\mathbf{E}_1 follows *tangent* to $q^2 = \text{const.}$...
 since only q^1 varies in $\frac{\partial \mathbf{r}}{\partial q^1}$
 while q^2, q^3, \dots remain constant


\mathbf{E}_m are convenient bases for *extensive* quantities like distance and velocity.

$$\mathbf{V} = V^1 \mathbf{E}_1 + V^2 \mathbf{E}_2 = V^1 \frac{\partial \mathbf{r}}{\partial q^1} + V^2 \frac{\partial \mathbf{r}}{\partial q^2}$$

NOTE: These
 are 2D drawings!
No 3D perspective

Comparison: Covariant $\mathbf{E}_m = \frac{\partial \mathbf{r}}{\partial q^m}$ vs. Contravariant $\mathbf{E}^m = \frac{\partial q^m}{\partial \mathbf{r}} = \nabla q^m$

Covariant bases $\{\mathbf{E}_1, \mathbf{E}_2\}$ match ^{geometric unit} cell walls
 (Tangent)




\mathbf{E}_1 follows *tangent* to $q^2 = \text{const.}$...
 since only q^1 varies in $\frac{\partial \mathbf{r}}{\partial q^1}$
 while q^2, q^3, \dots remain constant

\mathbf{E}_m are convenient bases for *extensive* quantities like distance and velocity.

$$\mathbf{V} = V^1 \mathbf{E}_1 + V^2 \mathbf{E}_2 = V^1 \frac{\partial \mathbf{r}}{\partial q^1} + V^2 \frac{\partial \mathbf{r}}{\partial q^2}$$

Contravariant $\{\mathbf{E}^1, \mathbf{E}^2\}$ match reciprocal cells

(Normal)




NOTE: These are 2D drawings!
No 3D perspective

\mathbf{E}^1 is *normal* to $q^2 = \text{const.}$ since **gradient** of q^1 is vector sum $\nabla q^1 = \left(\begin{array}{c} \frac{\partial q^1}{\partial x} \\ \frac{\partial q^1}{\partial y} \end{array} \right)$ of all its partial derivatives

Comparison: Covariant $\mathbf{E}_m = \frac{\partial \mathbf{r}}{\partial q^m}$ vs. Contravariant $\mathbf{E}^m = \frac{\partial q^m}{\partial \mathbf{r}} = \nabla q^m$

geometric unit

Covariant bases $\{\mathbf{E}_1 \mathbf{E}_2\}$ match cell walls
(Tangent)




\mathbf{E}_m are convenient bases for extensive quantities like distance and velocity.

$$\mathbf{V} = V^1 \mathbf{E}_1 + V^2 \mathbf{E}_2 = V^1 \frac{\partial \mathbf{r}}{\partial q^1} + V^2 \frac{\partial \mathbf{r}}{\partial q^2}$$

Contravariant $\{\mathbf{E}^1 \mathbf{E}^2\}$ match reciprocal cells

(Normal)



NOTE: These are 2D drawings! No 3D perspective


\mathbf{E}^1 is normal to $q^2 = \text{const.}$ since gradient of q^1 is vector sum $\nabla q^1 = \left(\begin{array}{c} \frac{\partial q^1}{\partial x} \\ \frac{\partial q^1}{\partial y} \end{array} \right)$ of all its partial derivatives

\mathbf{E}^m are convenient bases for intensive quantities like force and momentum.

$$\mathbf{F} = F_1 \mathbf{E}^1 + F_2 \mathbf{E}^2 = F_1 \frac{\partial q^1}{\partial \mathbf{r}} + F_2 \frac{\partial q^2}{\partial \mathbf{r}} = F_1 \nabla q^1 + F_2 \nabla q^2$$

Comparison: Covariant $\mathbf{E}_m = \frac{\partial \mathbf{r}}{\partial q^m}$ vs. Contravariant $\mathbf{E}^n = \frac{\partial q^n}{\partial \mathbf{r}} = \nabla q^n$

Covariant bases $\{\mathbf{E}_1, \mathbf{E}_2\}$ match ^{geometric unit} cell walls
(Tangent)



\mathbf{E}_m are convenient bases for *extensive* quantities like distance and velocity.


$$\mathbf{V} = V^1 \mathbf{E}_1 + V^2 \mathbf{E}_2 = V^1 \frac{\partial \mathbf{r}}{\partial q^1} + V^2 \frac{\partial \mathbf{r}}{\partial q^2}$$

Co-Contr dot products $\mathbf{E}_m \cdot \mathbf{E}^n$ are *orthonormal*:

$$\mathbf{E}_m \cdot \mathbf{E}^n = \frac{\partial \mathbf{r}}{\partial q^m} \cdot \frac{\partial q^n}{\partial \mathbf{r}} = \delta_m^n$$

Contravariant $\{\mathbf{E}^1, \mathbf{E}^2\}$ match reciprocal cells

(Normal)



\mathbf{E}^1 is *normal* to $q^1 = \text{const.}$ since **gradient** of q^1 is vector sum $\nabla q^1 =$

$$\left(\begin{array}{c} \frac{\partial q^1}{\partial x} \\ \frac{\partial q^1}{\partial y} \end{array} \right)$$

\mathbf{E}^m are convenient bases for *intensive* quantities like force and momentum.

$$\mathbf{F} = F_1 \mathbf{E}^1 + F_2 \mathbf{E}^2 = F_1 \frac{\partial q^1}{\partial \mathbf{r}} + F_2 \frac{\partial q^2}{\partial \mathbf{r}} = F_1 \nabla q^1 + F_2 \nabla q^2$$

GCC Cells, base vectors, and metric tensors

Polar coordinate examples: Covariant \mathbf{E}_m vs. Contravariant \mathbf{E}^m
 *Covariant g_{mn} vs. Invariant δ_m^n vs. Contravariant g^{mn}*

Covariant g_{mn} vs. Invariant δ_m^n vs. Contravariant g^{mn}

$$\mathbf{E}_m \cdot \mathbf{E}_n = \frac{\partial \mathbf{r}}{\partial q^m} \cdot \frac{\partial \mathbf{r}}{\partial q^n} \equiv g_{mn}$$

$$\mathbf{E}_m \cdot \mathbf{E}^n = \frac{\partial \mathbf{r}}{\partial q^m} \cdot \frac{\partial q^n}{\partial \mathbf{r}} = \delta_m^n$$

$$\mathbf{E}^m \cdot \mathbf{E}^n = \frac{\partial q^m}{\partial \mathbf{r}} \cdot \frac{\partial q^n}{\partial \mathbf{r}} \equiv g^{mn}$$

Covariant
metric tensor

g_{mn}

Invariant
Kronecker unit tensor

$$\delta_m^n \equiv \begin{cases} 1 & \text{if } m = n \\ 0 & \text{if } m \neq n \end{cases}$$

Contravariant
metric tensor

g^{mn}

Covariant g_{mn} vs. Invariant δ_m^n vs. Contravariant g^{mn}

$$\mathbf{E}_m \cdot \mathbf{E}_n = \frac{\partial \mathbf{r}}{\partial q^m} \cdot \frac{\partial \mathbf{r}}{\partial q^n} \equiv g_{mn}$$

$$\mathbf{E}_m \cdot \mathbf{E}^n = \frac{\partial \mathbf{r}}{\partial q^m} \cdot \frac{\partial q^n}{\partial \mathbf{r}} = \delta_m^n$$

$$\mathbf{E}^m \cdot \mathbf{E}^n = \frac{\partial q^m}{\partial \mathbf{r}} \cdot \frac{\partial q^n}{\partial \mathbf{r}} \equiv g^{mn}$$

Covariant
metric tensor

g_{mn}

Invariant
Kronecker unit tensor

$$\delta_m^n \equiv \begin{cases} 1 & \text{if } m = n \\ 0 & \text{if } m \neq n \end{cases}$$

Contravariant
metric tensor

g^{mn}

Polar coordinate examples (again):

$$\langle J \rangle = \begin{pmatrix} \frac{\partial x^1}{\partial q^1} & \frac{\partial x^1}{\partial q^2} \\ \frac{\partial x^2}{\partial q^1} & \frac{\partial x^2}{\partial q^2} \end{pmatrix} = \begin{pmatrix} \frac{\partial x}{\partial r} = \cos \phi & \frac{\partial x}{\partial \phi} = -r \sin \phi \\ \frac{\partial y}{\partial r} = \sin \phi & \frac{\partial y}{\partial \phi} = r \cos \phi \end{pmatrix}$$

$\uparrow \mathbf{E}_1$ $\uparrow \mathbf{E}_2$ $\uparrow \mathbf{E}_r$ $\uparrow \mathbf{E}_\phi$

$$\langle K \rangle = \langle J^{-1} \rangle = \begin{pmatrix} \frac{\partial r}{\partial x} = \cos \phi & \frac{\partial r}{\partial y} = \sin \phi \\ \frac{\partial \phi}{\partial x} = \frac{-\sin \phi}{r} & \frac{\partial \phi}{\partial y} = \frac{\cos \phi}{r} \end{pmatrix} \leftarrow \mathbf{E}^r = \mathbf{E}^1$$

$$\leftarrow \mathbf{E}^\phi = \mathbf{E}^2$$

Covariant g_{mn} vs. Invariant δ_m^n vs. Contravariant g^{mn}

$$\mathbf{E}_m \cdot \mathbf{E}_n = \frac{\partial \mathbf{r}}{\partial q^m} \cdot \frac{\partial \mathbf{r}}{\partial q^n} \equiv g_{mn}$$

$$\mathbf{E}_m \cdot \mathbf{E}^n = \frac{\partial \mathbf{r}}{\partial q^m} \cdot \frac{\partial q^n}{\partial \mathbf{r}} = \delta_m^n$$

$$\mathbf{E}^m \cdot \mathbf{E}^n = \frac{\partial q^m}{\partial \mathbf{r}} \cdot \frac{\partial q^n}{\partial \mathbf{r}} \equiv g^{mn}$$

Covariant
metric tensor

g_{mn}

Invariant
Kronecker unit tensor

$$\delta_m^n \equiv \begin{cases} 1 & \text{if } m = n \\ 0 & \text{if } m \neq n \end{cases}$$

Contravariant
metric tensor

g^{mn}

Polar coordinate examples (again):

$$\langle J \rangle = \begin{pmatrix} \frac{\partial x^1}{\partial q^1} & \frac{\partial x^1}{\partial q^2} \\ \frac{\partial x^2}{\partial q^1} & \frac{\partial x^2}{\partial q^2} \end{pmatrix} = \begin{pmatrix} \frac{\partial x}{\partial r} = \cos \phi & \frac{\partial x}{\partial \phi} = -r \sin \phi \\ \frac{\partial y}{\partial r} = \sin \phi & \frac{\partial y}{\partial \phi} = r \cos \phi \end{pmatrix} \quad \langle K \rangle = \langle J^{-1} \rangle = \begin{pmatrix} \frac{\partial r}{\partial x} = \cos \phi & \frac{\partial r}{\partial y} = \sin \phi \\ \frac{\partial \phi}{\partial x} = \frac{-\sin \phi}{r} & \frac{\partial \phi}{\partial y} = \frac{\cos \phi}{r} \end{pmatrix}$$

$\uparrow \mathbf{E}_1 \quad \uparrow \mathbf{E}_2 \quad \quad \uparrow \mathbf{E}_r \quad \quad \uparrow \mathbf{E}_\phi \quad \quad \quad \leftarrow \mathbf{E}^r = \mathbf{E}^1$
 $\quad \leftarrow \mathbf{E}^\phi = \mathbf{E}^2$

Covariant g_{mn}

Invariant δ_m^n

Contravariant g^{mn}

$$\begin{pmatrix} g_{rr} & g_{r\phi} \\ g_{\phi r} & g_{\phi\phi} \end{pmatrix} = \begin{pmatrix} \mathbf{E}_r \cdot \mathbf{E}_r & \mathbf{E}_r \cdot \mathbf{E}_\phi \\ \mathbf{E}_\phi \cdot \mathbf{E}_r & \mathbf{E}_\phi \cdot \mathbf{E}_\phi \end{pmatrix} \quad \begin{pmatrix} \delta_r^r & \delta_r^\phi \\ \delta_\phi^r & \delta_\phi^\phi \end{pmatrix} = \begin{pmatrix} \mathbf{E}_r \cdot \mathbf{E}^r & \mathbf{E}_r \cdot \mathbf{E}^\phi \\ \mathbf{E}_\phi \cdot \mathbf{E}^r & \mathbf{E}_\phi \cdot \mathbf{E}^\phi \end{pmatrix} \quad \begin{pmatrix} g^{rr} & g^{r\phi} \\ g^{\phi r} & g^{\phi\phi} \end{pmatrix} = \begin{pmatrix} \mathbf{E}^r \cdot \mathbf{E}^r & \mathbf{E}^r \cdot \mathbf{E}^\phi \\ \mathbf{E}^\phi \cdot \mathbf{E}^r & \mathbf{E}^\phi \cdot \mathbf{E}^\phi \end{pmatrix}$$

$$= \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix} \quad = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad = \begin{pmatrix} 1 & 0 \\ 0 & 1/r^2 \end{pmatrix}$$

Lagrange prefers Covariant g_{mn} with Contravariant velocity \dot{q}^m



GCC Lagrangian definition

GCC “canonical” momentum p_m definition

GCC “canonical” force F_m definition

Coriolis “fictitious” forces (... and weather effects)

Lagrange prefers Covariant g_{mn} with Contravariant velocity

Lagrangian $L=KE-U$ is supposed to be explicit function of velocity.

$$L(\mathbf{v}) = \frac{1}{2} M \mathbf{v} \cdot \mathbf{v} - U = \frac{1}{2} M \dot{\mathbf{r}} \cdot \dot{\mathbf{r}} - U = \frac{1}{2} M (\mathbf{E}_m \dot{q}^m) \cdot (\mathbf{E}_n \dot{q}^n) - U = \frac{1}{2} M (g_{mn} \dot{q}^m \dot{q}^n) - U = L(\dot{q})$$

Lagrange prefers Covariant g_{mn} with Contravariant velocity

Lagrangian KE-U is supposed to be explicit function of velocity.

$$L(\mathbf{v}) = \frac{1}{2} M \mathbf{v} \cdot \mathbf{v} - U = \frac{1}{2} M \dot{\mathbf{r}} \cdot \dot{\mathbf{r}} - U = \frac{1}{2} M (\mathbf{E}_m \dot{q}^m) \cdot (\mathbf{E}_n \dot{q}^n) - U = \frac{1}{2} M (g_{mn} \dot{q}^m \dot{q}^n) - U = L(\dot{q})$$

Use polar coordinate Covariant g_{mn} metric (page 53)

$$\begin{pmatrix} g_{rr} & g_{r\phi} \\ g_{\phi r} & g_{\phi\phi} \end{pmatrix} = \begin{pmatrix} \mathbf{E}_r \cdot \mathbf{E}_r & \mathbf{E}_r \cdot \mathbf{E}_\phi \\ \mathbf{E}_\phi \cdot \mathbf{E}_r & \mathbf{E}_\phi \cdot \mathbf{E}_\phi \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$$

Lagrange prefers Covariant g_{mn} with Contravariant velocity

Lagrangian KE-U is supposed to be explicit function of velocity.

$$L(\mathbf{v}) = \frac{1}{2} M \mathbf{v} \cdot \mathbf{v} - U = \frac{1}{2} M \dot{\mathbf{r}} \cdot \dot{\mathbf{r}} - U = \frac{1}{2} M (\mathbf{E}_m \dot{q}^m) \cdot (\mathbf{E}_n \dot{q}^n) - U = \frac{1}{2} M (g_{mn} \dot{q}^m \dot{q}^n) - U = L(\dot{q})$$

Use polar coordinate Covariant g_{mn} metric (page 53)

$$\begin{pmatrix} g_{rr} & g_{r\phi} \\ g_{\phi r} & g_{\phi\phi} \end{pmatrix} = \begin{pmatrix} \mathbf{E}_r \cdot \mathbf{E}_r & \mathbf{E}_r \cdot \mathbf{E}_\phi \\ \mathbf{E}_\phi \cdot \mathbf{E}_r & \mathbf{E}_\phi \cdot \mathbf{E}_\phi \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$$

This gives polar GCC form (Actually it's an OCC or Orthogonal Curvilinear Coordinate form)

$$L(\dot{r}, \dot{\phi}) = \frac{1}{2} M (g_{rr} \dot{r}^2 + g_{\phi\phi} \dot{\phi}^2) - U(r, \phi) = \frac{1}{2} M (1 \cdot \dot{r}^2 + r^2 \dot{\phi}^2) - U(r, \phi)$$

Lagrange prefers Covariant g_{mn} with Contravariant velocity \dot{q}^m

GCC Lagrangian definition

 *GCC “canonical” momentum p_m definition*

GCC “canonical” force F_m definition

Coriolis “fictitious” forces (... and weather effects)

Lagrange prefers Covariant g_{mn} with Contravariant velocity

Lagrangian KE-U is supposed to be explicit function of velocity.

$$L(\mathbf{v}) = \frac{1}{2} M \mathbf{v} \cdot \mathbf{v} - U = \frac{1}{2} M \dot{\mathbf{r}} \cdot \dot{\mathbf{r}} - U = \frac{1}{2} M (\mathbf{E}_m \dot{q}^m) \cdot (\mathbf{E}_n \dot{q}^n) - U = \frac{1}{2} M (g_{mn} \dot{q}^m \dot{q}^n) - U = L(\dot{q})$$

Use polar coordinate Covariant g_{mn} metric (page 53)

$$\begin{pmatrix} g_{rr} & g_{r\phi} \\ g_{\phi r} & g_{\phi\phi} \end{pmatrix} = \begin{pmatrix} \mathbf{E}_r \cdot \mathbf{E}_r & \mathbf{E}_r \cdot \mathbf{E}_\phi \\ \mathbf{E}_\phi \cdot \mathbf{E}_r & \mathbf{E}_\phi \cdot \mathbf{E}_\phi \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$$

This gives polar GCC form (Actually it's an OCC or Orthogonal Curvilinear Coordinate form)

$$L(\dot{r}, \dot{\phi}) = \frac{1}{2} M (g_{rr} \dot{r}^2 + g_{\phi\phi} \dot{\phi}^2) - U(r, \phi) = \frac{1}{2} M (1 \cdot \dot{r}^2 + r^2 \dot{\phi}^2) - U(r, \phi)$$

(From preceding page)

Lagrange prefers Covariant g_{mn} with Contravariant velocity

Lagrangian KE-U is supposed to be explicit function of velocity.

$$L(\mathbf{v}) = \frac{1}{2} M \mathbf{v} \cdot \mathbf{v} - U = \frac{1}{2} M \dot{\mathbf{r}} \cdot \dot{\mathbf{r}} - U = \frac{1}{2} M (\mathbf{E}_m \dot{q}^m) \cdot (\mathbf{E}_n \dot{q}^n) - U = \frac{1}{2} M (g_{mn} \dot{q}^m \dot{q}^n) - U = L(\dot{q})$$

Use polar coordinate Covariant g_{mn} metric (page 53)

$$\begin{pmatrix} g_{rr} & g_{r\phi} \\ g_{\phi r} & g_{\phi\phi} \end{pmatrix} = \begin{pmatrix} \mathbf{E}_r \cdot \mathbf{E}_r & \mathbf{E}_r \cdot \mathbf{E}_\phi \\ \mathbf{E}_\phi \cdot \mathbf{E}_r & \mathbf{E}_\phi \cdot \mathbf{E}_\phi \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$$

This gives polar GCC form (Actually it's an OCC or Orthogonal Curvilinear Coordinate form)

$$L(\dot{r}, \dot{\phi}) = \frac{1}{2} M (g_{rr} \dot{r}^2 + g_{\phi\phi} \dot{\phi}^2) - U(r, \phi) = \frac{1}{2} M (1 \cdot \dot{r}^2 + r^2 \dot{\phi}^2) - U(r, \phi)$$

GCC Lagrange equations follow. 1st L-equation is momentum p_m definition for each coordinate q^m :

$$p_r = \frac{\partial L}{\partial \dot{r}} = M g_{rr} \dot{r} = M \dot{r}$$

Nothing too surprising;
radial momentum p_r has the
usual linear $M \cdot v$ form

Lagrange prefers Covariant g_{mn} with Contravariant velocity

Lagrangian KE-U is supposed to be explicit function of velocity.

$$L(\mathbf{v}) = \frac{1}{2} M \mathbf{v} \cdot \mathbf{v} - U = \frac{1}{2} M \dot{\mathbf{r}} \cdot \dot{\mathbf{r}} - U = \frac{1}{2} M (\mathbf{E}_m \dot{q}^m) \cdot (\mathbf{E}_n \dot{q}^n) - U = \frac{1}{2} M (g_{mn} \dot{q}^m \dot{q}^n) - U = L(\dot{q})$$

Use polar coordinate Covariant g_{mn} metric (page 53)

$$\begin{pmatrix} g_{rr} & g_{r\phi} \\ g_{\phi r} & g_{\phi\phi} \end{pmatrix} = \begin{pmatrix} \mathbf{E}_r \cdot \mathbf{E}_r & \mathbf{E}_r \cdot \mathbf{E}_\phi \\ \mathbf{E}_\phi \cdot \mathbf{E}_r & \mathbf{E}_\phi \cdot \mathbf{E}_\phi \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$$

This gives polar GCC form (Actually it's an OCC or Orthogonal Curvilinear Coordinate form)

$$L(\dot{r}, \dot{\phi}) = \frac{1}{2} M (g_{rr} \dot{r}^2 + g_{\phi\phi} \dot{\phi}^2) - U(r, \phi) = \frac{1}{2} M (1 \cdot \dot{r}^2 + r^2 \dot{\phi}^2) - U(r, \phi)$$

GCC Lagrange equations follow. 1st L-equation is momentum p_m definition for each coordinate q^m :

$$p_r = \frac{\partial L}{\partial \dot{r}} = M g_{rr} \dot{r} = M \dot{r}$$

Nothing too surprising;
radial momentum p_r has the
usual linear $M \cdot v$ form

$$p_\phi = \frac{\partial L}{\partial \dot{\phi}} = M g_{\phi\phi} \dot{\phi} = M r^2 \dot{\phi}$$

Wow! $g_{\phi\phi}$ gives moment-of-inertia
factor Mr^2 automatically for the
angular momentum $p_\phi = Mr^2 \omega$.

Lagrange's 1st GCC equation
(Defining GCC momentum)

$$p_m = \frac{\partial L}{\partial \dot{q}^m}$$

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{q}^m} = \frac{\partial L}{\partial q^m}$$

Recall:

$$\mathbf{p} = \frac{\partial L}{\partial \mathbf{v}}$$


Lagrange's 2nd GCC equation
(Change of GCC momentum)

$$\frac{dp_m}{dt} \equiv \dot{p}_m = \frac{\partial L}{\partial q^m}$$

Lagrange prefers Covariant g_{mn} with Contravariant velocity \dot{q}^m

GCC Lagrangian definition

GCC “canonical” momentum p_m definition

 *GCC “canonical” force F_m definition*

Coriolis “fictitious” forces (... and weather effects)

Lagrange prefers Covariant g_{mn} with Contravariant velocity

Lagrangian KE-U is supposed to be explicit function of velocity.

$$L(\mathbf{v}) = \frac{1}{2} M \mathbf{v} \cdot \mathbf{v} - U = \frac{1}{2} M \dot{\mathbf{r}} \cdot \dot{\mathbf{r}} - U = \frac{1}{2} M (\mathbf{E}_m \dot{q}^m) \cdot (\mathbf{E}_n \dot{q}^n) - U = \frac{1}{2} M (g_{mn} \dot{q}^m \dot{q}^n) - U = L(\dot{q})$$

Use polar coordinate Covariant g_{mn} metric (page 53)

$$\begin{pmatrix} g_{rr} & g_{r\phi} \\ g_{\phi r} & g_{\phi\phi} \end{pmatrix} = \begin{pmatrix} \mathbf{E}_r \cdot \mathbf{E}_r & \mathbf{E}_r \cdot \mathbf{E}_\phi \\ \mathbf{E}_\phi \cdot \mathbf{E}_r & \mathbf{E}_\phi \cdot \mathbf{E}_\phi \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$$

This gives polar GCC form (Actually it's an OCC or Orthogonal Curvilinear Coordinate form)

$$L(\dot{r}, \dot{\phi}) = \frac{1}{2} M (g_{rr} \dot{r}^2 + g_{\phi\phi} \dot{\phi}^2) - U(r, \phi) = \frac{1}{2} M (1 \cdot \dot{r}^2 + r^2 \dot{\phi}^2) - U(r, \phi)$$

GCC Lagrange equations follow. 1st L-equation is momentum p_m definition for each coordinate q^m :

$$p_r = \frac{\partial L}{\partial \dot{r}} = M g_{rr} \dot{r} = M \dot{r}$$

Nothing too surprising;
radial momentum p_r has the
usual linear $M \cdot v$ form

$$p_\phi = \frac{\partial L}{\partial \dot{\phi}} = M g_{\phi\phi} \dot{\phi} = M r^2 \dot{\phi}$$

Wow! $g_{\phi\phi}$ gives moment-of-inertia
factor Mr^2 automatically for the
angular momentum $p_\phi = Mr^2 \omega$.

(From preceding page)

Lagrange prefers Covariant g_{mn} with Contravariant velocity

Lagrangian KE-U is supposed to be explicit function of velocity.

$$L(\mathbf{v}) = \frac{1}{2} M \mathbf{v} \cdot \mathbf{v} - U = \frac{1}{2} M \dot{\mathbf{r}} \cdot \dot{\mathbf{r}} - U = \frac{1}{2} M (\mathbf{E}_m \dot{q}^m) \cdot (\mathbf{E}_n \dot{q}^n) - U = \frac{1}{2} M (g_{mn} \dot{q}^m \dot{q}^n) - U = L(\dot{q})$$

Use polar coordinate Covariant g_{mn} metric (page 53)

$$\begin{pmatrix} g_{rr} & g_{r\phi} \\ g_{\phi r} & g_{\phi\phi} \end{pmatrix} = \begin{pmatrix} \mathbf{E}_r \cdot \mathbf{E}_r & \mathbf{E}_r \cdot \mathbf{E}_\phi \\ \mathbf{E}_\phi \cdot \mathbf{E}_r & \mathbf{E}_\phi \cdot \mathbf{E}_\phi \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$$

This gives polar GCC form (Actually it's an OCC or Orthogonal Curvilinear Coordinate form)

$$L(\dot{r}, \dot{\phi}) = \frac{1}{2} M (g_{rr} \dot{r}^2 + g_{\phi\phi} \dot{\phi}^2) - U(r, \phi) = \frac{1}{2} M (1 \cdot \dot{r}^2 + r^2 \dot{\phi}^2) - U(r, \phi)$$

GCC Lagrange equations follow. 1st L-equation is momentum p_m definition for each coordinate q^m :

$$p_r = \frac{\partial L}{\partial \dot{r}} = M g_{rr} \dot{r} = M \dot{r}$$

Nothing too surprising;
radial momentum p_r has the
usual linear $M \cdot v$ form

$$p_\phi = \frac{\partial L}{\partial \dot{\phi}} = M g_{\phi\phi} \dot{\phi} = M r^2 \dot{\phi}$$

Wow! $g_{\phi\phi}$ gives moment-of-inertia
factor $M r^2$ automatically for the
angular momentum $p_\phi = M r^2 \omega$.

2nd L-equation involves total time derivative \dot{p}_m for each momentum p_m :

$$\dot{p}_r = \frac{\partial L}{\partial r} = \frac{M}{2} \frac{\partial g_{\phi\phi}}{\partial r} \dot{\phi}^2 - \frac{\partial U}{\partial r} = M r \dot{\phi}^2 - \frac{\partial U}{\partial r}$$

Centrifugal
force $M r \omega^2$

$$\dot{p}_\phi = \frac{\partial L}{\partial \phi} = 0 - \frac{\partial U}{\partial \phi}$$

Angular momentum p_ϕ is **conserved** if
potential U has no explicit ϕ -dependence

Lagrange's 1st GCC equation
(Defining GCC momentum)

$$p_m = \frac{\partial L}{\partial \dot{q}^m}$$

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{q}^m} = \frac{\partial L}{\partial q^m}$$

Recall:

$$\mathbf{p} = \frac{\partial L}{\partial \mathbf{v}}$$

Lagrange's 2nd GCC equation
(Change of GCC momentum)

$$\frac{dp_m}{dt} \equiv \dot{p}_m = \frac{\partial L}{\partial q^m}$$

Lagrange prefers Covariant g_{mn} with Contravariant velocity

Lagrangian KE-U is supposed to be explicit function of velocity.

$$L(\mathbf{v}) = \frac{1}{2} M \mathbf{v} \cdot \mathbf{v} - U = \frac{1}{2} M \dot{\mathbf{r}} \cdot \dot{\mathbf{r}} - U = \frac{1}{2} M (\mathbf{E}_m \dot{q}^m) \cdot (\mathbf{E}_n \dot{q}^n) - U = \frac{1}{2} M (g_{mn} \dot{q}^m \dot{q}^n) - U = L(\dot{q})$$

Use polar coordinate Covariant g_{mn} metric (page 53)

$$\begin{pmatrix} g_{rr} & g_{r\phi} \\ g_{\phi r} & g_{\phi\phi} \end{pmatrix} = \begin{pmatrix} \mathbf{E}_r \cdot \mathbf{E}_r & \mathbf{E}_r \cdot \mathbf{E}_\phi \\ \mathbf{E}_\phi \cdot \mathbf{E}_r & \mathbf{E}_\phi \cdot \mathbf{E}_\phi \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$$

This gives polar GCC form (Actually it's an OCC or Orthogonal Curvilinear Coordinate form)

$$L(\dot{r}, \dot{\phi}) = \frac{1}{2} M (g_{rr} \dot{r}^2 + g_{\phi\phi} \dot{\phi}^2) - U(r, \phi) = \frac{1}{2} M (1 \cdot \dot{r}^2 + r^2 \dot{\phi}^2) - U(r, \phi)$$

GCC Lagrange equations follow. 1st L-equation is momentum p_m definition for each coordinate q^m :

$$p_r = \frac{\partial L}{\partial \dot{r}} = M g_{rr} \dot{r} = M \dot{r}$$

Nothing too surprising;
radial momentum p_r has the
usual linear $M \cdot v$ form

$$p_\phi = \frac{\partial L}{\partial \dot{\phi}} = M g_{\phi\phi} \dot{\phi} = M r^2 \dot{\phi}$$

Wow! $g_{\phi\phi}$ gives moment-of-inertia
factor $M r^2$ automatically for the
angular momentum $p_\phi = M r^2 \omega$.

2nd L-equation involves total time derivative \dot{p}_m for each momentum p_m :

$$\dot{p}_r = \frac{\partial L}{\partial r} = \frac{M}{2} \frac{\partial g_{\phi\phi}}{\partial r} \dot{\phi}^2 - \frac{\partial U}{\partial r} = M r \dot{\phi}^2 - \frac{\partial U}{\partial r}$$

Centrifugal
force $M r \omega^2$

$$\dot{p}_\phi = \frac{\partial L}{\partial \phi} = 0 - \frac{\partial U}{\partial \phi}$$

Angular momentum p_ϕ is **conserved** if
potential U has no explicit ϕ -dependence

Find \dot{p}_m directly from 1st L-equation: $\dot{p}_m \equiv \frac{dp_m}{dt} = \frac{d}{dt} M (g_{mn} \dot{q}^n) = M (\dot{g}_{mn} \dot{q}^n + g_{mn} \ddot{q}^n)$ Equate it to \dot{p}_m in 2nd L-equation:

Lagrange prefers Covariant g_{mn} with Contravariant velocity \dot{q}^m

GCC Lagrangian definition

GCC “canonical” momentum p_m definition

 *GCC “canonical” force F_m definition*

Coriolis “fictitious” forces (... and weather effects)

Lagrange prefers Covariant g_{mn} with Contravariant velocity

Lagrangian KE-U is supposed to be explicit function of velocity.

$$L(\mathbf{v}) = \frac{1}{2} M \mathbf{v} \cdot \mathbf{v} - U = \frac{1}{2} M \dot{\mathbf{r}} \cdot \dot{\mathbf{r}} - U = \frac{1}{2} M (\mathbf{E}_m \dot{q}^m) \cdot (\mathbf{E}_n \dot{q}^n) - U = \frac{1}{2} M (g_{mn} \dot{q}^m \dot{q}^n) - U = L(\dot{q})$$

Use polar coordinate Covariant g_{mn} metric (page 53)

$$\begin{pmatrix} g_{rr} & g_{r\phi} \\ g_{\phi r} & g_{\phi\phi} \end{pmatrix} = \begin{pmatrix} \mathbf{E}_r \cdot \mathbf{E}_r & \mathbf{E}_r \cdot \mathbf{E}_\phi \\ \mathbf{E}_\phi \cdot \mathbf{E}_r & \mathbf{E}_\phi \cdot \mathbf{E}_\phi \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$$

This gives polar GCC form (Actually it's an OCC or Orthogonal Curvilinear Coordinate form)

$$L(\dot{r}, \dot{\phi}) = \frac{1}{2} M (g_{rr} \dot{r}^2 + g_{\phi\phi} \dot{\phi}^2) - U(r, \phi) = \frac{1}{2} M (1 \cdot \dot{r}^2 + r^2 \dot{\phi}^2) - U(r, \phi)$$

GCC Lagrange equations follow. 1st L-equation is momentum p_m definition for each coordinate q^m :

$$p_r = \frac{\partial L}{\partial \dot{r}} = M g_{rr} \dot{r} = M \dot{r}$$

Nothing too surprising;
radial momentum p_r has the
usual linear $M \cdot v$ form

$$p_\phi = \frac{\partial L}{\partial \dot{\phi}} = M g_{\phi\phi} \dot{\phi} = M r^2 \dot{\phi}$$

Wow! $g_{\phi\phi}$ gives moment-of-inertia
factor Mr^2 automatically for the
angular momentum $p_\phi = Mr^2 \omega$.

2nd L-equation involves total time derivative \dot{p}_m for each momentum p_m :

$$\dot{p}_r = \frac{\partial L}{\partial r} = \frac{M}{2} \frac{\partial g_{\phi\phi}}{\partial r} \dot{\phi}^2 - \frac{\partial U}{\partial r} = M r \dot{\phi}^2 - \frac{\partial U}{\partial r}$$

Centrifugal
force $Mr\omega^2$

$$\dot{p}_\phi = \frac{\partial L}{\partial \phi} = 0 - \frac{\partial U}{\partial \phi}$$

Angular momentum p_ϕ is **conserved** if
potential U has no explicit ϕ -dependence

Find \dot{p}_m directly from 1st L-equation: $\dot{p}_m \equiv \frac{dp_m}{dt} = \frac{d}{dt} M (g_{mn} \dot{q}^n) = M (\dot{g}_{mn} \dot{q}^n + g_{mn} \ddot{q}^n)$ Equate it to \dot{p}_m in 2nd L-equation:

(From preceding page)

Lagrange prefers Covariant g_{mn} with Contravariant velocity

Lagrangian KE-U is supposed to be explicit function of velocity.

$$L(\mathbf{v}) = \frac{1}{2} M \mathbf{v} \cdot \mathbf{v} - U = \frac{1}{2} M \dot{\mathbf{r}} \cdot \dot{\mathbf{r}} - U = \frac{1}{2} M (\mathbf{E}_m \dot{q}^m) \cdot (\mathbf{E}_n \dot{q}^n) - U = \frac{1}{2} M (g_{mn} \dot{q}^m \dot{q}^n) - U = L(\dot{q})$$

Use polar coordinate Covariant g_{mn} metric (page 53)

$$\begin{pmatrix} g_{rr} & g_{r\phi} \\ g_{\phi r} & g_{\phi\phi} \end{pmatrix} = \begin{pmatrix} \mathbf{E}_r \cdot \mathbf{E}_r & \mathbf{E}_r \cdot \mathbf{E}_\phi \\ \mathbf{E}_\phi \cdot \mathbf{E}_r & \mathbf{E}_\phi \cdot \mathbf{E}_\phi \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$$

This gives polar GCC form (Actually it's an OCC or Orthogonal Curvilinear Coordinate form)

$$L(\dot{r}, \dot{\phi}) = \frac{1}{2} M (g_{rr} \dot{r}^2 + g_{\phi\phi} \dot{\phi}^2) - U(r, \phi) = \frac{1}{2} M (1 \cdot \dot{r}^2 + r^2 \dot{\phi}^2) - U(r, \phi)$$

GCC Lagrange equations follow. 1st L-equation is momentum p_m definition for each coordinate q^m :

$$p_r = \frac{\partial L}{\partial \dot{r}} = M g_{rr} \dot{r} = M \dot{r}$$

Nothing too surprising;
radial momentum p_r has the
usual linear $M \cdot v$ form

$$p_\phi = \frac{\partial L}{\partial \dot{\phi}} = M g_{\phi\phi} \dot{\phi} = M r^2 \dot{\phi}$$

Wow! $g_{\phi\phi}$ gives moment-of-inertia
factor $M r^2$ automatically for the
angular momentum $p_\phi = M r^2 \omega$.

2nd L-equation involves total time derivative \dot{p}_m for each momentum p_m :

$$\dot{p}_r = \frac{\partial L}{\partial r} = \frac{M}{2} \frac{\partial g_{\phi\phi}}{\partial r} \dot{\phi}^2 - \frac{\partial U}{\partial r} = M r \dot{\phi}^2 - \frac{\partial U}{\partial r}$$

Centrifugal
force $M r \omega^2$

$$\dot{p}_\phi = \frac{\partial L}{\partial \phi} = 0 - \frac{\partial U}{\partial \phi}$$

Angular momentum p_ϕ is **conserved** if
potential U has no explicit ϕ -dependence

Find \dot{p}_m directly from 1st L-equation: $\dot{p}_m \equiv \frac{dp_m}{dt} = \frac{d}{dt} M (g_{mn} \dot{q}^n) = M (\dot{g}_{mn} \dot{q}^n + g_{mn} \ddot{q}^n)$ Equate it to \dot{p}_m in 2nd L-equation:

$$\begin{aligned} \dot{p}_r &\equiv \frac{dp_r}{dt} = M \ddot{r} \\ &= M r \dot{\phi}^2 - \frac{\partial U}{\partial r} \end{aligned}$$

Centrifugal (center-fleeing) force
equals total
Centripetal (center-pulling) force

Lagrange prefers Covariant g_{mn} with Contravariant velocity

Lagrangian KE-U is supposed to be explicit function of velocity.

$$L(\mathbf{v}) = \frac{1}{2} M \mathbf{v} \cdot \mathbf{v} - U = \frac{1}{2} M \dot{\mathbf{r}} \cdot \dot{\mathbf{r}} - U = \frac{1}{2} M (\mathbf{E}_m \dot{q}^m) \cdot (\mathbf{E}_n \dot{q}^n) - U = \frac{1}{2} M (g_{mn} \dot{q}^m \dot{q}^n) - U = L(\dot{q})$$

Use polar coordinate Covariant g_{mn} metric (page 53)

$$\begin{pmatrix} g_{rr} & g_{r\phi} \\ g_{\phi r} & g_{\phi\phi} \end{pmatrix} = \begin{pmatrix} \mathbf{E}_r \cdot \mathbf{E}_r & \mathbf{E}_r \cdot \mathbf{E}_\phi \\ \mathbf{E}_\phi \cdot \mathbf{E}_r & \mathbf{E}_\phi \cdot \mathbf{E}_\phi \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$$

This gives polar GCC form (Actually it's an OCC or Orthogonal Curvilinear Coordinate form)

$$L(\dot{r}, \dot{\phi}) = \frac{1}{2} M (g_{rr} \dot{r}^2 + g_{\phi\phi} \dot{\phi}^2) - U(r, \phi) = \frac{1}{2} M (1 \cdot \dot{r}^2 + r^2 \dot{\phi}^2) - U(r, \phi)$$

GCC Lagrange equations follow. 1st L-equation is momentum p_m definition for each coordinate q^m :

$$p_r = \frac{\partial L}{\partial \dot{r}} = M g_{rr} \dot{r} = M \dot{r}$$

Nothing too surprising;
radial momentum p_r has the
usual linear $M \cdot v$ form

$$p_\phi = \frac{\partial L}{\partial \dot{\phi}} = M g_{\phi\phi} \dot{\phi} = M r^2 \dot{\phi}$$

Wow! $g_{\phi\phi}$ gives moment-of-inertia
factor $M r^2$ automatically for the
angular momentum $p_\phi = M r^2 \omega$.

2nd L-equation involves total time derivative \dot{p}_m for each momentum p_m :

$$\dot{p}_r = \frac{\partial L}{\partial r} = \frac{M}{2} \frac{\partial g_{\phi\phi}}{\partial r} \dot{\phi}^2 - \frac{\partial U}{\partial r} = M r \dot{\phi}^2 - \frac{\partial U}{\partial r}$$

Centrifugal
force $M r \omega^2$

$$\dot{p}_\phi = \frac{\partial L}{\partial \phi} = 0 - \frac{\partial U}{\partial \phi}$$

Angular momentum p_ϕ is **conserved** if
potential U has no explicit ϕ -dependence

Find \dot{p}_m directly from 1st L-equation: $\dot{p}_m \equiv \frac{dp_m}{dt} = \frac{d}{dt} M (g_{mn} \dot{q}^n) = M (\dot{g}_{mn} \dot{q}^n + g_{mn} \ddot{q}^n)$ Equate it to \dot{p}_m in 2nd L-equation:

$$\dot{p}_r \equiv \frac{dp_r}{dt} = M \ddot{r}$$

Centrifugal (center-fleeing) force
equals total
Centripetal (center-pulling) force

$$= M r \dot{\phi}^2 - \frac{\partial U}{\partial r}$$

$$\dot{p}_\phi \equiv \frac{dp_\phi}{dt} = 2 M r \dot{r} \dot{\phi} + M r^2 \ddot{\phi}$$

Torque relates to two distinct parts:
Coriolis and angular acceleration

$$= 0 - \frac{\partial U}{\partial \phi}$$

Angular momentum p_ϕ is **conserved** if
potential U has no explicit ϕ -dependence

Rewriting GCC Lagrange equations :

$$\dot{p}_r \equiv \frac{dp_r}{dt} = M \ddot{r}$$

Centrifugal (center-fleeing) force equals total

$$= M r \dot{\phi}^2 - \frac{\partial U}{\partial r}$$

Centripetal (center-pulling) force

$$\dot{p}_\phi \equiv \frac{dp_\phi}{dt} = 2Mr\dot{\phi} + Mr^2\ddot{\phi}$$

Torque relates to two distinct parts: Coriolis and angular acceleration

$$= 0 - \frac{\partial U}{\partial \phi}$$

Angular momentum p_ϕ is conserved if potential U has no explicit ϕ -dependence

Conventional forms


radial force: $M \ddot{r} = M r \dot{\phi}^2 - \frac{\partial U}{\partial r}$

angular force or torque: $Mr^2\ddot{\phi} = -2Mr\dot{\phi} - \frac{\partial U}{\partial \phi}$

Field-free ($U=0$)

radial acceleration: $\ddot{r} = r \dot{\phi}^2$

angular acceleration: $\ddot{\phi} = -2 \frac{\dot{r}\dot{\phi}}{r}$



Rewriting GCC Lagrange equations :

$$\dot{p}_r \equiv \frac{dp_r}{dt} = M \ddot{r}$$

Centrifugal (center-fleeing) force equals total

$$= M r \dot{\phi}^2 - \frac{\partial U}{\partial r}$$

Centripetal (center-pulling) force

$$\dot{p}_\phi \equiv \frac{dp_\phi}{dt} = 2Mr\dot{\phi} + Mr^2\ddot{\phi}$$

Torque relates to two distinct parts: Coriolis and angular acceleration

$$= 0 - \frac{\partial U}{\partial \phi}$$

Angular momentum p_ϕ is conserved if potential U has no explicit ϕ -dependence

Conventional forms


radial force: $M \ddot{r} = M r \dot{\phi}^2 - \frac{\partial U}{\partial r}$

angular force or torque: $Mr^2\ddot{\phi} = -2Mr\dot{\phi} - \frac{\partial U}{\partial \phi}$

Field-free ($U=0$)

radial acceleration: $\ddot{r} = r \dot{\phi}^2$

angular acceleration: $\ddot{\phi} = -2 \frac{\dot{r}\dot{\phi}}{r}$



Effect on Northern Hemisphere local weather

Cyclonic flow around lows

Rewriting GCC Lagrange equations :

$$\dot{p}_r \equiv \frac{dp_r}{dt} = M \ddot{r}$$

Centrifugal (center-fleeing) force equals total

$$= M r \dot{\phi}^2 - \frac{\partial U}{\partial r}$$

Centripetal (center-pulling) force

$$\dot{p}_\phi \equiv \frac{dp_\phi}{dt} = 2Mr\dot{\phi} + Mr^2\ddot{\phi}$$

Torque relates to two distinct parts: Coriolis and angular acceleration

$$= 0 - \frac{\partial U}{\partial \phi}$$

Angular momentum p_ϕ is conserved if potential U has no explicit ϕ -dependence

Conventional forms


radial force: $M \ddot{r} = M r \dot{\phi}^2 - \frac{\partial U}{\partial r}$

angular force or torque: $Mr^2\ddot{\phi} = -2Mr\dot{\phi} - \frac{\partial U}{\partial \phi}$


Field-free ($U=0$)

radial acceleration: $\ddot{r} = r \dot{\phi}^2$


angular acceleration: $\ddot{\phi} = -2 \frac{\dot{r}\dot{\phi}}{r}$



GOES-16 captured this geocolor image of Hurricane Irma approaching Anguilla at about 7:15 am (eastern), September 6, 2017. Irma's maximum sustained winds remain near 185 mph with higher gusts, making it a category 5 hurricane on the Saffir-Simpson Hurricane Wind Scale. According to the latest information from NOAA's National Hurricane Center (issued at 8:00 am eastern), Irma was located about 15 miles west-southwest of Anguilla and moving toward the west-northwest near 16 miles per hour.



Science News link



Saturn's north pole was dark when Cassini arrived in 2004. But as the seasons changed, light illuminated a bizarre six-sided swirl of gases at the pole (shown here in false color). The hexagon has been known since the 1980s. It is about 30,000 kilometers (18,600 miles) wide with a massive hurricane centered on the north pole.

JPL-CALTECH/NASA, SPACE SCIENCE INSTITUTE

Lecture 9 ends here
Thu 9/19/2017