


Assignment 12 - PHYS 5103-11/13/17-Due Mon. Nov. 21 CMwBang! Ch 4.1 thru Ch.4.4. and Lectures 20-24

Ex.1 The “standard” Lorentzian (Note: Review complex 2-pole potential $\phi(z)=1/z$ (10.42) in Unit 1-Ch. 10 Fig. 10.11.)

In physics literature, a standard Lorentzian function generally means a form $L(\Delta) = A/(\Delta^2 + A^2)$ with constant A . In the *Near-Resonant Approximation* (NRA is (4.2.18)) the $L(\Delta)$ or its derivative is an approximation to exact G -equations (4.2.15).

- (a) Use NRA (4.2.18) to reduce (4.2.15a-d) and identify a standard Lorentzian function of the detuning parameter $\Delta=\omega_s-\omega_0$.
- (b) Show that NRA for complex response $G=\text{Re } G + i\text{Im } G$ gives circular arcs in the complex $\omega=|\omega| e^{i\theta} = \Delta + i\Gamma$ plane for constant decay rate Γ and variable detuning or beat rate Δ . How does this circle deviate from what is almost a circle in Fig. 4.2.6? (Consider higher Γ values for which NRA breaks down such as Fig. 4.2.14.) Fixed Δ and varying Γ give what curve?
- (c) Do ruler-&-compass construction of NRA Lorentz-Green functions as in figures below for $b=1/2$ and $b=1/3$.

$\text{Re } G_{\omega_0}(\omega_s) = y = \frac{b}{x^2 + b^2}$, $\text{Im } G_{\omega_0}(\omega_s) = y = \frac{x}{x^2 + b^2}$, and $|G_{\omega_0}(\omega_s)|$. (See p. 60-62 of Lect. 20.)




Ex.2 Max and min G -values (Part (b-c) involves some algebra!)

Derive equations for the extreme values for the response function or function related to G as asked below.

For part (a) *only* use *Near-Resonant Approximation (NRA)*: See preceding **Ex.1**.

- (a1) Find values which give maxima for: $\text{Re } G_{\omega_0}(\omega_s)$, $\text{Im } G_{\omega_0}(\omega_s)$, and $|G_{\omega_0}(\omega_s)|$ assuming ω_0 is constant and ω_s varies.
- (a2) Find values which give maxima for: $\text{Re } G_{\omega_0}(\omega_s)$, $\text{Im } G_{\omega_0}(\omega_s)$, and $|G_{\omega_0}(\omega_s)|$ assuming ω_s is constant and ω_0 varies.
- (b) Do (a) for *exact* $G_{\omega_0}(\omega_s)$. Exact plots by calculator help check these answers.
- (c) Find exact value to maximize peak KE of responding oscillator. (1st show total $KE = \frac{1}{2} m \omega^2 A^2$ for oscillation of amplitude A .)



Ex.3 Coupled oscillation

Two identical mass $M=1\text{kg}$ blocks slide friction-free on a rod and are connected by springs $k_1=16\text{N}\cdot\text{m}^{-1}$ and $k_2=37\text{N}\cdot\text{m}^{-1}$ to ends of a box and coupled to each other by spring $k_{12}=36\text{N}\cdot\text{m}^{-1}$.

- (a) Write Lagrangian equations of motion and derive a \mathbf{K} -matrix form of them.
- (b) Solve for eigenmodes and eigenfrequencies of system and plot their directions on an X,Y-graph. Use spectral decomposition methods (Lect. 20 or Appendix 4.C) to derive eigensolution projectors and eigenvectors.
- (c) Given initial conditions $(X(0)=1, Y(0)=0)$, plot the resulting path in the XY-plane. Show algebraically that it is a parabola.
- (d) Use spectral decomposition (Lect. 20 or Appendix 4.C) to derive square-roots $\mathbf{H}=\sqrt{\mathbf{K}}$. (How many square-roots does \mathbf{K} have?)

Ex.4 $U(2)$ view of coupled oscillation

- (a) Rewrite the spring \mathbf{K} -matrix for **Ex.3** into an \mathbf{H} -matrix where $\mathbf{K}=\mathbf{H}^2$ as in (4.4.8).
- (b) Give the resulting \mathbf{H} -matrix as an (A,B,C,D) combination of $\mathbf{1}$, σ_A , σ_B , and σ_C as in (4.4.9). (++) root of \mathbf{K} results for \mathbf{H} .)
- (c) Sketch the resulting Ω -whirl vector or “crank” in real 3D (A,B,C) -space as in (4.4.10).
- (d) For $(X(0)=1, Y(0)=0)$ find initial \mathbf{S} -state (“spin”)vector in (A,B,C) -space as in (4.4.16). Show its evolution by Ω as in Fig. 4.4.2.
- (e) Plot \mathbf{H} -eigenvalues (ϵ_1, ϵ_2) as though they were energy levels and indicate transition rate $\Omega=\epsilon_1-\epsilon_2$ and mean rate $\omega=(\epsilon_1+\epsilon_2)/2$.