

# Lecture 12

## Tue. 10.03.2017

# Complex Variables, Series, and Field Coordinates I.

(Ch. 10 of Unit 1)

## 1. The Story of $e$ (A Tale of Great \$Interest\$)

How good are those power series?

Taylor-Maclaurin series, imaginary interest, and complex exponentials

Lecture 14 Tue. 10.15  
starts here

## 2. What good are complex exponentials?

Easy trig

Easy 2D vector analysis

Easy oscillator phase analysis

Easy rotation and “dot” or “cross” products

## 3. Easy 2D vector calculus

Easy 2D vector derivatives

Easy 2D source-free field theory

Easy 2D vector field-potential theory

## 4. Riemann-Cauchy relations (What's analytic? What's not?)

Easy 2D curvilinear coordinate discovery

Lect. 12

Easy 2D circulation and flux integrals

ends here

Easy 2D monopole, dipole, and  $2^n$ -pole analysis

Easy  $2^n$ -multipole field and potential expansion

Easy stereo-projection visualization

Cauchy integrals, Laurent-Maclaurin series

1. Complex numbers provide "automatic trigonometry"

2. Complex numbers add like vectors.

3. Complex exponentials  $Ae^{-i\omega t}$  track position and velocity using Phasor Clock.

4. Complex products provide 2D rotation operations.

5. Complex products provide 2D “dot”(•) and “cross”(x) products.

6. Complex derivative contains “divergence”(∇•F) and “curl”(∇x F) of 2D vector field

7. Invent source-free 2D vector fields [∇•F=0 and ∇x F=0]

8. Complex potential  $\phi$  contains “scalar”(F=∇Φ) and “vector”(F=∇xA) potentials

The **half-n'-half** results: (Riemann-Cauchy Derivative Relations)

9. Complex potentials define 2D Orthogonal Curvilinear Coordinates (OCC) of field

10. Complex integrals  $\int f(z)dz$  count 2D “circulation”(∫F•dr) and “flux”(∫Fxdr)

11. Complex integrals define 2D **monopole** fields and potentials

12. Complex derivatives give 2D dipole fields

Lecture 15 Thur. 10.17  
starts here

13. More derivatives give 2D  $2^N$ -pole fields...

14. ...and  $2^N$ -pole multipole expansions of fields and potentials...

15. ...and Laurent Series...

16. Mapping and non-analytic source analysis.

## *The Story of $e$ (A Tale of Great \$Interest\$)*

Simple *interest* at some rate  $r$  based on a 1 year period.

You gave a principal  $p(0)$  to the bank and some time  $t$  later they would pay you  $p(t)=(1+r\cdot t)p(0)$ .

\$1.00 at rate  $r=1$  (like Israel and Brazil that once had 100% interest.) gives \$2.00 at  $t=1$  year.

## *The Story of e (A Tale of Great \$Interest\$)*

Simple *interest* at some rate  $r$  based on a 1 year period.

You gave a principal  $p(0)$  to the bank and some time  $t$  later they would pay you  $p(t)=(1+r\cdot t)p(0)$ .

\$1.00 at rate  $r=1$  (like Israel and Brazil that once had 100% interest.) gives \$2.00 at  $t=1$  year.

*Semester compounded* interest gives  $p(\frac{t}{2})=(1+r\cdot\frac{t}{2})p(0)$  at the half-period  $\frac{t}{2}$  and then use  $p(\frac{t}{2})$  during the last half to figure final payment. Now \$1.00 at rate  $r=1$  earns \$2.25.

$$p^{\frac{1}{2}}(t) = (1+r\cdot\frac{t}{2})p(\frac{t}{2}) = (1+r\cdot\frac{t}{2})\cdot(1+r\cdot\frac{t}{2})p(0) = \frac{3}{2}\cdot\frac{3}{2}\cdot 1 = \frac{9}{4} = 2.25$$

## The Story of $e$ (A Tale of Great \$Interest\$)

Simple *interest* at some rate  $r$  based on a 1 year period.

You gave a principal  $p(0)$  to the bank and some time  $t$  later they would pay you  $p(t)=(1+r\cdot t)p(0)$ .

\$1.00 at rate  $r=1$  (like Israel and Brazil that once had 100% interest.) gives \$2.00 at  $t=1$  year.

*Semester compounded* interest gives  $p(\frac{t}{2})=(1+r\cdot\frac{t}{2})p(0)$  at the half-period  $\frac{t}{2}$  and then use  $p(\frac{t}{2})$  during the last half to figure final payment. Now \$1.00 at rate  $r=1$  earns \$2.25.

$$p^{\frac{1}{2}}(t) = (1+r\cdot\frac{t}{2})p(\frac{t}{2}) = (1+r\cdot\frac{t}{2})\cdot(1+r\cdot\frac{t}{2})p(0) = \frac{3}{2}\cdot\frac{3}{2}\cdot 1 = \frac{9}{4} = 2.25$$

*Trimester compounded* interest gives  $p(\frac{t}{3})=(1+r\cdot\frac{t}{3})p(0)$  at the  $1/3^{\text{rd}}$ -period  $\frac{t}{3}$  or 1<sup>st</sup> trimester and then use that to figure the 2<sup>nd</sup> trimester and so on. Now \$1.00 at rate  $r=1$  earns \$2.37.

$$p^{\frac{1}{3}}(t) = (1+r\cdot\frac{t}{3})p(2\frac{t}{3}) = (1+r\cdot\frac{t}{3})\cdot(1+r\cdot\frac{t}{3})p(\frac{t}{3}) = (1+r\cdot\frac{t}{3})\cdot(1+r\cdot\frac{t}{3})\cdot(1+r\cdot\frac{t}{3})p(0) = \frac{4}{3}\cdot\frac{4}{3}\cdot\frac{4}{3}\cdot 1 = \frac{64}{27} = 2.37$$



# The Story of $e$ (A Tale of Great \$Interest\$)

Simple *interest* at some rate  $r$  based on a 1 year period.

You gave a principal  $p(0)$  to the bank and some time  $t$  later they would pay you  $p(t) = (1 + r \cdot t)p(0)$ .


\$1.00 at rate  $r=1$  (like Israel and Brazil that once had 100% interest.) gives \$2.00 at  $t=1$  year.

*Semester compounded* interest gives  $p(\frac{t}{2}) = (1 + r \cdot \frac{t}{2})p(0)$  at the half-period  $\frac{t}{2}$  and then use  $p(\frac{t}{2})$  during the last half to figure final payment. Now \$1.00 at rate  $r=1$  earns \$2.25.

$$p^{\frac{1}{2}}(t) = (1 + r \cdot \frac{t}{2})p(\frac{t}{2}) = (1 + r \cdot \frac{t}{2}) \cdot (1 + r \cdot \frac{t}{2})p(0) = \frac{3}{2} \cdot \frac{3}{2} \cdot 1 = \frac{9}{4} = 2.25$$

*Trimester compounded* interest gives  $p(\frac{t}{3}) = (1 + r \cdot \frac{t}{3})p(0)$  at the  $1/3^{\text{rd}}$ -period  $\frac{t}{3}$  or 1<sup>st</sup> trimester and then use that to figure the 2<sup>nd</sup> trimester and so on. Now \$1.00 at rate  $r=1$  earns \$2.37.

$$p^{\frac{1}{3}}(t) = (1 + r \cdot \frac{t}{3})p(2\frac{t}{3}) = (1 + r \cdot \frac{t}{3}) \cdot (1 + r \cdot \frac{t}{3})p(\frac{t}{3}) = (1 + r \cdot \frac{t}{3}) \cdot (1 + r \cdot \frac{t}{3}) \cdot (1 + r \cdot \frac{t}{3})p(0) = \frac{4}{3} \cdot \frac{4}{3} \cdot \frac{4}{3} \cdot 1 = \frac{64}{27} = 2.37$$



So if you compound interest more and more frequently, do you approach **INFININTEREST?**

# The Story of $e$ (A Tale of Great \$Interest\$)

Simple *interest* at some rate  $r$  based on a 1 year period.

You gave a principal  $p(0)$  to the bank and some time  $t$  later they would pay you  $p(t)=(1+r\cdot t)p(0)$ .

\$1.00 at rate  $r=1$  (like Israel and Brazil that once had 100% interest.) gives \$2.00 at  $t=1$  year.

*Semester compounded* interest gives  $p(\frac{t}{2})=(1+r\cdot\frac{t}{2})p(0)$  at the half-period  $\frac{t}{2}$  and then use  $p(\frac{t}{2})$  during the last half to figure final payment. Now \$1.00 at rate  $r=1$  earns \$2.25.

$$p^{\frac{1}{2}}(t) = (1+r\cdot\frac{t}{2})p(\frac{t}{2}) = (1+r\cdot\frac{t}{2})\cdot(1+r\cdot\frac{t}{2})p(0) = \frac{3}{2}\cdot\frac{3}{2}\cdot 1 = \frac{9}{4} = 2.25$$

*Trimester compounded* interest gives  $p(\frac{t}{3})=(1+r\cdot\frac{t}{3})p(0)$  at the  $1/3^{\text{rd}}$ -period  $\frac{t}{3}$  or 1<sup>st</sup> trimester and then use that to figure the 2<sup>nd</sup> trimester and so on. Now \$1.00 at rate  $r=1$  earns \$2.37.

$$p^{\frac{1}{3}}(t) = (1+r\cdot\frac{t}{3})p(2\frac{t}{3}) = (1+r\cdot\frac{t}{3})\cdot(1+r\cdot\frac{t}{3})p(\frac{t}{3}) = (1+r\cdot\frac{t}{3})\cdot(1+r\cdot\frac{t}{3})\cdot(1+r\cdot\frac{t}{3})p(0) = \frac{4}{3}\cdot\frac{4}{3}\cdot\frac{4}{3}\cdot 1 = \frac{64}{27} = 2.37$$



**NOT!!**



So if you compound interest more and more frequently, do you approach **INFININTEREST?**

# The Story of $e$ (A Tale of Great \$Interest\$)

Simple *interest* at some rate  $r$  based on a 1 year period.

You gave a principal  $p(0)$  to the bank and some time  $t$  later they would pay you  $p(t) = (1 + r \cdot t)p(0)$ .


\$1.00 at rate  $r=1$  (like Israel and Brazil that once had 100% interest.) gives \$2.00 at  $t=1$  year.

*Semester compounded* interest gives  $p(\frac{t}{2}) = (1 + r \cdot \frac{t}{2})p(0)$  at the half-period  $\frac{t}{2}$  and then use  $p(\frac{t}{2})$  during the last half to figure final payment. Now \$1.00 at rate  $r=1$  earns \$2.25.

$$p^{\frac{1}{2}}(t) = (1 + r \cdot \frac{t}{2})p(\frac{t}{2}) = (1 + r \cdot \frac{t}{2}) \cdot (1 + r \cdot \frac{t}{2})p(0) = \frac{3}{2} \cdot \frac{3}{2} \cdot 1 = \frac{9}{4} = 2.25$$

*Trimester compounded* interest gives  $p(\frac{t}{3}) = (1 + r \cdot \frac{t}{3})p(0)$  at the  $1/3^{\text{rd}}$ -period  $\frac{t}{3}$  or 1<sup>st</sup> trimester and then use that to figure the 2<sup>nd</sup> trimester and so on. Now \$1.00 at rate  $r=1$  earns \$2.37.

$$p^{\frac{1}{3}}(t) = (1 + r \cdot \frac{t}{3})p(2\frac{t}{3}) = (1 + r \cdot \frac{t}{3}) \cdot (1 + r \cdot \frac{t}{3})p(\frac{t}{3}) = (1 + r \cdot \frac{t}{3}) \cdot (1 + r \cdot \frac{t}{3}) \cdot (1 + r \cdot \frac{t}{3})p(0) = \frac{4}{3} \cdot \frac{4}{3} \cdot \frac{4}{3} \cdot 1 = \frac{64}{27} = 2.37$$



**NOT!!**



So if you compound interest more and more frequently, do you approach **INFININTEREST?**

$$p^{\frac{1}{1}}(t) = (1 + r \cdot \frac{t}{1})^1 p(0) = \left(\frac{2}{1}\right)^1 \cdot 1 = \frac{2}{1} = 2.00$$

$$p^{\frac{1}{2}}(t) = (1 + r \cdot \frac{t}{2})^2 p(0) = \left(\frac{3}{2}\right)^2 \cdot 1 = \frac{9}{4} = 2.25$$

$$p^{\frac{1}{3}}(t) = (1 + r \cdot \frac{t}{3})^3 p(0) = \left(\frac{4}{3}\right)^3 \cdot 1 = \frac{64}{27} = 2.37$$

$$p^{\frac{1}{4}}(t) = (1 + r \cdot \frac{t}{4})^4 p(0) = \left(\frac{5}{4}\right)^4 \cdot 1 = \frac{625}{256} = 2.44$$

# The Story of $e$ (A Tale of Great \$Interest\$)

Simple *interest* at some rate  $r$  based on a 1 year period.

You gave a principal  $p(0)$  to the bank and some time  $t$  later they would pay you  $p(t) = (1+r \cdot t)p(0)$ .

\$1.00 at rate  $r=1$  (like Israel and Brazil that once had 100% interest.) gives \$2.00 at  $t=1$  year.

*Semester compounded* interest gives  $p(\frac{t}{2}) = (1+r \cdot \frac{t}{2})p(0)$  at the half-period  $\frac{t}{2}$  and then use  $p(\frac{t}{2})$  during the last half to figure final payment. Now \$1.00 at rate  $r=1$  earns \$2.25.

$$p^{\frac{1}{2}}(t) = (1+r \cdot \frac{t}{2})p(\frac{t}{2}) = (1+r \cdot \frac{t}{2}) \cdot (1+r \cdot \frac{t}{2})p(0) = \frac{3}{2} \cdot \frac{3}{2} \cdot 1 = \frac{9}{4} = 2.25$$

*Trimester compounded* interest gives  $p(\frac{t}{3}) = (1+r \cdot \frac{t}{3})p(0)$  at the  $1/3^{\text{rd}}$ -period  $\frac{t}{3}$  or 1<sup>st</sup> trimester and then use that to figure the 2<sup>nd</sup> trimester and so on. Now \$1.00 at rate  $r=1$  earns \$2.37.

$$p^{\frac{1}{3}}(t) = (1+r \cdot \frac{t}{3})p(2\frac{t}{3}) = (1+r \cdot \frac{t}{3}) \cdot (1+r \cdot \frac{t}{3})p(\frac{t}{3}) = (1+r \cdot \frac{t}{3}) \cdot (1+r \cdot \frac{t}{3}) \cdot (1+r \cdot \frac{t}{3})p(0) = \frac{4}{3} \cdot \frac{4}{3} \cdot \frac{4}{3} \cdot 1 = \frac{64}{27} = 2.37$$



So if you compound interest more and more frequently, do you approach **INFININTEREST?**

**NOT!!**

$$p^{\frac{1}{1}}(t) = (1+r \cdot \frac{t}{1})^1 p(0) = \left(\frac{2}{1}\right)^1 \cdot 1 = \frac{2}{1} = 2.00$$

$$p^{\frac{1}{2}}(t) = (1+r \cdot \frac{t}{2})^2 p(0) = \left(\frac{3}{2}\right)^2 \cdot 1 = \frac{9}{4} = 2.25$$

$$p^{\frac{1}{3}}(t) = (1+r \cdot \frac{t}{3})^3 p(0) = \left(\frac{4}{3}\right)^3 \cdot 1 = \frac{64}{27} = 2.37$$

$$p^{\frac{1}{4}}(t) = (1+r \cdot \frac{t}{4})^4 p(0) = \left(\frac{5}{4}\right)^4 \cdot 1 = \frac{625}{256} = 2.44$$

Monthly:  $p^{\frac{1}{12}}(t) = (1+r \cdot \frac{t}{12})^{12} p(0) = \left(\frac{13}{12}\right)^{12} \cdot 1 = 2.613$

Weekly:  $p^{\frac{1}{52}}(t) = (1+r \cdot \frac{t}{52})^{52} p(0) = \left(\frac{53}{52}\right)^{52} \cdot 1 = 2.693$

Daily:  $p^{\frac{1}{365}}(t) = (1+r \cdot \frac{t}{365})^{365} p(0) = \left(\frac{366}{365}\right)^{365} \cdot 1 = 2.7145$

Hrly:  $p^{\frac{1}{8760}}(t) = (1+r \cdot \frac{t}{8760})^{8760} p(0) = \left(\frac{8761}{8760}\right)^{8760} \cdot 1 = 2.7181$



Interest product formula is really inefficient:  $10^6$  products for 6-figures! ..  $10^9$  products for 9 ...

$$p^{1/m}(1) = \left(1 + \frac{1}{m}\right)^m \xrightarrow{m \rightarrow \infty} \mathbf{2.718281828459..} = e$$

Let:  $m \cdot r \cdot t = n$

$$\left(1 + \frac{1}{m}\right)^{m \cdot r \cdot t} \xrightarrow{m \rightarrow \infty} e^{r \cdot t}$$

or:  $1/m = r \cdot t / n$

$$\left(1 + \frac{r \cdot t}{n}\right)^n \xrightarrow{n \rightarrow \infty} e^{r \cdot t}$$

$p^{1/m}(1) = \mathbf{2.7169239322}$	for $m = 1,000$
$p^{1/m}(1) = \mathbf{2.7181459268}$	for $m = 10,000$
$p^{1/m}(1) = \mathbf{2.7182682372}$	for $m = 100,000$
$p^{1/m}(1) = \mathbf{2.7182804693}$	for $m = 1,000,000$
$p^{1/m}(1) = \mathbf{2.7182816925}$	for $m = 10,000,000$
$p^{1/m}(1) = \mathbf{2.7182818149}$	for $m = 100,000,000$
$p^{1/m}(1) = \mathbf{2.7182818271}$	for $m = 1,000,000,000$

Interest product formula is really inefficient:  $10^6$  products for 6-figures! ..  $10^9$  products for 9 ...

	$p^{1/m}(1) = (1 + \frac{1}{m})^m \xrightarrow{m \rightarrow \infty} 2.718281828459.. = e$	$p^{1/m}(1) = 2.7169239322$	for $m = 1,000$
		$p^{1/m}(1) = 2.7181459268$	for $m = 10,000$
		$p^{1/m}(1) = 2.7182682372$	for $m = 100,000$
		$p^{1/m}(1) = 2.7182804693$	for $m = 1,000,000$
Let: $m \cdot r \cdot t = n$	$(1 + \frac{1}{m})^{m \cdot r \cdot t} \xrightarrow{m \rightarrow \infty} e^{r \cdot t}$	$p^{1/m}(1) = 2.7182816925$	for $m = 10,000,000$
or: $1/m = r \cdot t/n$	$(1 + \frac{r \cdot t}{n})^n \xrightarrow{n \rightarrow \infty} e^{r \cdot t}$	$p^{1/m}(1) = 2.7182818149$	for $m = 100,000,000$
		$p^{1/m}(1) = 2.7182818271$	for $m = 1,000,000,000$

Can improve computational efficiency using binomial theorem:

$$(x + y)^n = x^n + n \cdot x^{n-1}y + \frac{n(n-1)}{2!} x^{n-2}y^2 + \frac{n(n-1)(n-2)}{3!} x^{n-3}y^3 + \dots + n \cdot xy^{n-1} + y^n$$

$$(1 + \frac{r \cdot t}{n})^n = 1 + n \cdot \left(\frac{r \cdot t}{n}\right) + \frac{n(n-1)}{2!} \left(\frac{r \cdot t}{n}\right)^2 + \frac{n(n-1)(n-2)}{3!} \left(\frac{r \cdot t}{n}\right)^3 + \dots$$

Define: Factorials(!):  
 $0! = 1 = 1!$ ,  $2! = 1 \cdot 2$ ,  $3! = 1 \cdot 2 \cdot 3, \dots$

Interest product formula is really inefficient:  $10^6$  products for 6-figures! ..  $10^9$  products for 9 ...

	$p^{1/m}(1) = (1 + \frac{1}{m})^m \xrightarrow{m \rightarrow \infty} 2.718281828459.. = e$	$p^{1/m}(1) = 2.7169239322$	for $m = 1,000$
		$p^{1/m}(1) = 2.7181459268$	for $m = 10,000$
		$p^{1/m}(1) = 2.7182682372$	for $m = 100,000$
		$p^{1/m}(1) = 2.7182804693$	for $m = 1,000,000$
Let: $m \cdot r \cdot t = n$	$(1 + \frac{1}{m})^{m \cdot r \cdot t} \xrightarrow{m \rightarrow \infty} e^{r \cdot t}$	$p^{1/m}(1) = 2.7182816925$	for $m = 10,000,000$
or: $1/m = r \cdot t/n$	$(1 + \frac{r \cdot t}{n})^n \xrightarrow{n \rightarrow \infty} e^{r \cdot t}$	$p^{1/m}(1) = 2.7182818149$	for $m = 100,000,000$
		$p^{1/m}(1) = 2.7182818271$	for $m = 1,000,000,000$

Can improve computational efficiency using binomial theorem:

$$(x + y)^n = x^n + n \cdot x^{n-1}y + \frac{n(n-1)}{2!} x^{n-2}y^2 + \frac{n(n-1)(n-2)}{3!} x^{n-3}y^3 + \dots + n \cdot xy^{n-1} + y^n$$

$$(1 + \frac{r \cdot t}{n})^n = 1 + n \cdot \left(\frac{r \cdot t}{n}\right) + \frac{n(n-1)}{2!} \left(\frac{r \cdot t}{n}\right)^2 + \frac{n(n-1)(n-2)}{3!} \left(\frac{r \cdot t}{n}\right)^3 + \dots$$

Define: Factorials(!):

$$0! = 1 = 1!, \quad 2! = 1 \cdot 2, \quad 3! = 1 \cdot 2 \cdot 3, \dots$$

As  $n \rightarrow \infty$  let :

$$n(n-1) \rightarrow n^2,$$

$$n(n-1)(n-2) \rightarrow n^3, \text{ etc.}$$

$$e^{r \cdot t} = 1 + r \cdot t + \frac{1}{2!} (r \cdot t)^2 + \frac{1}{3!} (r \cdot t)^3 + \dots = \sum_{p=0}^{\infty} \frac{(r \cdot t)^p}{p!}$$



Interest product formula is really inefficient:  $10^6$  products for 6-figures! ..  $10^9$  products for 9 ...

	$p^{1/m}(1) = (1 + \frac{1}{m})^m \xrightarrow{m \rightarrow \infty} 2.718281828459.. = e$	$p^{1/m}(1) = 2.7169239322$	for $m = 1,000$
		$p^{1/m}(1) = 2.7181459268$	for $m = 10,000$
		$p^{1/m}(1) = 2.7182682372$	for $m = 100,000$
		$p^{1/m}(1) = 2.7182804693$	for $m = 1,000,000$
Let: $m \cdot r \cdot t = n$	$(1 + \frac{1}{m})^{m \cdot r \cdot t} \xrightarrow{m \rightarrow \infty} e^{r \cdot t}$	$p^{1/m}(1) = 2.7182816925$	for $m = 10,000,000$
or: $1/m = r \cdot t/n$	$(1 + \frac{r \cdot t}{n})^n \xrightarrow{n \rightarrow \infty} e^{r \cdot t}$	$p^{1/m}(1) = 2.7182818149$	for $m = 100,000,000$
		$p^{1/m}(1) = 2.7182818271$	for $m = 1,000,000,000$

Can improve computational efficiency using binomial theorem:

$$(x + y)^n = x^n + n \cdot x^{n-1}y + \frac{n(n-1)}{2!} x^{n-2}y^2 + \frac{n(n-1)(n-2)}{3!} x^{n-3}y^3 + \dots + n \cdot xy^{n-1} + y^n$$

$$(1 + \frac{r \cdot t}{n})^n = 1 + n \cdot \left(\frac{r \cdot t}{n}\right) + \frac{n(n-1)}{2!} \left(\frac{r \cdot t}{n}\right)^2 + \frac{n(n-1)(n-2)}{3!} \left(\frac{r \cdot t}{n}\right)^3 + \dots$$

Define: Factorials(!):

$$0! = 1 = 1!, \quad 2! = 1 \cdot 2, \quad 3! = 1 \cdot 2 \cdot 3, \dots$$

As  $n \rightarrow \infty$  let :

$$n(n-1) \rightarrow n^2,$$

$$n(n-1)(n-2) \rightarrow n^3, \text{ etc.}$$

$$e^{r \cdot t} = 1 + r \cdot t + \frac{1}{2!} (r \cdot t)^2 + \frac{1}{3!} (r \cdot t)^3 + \dots = \sum_{p=0}^{\infty} \frac{(r \cdot t)^p}{p!}$$

Precision order:	$(o=1)$ -e-series = 2.00000 = 1+1
	$(o=2)$ -e-series = 2.50000 = 1+1+1/2
	$(o=3)$ -e-series = 2.66667 = 1+1+1/2+1/6
	$(o=4)$ -e-series = 2.70833 = 1+1+1/2+1/6+1/24
	$(o=5)$ -e-series = 2.71667 = 1+1+1/2+1/6+1/24+1/120
	$(o=6)$ -e-series = 2.71805 = 1+1+1/2+1/6+1/24+1/120+1/720
	$(o=7)$ -e-series = 2.71825
	$(o=8)$ -e-series = 2.71828

About 12 summed quotients for 6-figure precision (A lot better!)



## *Power Series Good! Need general power series development*

Start with a general power series with constant coefficients  $c_0, c_1, \text{ etc.}$

Set  $t=0$  to get  $c_0 = x(0)$ .

$$x(t) = c_0 + c_1t + c_2t^2 + c_3t^3 + c_4t^4 + c_5t^5 + \dots + c_nt^n +$$

## *Power Series Good! Need general power series development*

Start with a general power series with constant coefficients  $c_0, c_1, \text{ etc.}$

Set  $t=0$  to get  $c_0 = x(0)$ .

$$x(t) = c_0 + c_1t + c_2t^2 + c_3t^3 + c_4t^4 + c_5t^5 + \dots + c_nt^n +$$

Rate of change of position  $x(t)$  is *velocity*  $v(t)$ .

Set  $t=0$  to get  $c_1 = v(0)$ .

$$v(t) = \frac{d}{dt}x(t) = 0 + c_1 + 2c_2t + 3c_3t^2 + 4c_4t^3 + 5c_5t^4 + \dots + nc_nt^{n-1} +$$

## *Power Series Good! Need general power series development*

Start with a general power series with constant coefficients  $c_0, c_1, \text{ etc.}$

Set  $t=0$  to get  $c_0 = x(0)$ .

$$x(t) = c_0 + c_1t + c_2t^2 + c_3t^3 + c_4t^4 + c_5t^5 + \dots + c_nt^n +$$

Rate of change of position  $x(t)$  is *velocity*  $v(t)$ .

Set  $t=0$  to get  $c_1 = v(0)$ .

$$v(t) = \frac{d}{dt}x(t) = 0 + c_1 + 2c_2t + 3c_3t^2 + 4c_4t^3 + 5c_5t^4 + \dots + nc_nt^{n-1} +$$

Change of velocity  $v(t)$  is *acceleration*  $a(t)$ .

Set  $t=0$  to get  $c_2 = \frac{1}{2}a(0)$ .

$$a(t) = \frac{d}{dt}v(t) = 0 + 2c_2 + 2 \cdot 3c_3t + 3 \cdot 4c_4t^2 + 4 \cdot 5c_5t^3 + \dots + n(n-1)c_nt^{n-2} +$$

## *Power Series Good! Need general power series development*

Start with a general power series with constant coefficients  $c_0, c_1, \text{ etc.}$

Set  $t=0$  to get  $c_0 = x(0)$ .

$$x(t) = c_0 + c_1t + c_2t^2 + c_3t^3 + c_4t^4 + c_5t^5 + \dots + c_nt^n +$$

Rate of change of position  $x(t)$  is *velocity*  $v(t)$ .

Set  $t=0$  to get  $c_1 = v(0)$ .

$$v(t) = \frac{d}{dt}x(t) = 0 + c_1 + 2c_2t + 3c_3t^2 + 4c_4t^3 + 5c_5t^4 + \dots + nc_nt^{n-1} +$$

Change of velocity  $v(t)$  is *acceleration*  $a(t)$ .

Set  $t=0$  to get  $c_2 = \frac{1}{2}a(0)$ .

$$a(t) = \frac{d}{dt}v(t) = 0 + 2c_2 + 2 \cdot 3c_3t + 3 \cdot 4c_4t^2 + 4 \cdot 5c_5t^3 + \dots + n(n-1)c_nt^{n-2} +$$

Change of acceleration  $a(t)$  is *jerk*  $j(t)$ . (*Jerk* is NASA term.)

Set  $t=0$  to get  $c_3 = \frac{1}{3!}j(0)$ .

$$j(t) = \frac{d}{dt}a(t) = 0 + 2 \cdot 3c_3 + 2 \cdot 3 \cdot 4c_4t + 3 \cdot 4 \cdot 5c_5t^2 + \dots + n(n-1)(n-2)c_nt^{n-3} +$$

## Power Series Good! Need general power series development

Start with a general power series with constant coefficients  $c_0, c_1, \text{ etc.}$

Set  $t=0$  to get  $c_0 = x(0)$ .

$$x(t) = c_0 + c_1t + c_2t^2 + c_3t^3 + c_4t^4 + c_5t^5 + \dots + c_nt^n +$$

Rate of change of position  $x(t)$  is *velocity*  $v(t)$ .

Set  $t=0$  to get  $c_1 = v(0)$ .

$$v(t) = \frac{d}{dt}x(t) = 0 + c_1 + 2c_2t + 3c_3t^2 + 4c_4t^3 + 5c_5t^4 + \dots + nc_nt^{n-1} +$$

Change of velocity  $v(t)$  is *acceleration*  $a(t)$ .

Set  $t=0$  to get  $c_2 = \frac{1}{2}a(0)$ .

$$a(t) = \frac{d}{dt}v(t) = 0 + 2c_2 + 2 \cdot 3c_3t + 3 \cdot 4c_4t^2 + 4 \cdot 5c_5t^3 + \dots + n(n-1)c_nt^{n-2} +$$

Change of acceleration  $a(t)$  is *jerk*  $j(t)$ . (*Jerk* is NASA term.)

Set  $t=0$  to get  $c_3 = \frac{1}{3!}j(0)$ .

$$j(t) = \frac{d}{dt}a(t) = 0 + 2 \cdot 3c_3 + 2 \cdot 3 \cdot 4c_4t + 3 \cdot 4 \cdot 5c_5t^2 + \dots + n(n-1)(n-2)c_nt^{n-3} +$$

Change of jerk  $j(t)$  is *inauguration*  $i(t)$ . (Be silly like NASA!)

Set  $t=0$  to get  $c_4 = \frac{1}{4!}i(0)$ .

$$i(t) = \frac{d}{dt}j(t) = 0 + 2 \cdot 3 \cdot 4c_4 + 2 \cdot 3 \cdot 4 \cdot 5c_5t + \dots + n(n-1)(n-2)(n-3)c_nt^{n-4} +$$

## Power Series Good! Need general power series development

Start with a general power series with constant coefficients  $c_0, c_1, \text{ etc.}$

Set  $t=0$  to get  $c_0 = x(0)$ .

$$x(t) = c_0 + c_1 t + c_2 t^2 + c_3 t^3 + c_4 t^4 + c_5 t^5 + \dots + c_n t^n +$$

Rate of change of position  $x(t)$  is *velocity*  $v(t)$ .

Set  $t=0$  to get  $c_1 = v(0)$ .

$$v(t) = \frac{d}{dt} x(t) = 0 + c_1 + 2c_2 t + 3c_3 t^2 + 4c_4 t^3 + 5c_5 t^4 + \dots + n c_n t^{n-1} +$$

Change of velocity  $v(t)$  is *acceleration*  $a(t)$ .

Set  $t=0$  to get  $c_2 = \frac{1}{2} a(0)$ .

$$a(t) = \frac{d}{dt} v(t) = 0 + 2c_2 + 2 \cdot 3c_3 t + 3 \cdot 4c_4 t^2 + 4 \cdot 5c_5 t^3 + \dots + n(n-1)c_n t^{n-2} +$$

Change of acceleration  $a(t)$  is *jerk*  $j(t)$ . (*Jerk* is NASA term.)

Set  $t=0$  to get  $c_3 = \frac{1}{3!} j(0)$ .

$$j(t) = \frac{d}{dt} a(t) = 0 + 2 \cdot 3c_3 + 2 \cdot 3 \cdot 4c_4 t + 3 \cdot 4 \cdot 5c_5 t^2 + \dots + n(n-1)(n-2)c_n t^{n-3} +$$

Change of jerk  $j(t)$  is *inauguration*  $i(t)$ . (Be silly like NASA!)

Set  $t=0$  to get  $c_4 = \frac{1}{4!} i(0)$ .

$$i(t) = \frac{d}{dt} j(t) = 0 + 2 \cdot 3 \cdot 4c_4 + 2 \cdot 3 \cdot 4 \cdot 5c_5 t + \dots + n(n-1)(n-2)(n-3)c_n t^{n-4} +$$

*Gives Maclaurin (or Taylor) power series*

$$x(t) = x(0) + v(0)t + \frac{1}{2!} a(0)t^2 + \frac{1}{3!} j(0)t^3 + \frac{1}{4!} i(0)t^4 + \frac{1}{5!} r(0)t^5 + \dots + \frac{1}{n!} x^{(n)} t^n +$$

## Power Series Good! Need general power series development

Start with a general power series with constant coefficients  $c_0, c_1, \text{ etc.}$

Set  $t=0$  to get  $c_0 = x(0)$ .

$$x(t) = c_0 + c_1t + c_2t^2 + c_3t^3 + c_4t^4 + c_5t^5 + \dots + c_nt^n +$$

Rate of change of position  $x(t)$  is *velocity*  $v(t)$ .

Set  $t=0$  to get  $c_1 = v(0)$ .

$$v(t) = \frac{d}{dt}x(t) = 0 + c_1 + 2c_2t + 3c_3t^2 + 4c_4t^3 + 5c_5t^4 + \dots + nc_nt^{n-1} +$$

Change of velocity  $v(t)$  is *acceleration*  $a(t)$ .

Set  $t=0$  to get  $c_2 = \frac{1}{2}a(0)$ .

$$a(t) = \frac{d}{dt}v(t) = 0 + 2c_2 + 2 \cdot 3c_3t + 3 \cdot 4c_4t^2 + 4 \cdot 5c_5t^3 + \dots + n(n-1)c_nt^{n-2} +$$

Change of acceleration  $a(t)$  is *jerk*  $j(t)$ . (*Jerk* is NASA term.)

Set  $t=0$  to get  $c_3 = \frac{1}{3!}j(0)$ .

$$j(t) = \frac{d}{dt}a(t) = 0 + 2 \cdot 3c_3 + 2 \cdot 3 \cdot 4c_4t + 3 \cdot 4 \cdot 5c_5t^2 + \dots + n(n-1)(n-2)c_nt^{n-3} +$$

Change of jerk  $j(t)$  is *inauguration*  $i(t)$ . (Be silly like NASA!)

Set  $t=0$  to get  $c_4 = \frac{1}{4!}i(0)$ .

$$i(t) = \frac{d}{dt}j(t) = 0 + 2 \cdot 3 \cdot 4c_4 + 2 \cdot 3 \cdot 4 \cdot 5c_5t + \dots + n(n-1)(n-2)(n-3)c_nt^{n-4} +$$

*Gives Maclaurin (or Taylor) power series*

$$x(t) = x(0) + v(0)t + \frac{1}{2!}a(0)t^2 + \frac{1}{3!}j(0)t^3 + \frac{1}{4!}i(0)t^4 + \frac{1}{5!}r(0)t^5 + \dots + \frac{1}{n!}x^{(n)}t^n +$$

Good old UP I formula!

# Power Series Good! Need general power series development

Start with a general power series with constant coefficients  $c_0, c_1, \text{ etc.}$

Set  $t=0$  to get  $c_0 = x(0)$ .

$$x(t) = c_0 + c_1 t + c_2 t^2 + c_3 t^3 + c_4 t^4 + c_5 t^5 + \dots + c_n t^n +$$

Rate of change of position  $x(t)$  is *velocity*  $v(t)$ .

Set  $t=0$  to get  $c_1 = v(0)$ .

$$v(t) = \frac{d}{dt} x(t) = 0 + c_1 + 2c_2 t + 3c_3 t^2 + 4c_4 t^3 + 5c_5 t^4 + \dots + n c_n t^{n-1} +$$

Change of velocity  $v(t)$  is *acceleration*  $a(t)$ .

Set  $t=0$  to get  $c_2 = \frac{1}{2} a(0)$ .

$$a(t) = \frac{d}{dt} v(t) = 0 + 2c_2 + 2 \cdot 3c_3 t + 3 \cdot 4c_4 t^2 + 4 \cdot 5c_5 t^3 + \dots + n(n-1)c_n t^{n-2} +$$

Change of acceleration  $a(t)$  is *jerk*  $j(t)$ . (*Jerk* is NASA term.)

Set  $t=0$  to get  $c_3 = \frac{1}{3!} j(0)$ .

$$j(t) = \frac{d}{dt} a(t) = 0 + 2 \cdot 3c_3 + 2 \cdot 3 \cdot 4c_4 t + 3 \cdot 4 \cdot 5c_5 t^2 + \dots + n(n-1)(n-2)c_n t^{n-3} +$$

Change of jerk  $j(t)$  is *inauguration*  $i(t)$ . (Be silly like NASA!)

Set  $t=0$  to get  $c_4 = \frac{1}{4!} i(0)$ .

$$i(t) = \frac{d}{dt} j(t) = 0 + 2 \cdot 3 \cdot 4c_4 + 2 \cdot 3 \cdot 4 \cdot 5c_5 t + \dots + n(n-1)(n-2)(n-3)c_n t^{n-4} +$$

Gives Maclaurin (or Taylor) power series

$$x(t) = x(0) + v(0)t + \frac{1}{2!} a(0)t^2 + \frac{1}{3!} j(0)t^3 + \frac{1}{4!} i(0)t^4 + \frac{1}{5!} r(0)t^5 + \dots + \frac{1}{n!} x^{(n)} t^n +$$

Setting all initial values to  $1 = x(0) = v(0) = a(0) = j(0) = i(0) = \dots$


Good old UP I formula!

gives exponential:

$$e^t = 1 + t + \frac{1}{2!} t^2 + \frac{1}{3!} t^3 + \frac{1}{4!} t^4 + \frac{1}{5!} t^5 + \dots + \frac{1}{n!} t^n +$$



But, how good are power series?



Unit 1  
Fig. 10.2

Gives Maclaurin (or Taylor) power series


$$x(t) = x(0) + v(0)t + \frac{1}{2!} a(0)t^2 + \frac{1}{3!} j(0)t^3 + \frac{1}{4!} i(0)t^4 + \frac{1}{5!} r(0)t^5 + \dots + \frac{1}{n!} x^{(n)}t^n +$$

Setting all initial values to  $1 = x(0) = v(0) = a(0) = j(0) = i(0) = \dots$


gives exponential: 
$$e^t = 1 + t + \frac{1}{2!} t^2 + \frac{1}{3!} t^3 + \frac{1}{4!} t^4 + \frac{1}{5!} t^5 + \dots + \frac{1}{n!} t^n +$$

# How good are power series? Depends...

$$x(t) = \boxed{\cos t} = 1 + 0 - \frac{t^2}{2!} + 0 + \frac{t^4}{4!} + 0 - \frac{t^6}{6!} + 0 + \frac{t^8}{8!} \dots$$



$$x(t) = \boxed{\sin t} = 0 + t + 0 - \frac{t^3}{3!} + 0 + \frac{t^5}{5!} + 0 - \frac{t^7}{7!} + 0 + \frac{t^9}{9!} \dots$$



Unit 1  
Fig. 10.3

# *1. The Story of $e$ (A Tale of Great \$Interest\$)*

*How good are those power series?*

*Taylor-Maclaurin series,*



*imaginary interest, and complex exponentials*

Suppose the fancy bankers really went bonkers and made interest rate  $r$  an *imaginary number*  $r=i\theta$ .

Imaginary number  $i=\sqrt{-1}$  powers have *repeat-after-4-pattern*:  $i^0=1, i^1=i, i^2=-1, i^3=-i, i^4=1, etc...$

$$\begin{aligned} e^{i\theta} &= 1 + i\theta + \frac{(i\theta)^2}{2!} + \frac{(i\theta)^3}{3!} + \frac{(i\theta)^4}{4!} + \frac{(i\theta)^5}{5!} + \dots && \text{(From exponential series)} \\ &= 1 + i\theta - \frac{\theta^2}{2!} - i\frac{\theta^3}{3!} + \frac{\theta^4}{4!} + i\frac{\theta^5}{5!} - \dots && (i = \sqrt{-1} \text{ implies: } i^1=i, i^2=-1, i^3=-i, i^4=+1, i^5=i, \dots) \\ &= \left( 1 - \frac{\theta^2}{2!} + \frac{\theta^4}{4!} - \dots \right) + \left( i\theta - i\frac{\theta^3}{3!} + i\frac{\theta^5}{5!} - \dots \right) \end{aligned}$$

Suppose the fancy bankers really went bonkers and made interest rate  $r$  an *imaginary number*  $r=i\theta$ .

Imaginary number  $i=\sqrt{-1}$  powers have *repeat-after-4-pattern*:  $i^0=1, i^1=i, i^2=-1, i^3=-i, i^4=1, etc...$



$$e^{i\theta} = 1 + i\theta + \frac{(i\theta)^2}{2!} + \frac{(i\theta)^3}{3!} + \frac{(i\theta)^4}{4!} + \frac{(i\theta)^5}{5!} + \dots \quad (\text{From exponential series})$$

$$= 1 + i\theta - \frac{\theta^2}{2!} - i\frac{\theta^3}{3!} + \frac{\theta^4}{4!} + i\frac{\theta^5}{5!} - \dots \quad (i = \sqrt{-1} \text{ implies: } i^1=i, i^2=-1, i^3=-i, i^4=+1, i^5=i, \dots)$$

$$= \left( 1 - \frac{\theta^2}{2!} + \frac{\theta^4}{4!} - \dots \right) + \left( i\theta - i\frac{\theta^3}{3!} + i\frac{\theta^5}{5!} - \dots \right) \quad \text{To match series for } \begin{cases} \text{cosine : } \cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots \\ \text{sine : } \sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots \end{cases}$$

$$e^{i\theta} = \cos \theta + i \sin \theta$$

*Euler-DeMoivre Theorem*



Unit 1  
Fig. 10.3

Suppose the fancy bankers really went bonkers and made interest rate  $r$  an *imaginary number*  $r=i\theta$ .

Imaginary number  $i=\sqrt{-1}$  powers have *repeat-after-4-pattern*:  $i^0=1, i^1=i, i^2=-1, i^3=-i, i^4=1, etc...$




$$e^{i\theta} = 1 + i\theta + \frac{(i\theta)^2}{2!} + \frac{(i\theta)^3}{3!} + \frac{(i\theta)^4}{4!} + \frac{(i\theta)^5}{5!} + \dots \quad (\text{From exponential series})$$

$$= 1 + i\theta - \frac{\theta^2}{2!} - i\frac{\theta^3}{3!} + \frac{\theta^4}{4!} + i\frac{\theta^5}{5!} - \dots \quad (i = \sqrt{-1} \text{ implies: } i^1=i, i^2=-1, i^3=-i, i^4=+1, i^5=i, \dots)$$

$$= \left( 1 - \frac{\theta^2}{2!} + \frac{\theta^4}{4!} - \dots \right) + \left( i\theta - i\frac{\theta^3}{3!} + i\frac{\theta^5}{5!} - \dots \right) \quad \text{To match series for } \begin{cases} \text{cosine : } \cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots \\ \text{sine : } \sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots \end{cases}$$

$$e^{i\theta} = \cos \theta + i \sin \theta$$

*Euler-DeMoivre Theorem*



Unit 1  
Fig. 10.3

## 2. *What Good Are Complex Exponentials?*

*Easy trig*



*Easy 2D vector analysis*



*Easy oscillator phase analysis*

*Easy rotation and “dot” or “cross” products*

# What Good Are Complex Exponentials?

## 1. Complex numbers provide "automatic trigonometry"

Can't remember  $\cos(a+b)$  or  $\sin(a+b)$ ? Just factor  $e^{i(a+b)} = e^{ia}e^{ib} \dots$

$$\begin{aligned} e^{i(a+b)} &= e^{ia} e^{ib} \\ \cos(a+b) + i \sin(a+b) &= (\cos a + i \sin a) (\cos b + i \sin b) \\ \boxed{\cos(a+b)} + i \boxed{\sin(a+b)} &= \boxed{[\cos a \cos b - \sin a \sin b]} + i \boxed{[\sin a \cos b + \cos a \sin b]} \end{aligned}$$



# What Good Are Complex Exponentials?

## 1. Complex numbers provide "automatic trigonometry"

Can't remember  $\cos(a+b)$  or  $\sin(a+b)$ ? Just factor  $e^{i(a+b)} = e^{ia}e^{ib} \dots$


$$e^{i(a+b)} = e^{ia} e^{ib}$$

$$\cos(a+b) + i \sin(a+b) = (\cos a + i \sin a) (\cos b + i \sin b)$$

$$\boxed{\cos(a+b)} + i \boxed{\sin(a+b)} = \boxed{[\cos a \cos b - \sin a \sin b]} + i \boxed{[\sin a \cos b + \cos a \sin b]}$$

2. Complex numbers add like vectors.  $z_{sum} = z + z' = (x + iy) + (x' + iy') = (x + x') + i(y + y')$

$$z_{diff} = z - z' = (x + iy) - (x' + iy') = (x - x') + i(y - y')$$



Unit 1  
Fig. 10.6



$$|z_{SUM}| = \sqrt{(z+z')^* (z+z')} = \sqrt{(re^{i\phi} + r'e^{i\phi'})^* (re^{i\phi} + r'e^{i\phi'})} = \sqrt{(re^{-i\phi} + r'e^{-i\phi'}) (re^{i\phi} + r'e^{i\phi'})}$$

$$= \sqrt{r^2 + r'^2 + rr'(e^{i(\phi-\phi')} + e^{-i(\phi-\phi')})} = \sqrt{r^2 + r'^2 + 2rr' \cos(\phi - \phi')} \quad (\text{quick derivation of Cosine Law})$$

# What Good Are Complex Exponentials? (contd.)


## 3. Complex exponentials $Ae^{-i\omega t}$ track position and velocity using Phasor Clock.

(a) Complex plane and unit vectors



(b) Quantum Phasor Clock  $\psi = Ae^{-i\omega t} = A\cos\omega t - iA\sin\omega t = x + iy$


Unit 1  
Fig. 10.5




# What Good Are Complex Exponentials? (contd.)

## 3. Complex exponentials $Ae^{-i\omega t}$ track position and velocity using Phasor Clock.

(a) Complex plane and unit vectors



(b) Quantum Phasor Clock  $\psi = Ae^{-i\omega t} = A\cos\omega t - iA\sin\omega t = x + iy$



Unit 1  
Fig. 10.5

Some Rect-vs-Polar relations worth remembering

$$\text{Cartesian } (x,y) \text{ form } \begin{cases} \psi_x = \text{Re } \psi(t) = x(t) = A \cos \omega t = \frac{\psi + \psi^*}{2} \\ \psi_y = \text{Im } \psi(t) = \frac{v(t)}{\omega} = -A \sin \omega t = \frac{\psi - \psi^*}{2i} \end{cases}$$

$$\psi = r e^{+i\theta} = r e^{-i\omega t} = r(\cos \omega t - i \sin \omega t)$$

$$\psi^* = r e^{-i\theta} = r e^{+i\omega t} = r(\cos \omega t + i \sin \omega t)$$

$$\text{Polar } (r,\theta) \text{ form } \begin{cases} r = A = |\psi| = \sqrt{\psi_x^2 + \psi_y^2} = \sqrt{\psi^* \psi} \\ \theta = -\omega t = \arctan(\psi_y / \psi_x) \end{cases}$$

$$\cos \theta = \frac{1}{2}(e^{+i\theta} + e^{-i\theta}) \quad \text{Re } \psi = \frac{\psi + \psi^*}{2}$$

$$\sin \theta = \frac{1}{2i}(e^{+i\theta} - e^{-i\theta}) \quad \text{Im } \psi = \frac{\psi - \psi^*}{2i}$$

## *2. What Good Are Complex Exponentials?*

*Easy trig*

*Easy 2D vector analysis*

*Easy oscillator phase analysis*

 *Easy rotation and “dot” or “cross” products*

## What Good Are Complex Exponentials? (contd.)

4. Complex products provide 2D rotation operations.

$$e^{i\phi} \cdot z = (\cos\phi + i \sin\phi) \cdot (x + iy) = x \cos\phi - y \sin\phi + i (x \sin\phi + y \cos\phi)$$

$$\mathbf{R}_{+\phi} \cdot \mathbf{r} = (x \cos\phi - y \sin\phi) \hat{\mathbf{e}}_x + (x \sin\phi + y \cos\phi) \hat{\mathbf{e}}_y$$
$$\begin{pmatrix} \cos\phi & -\sin\phi \\ \sin\phi & \cos\phi \end{pmatrix} \cdot \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \cos\phi - y \sin\phi \\ x \sin\phi + y \cos\phi \end{pmatrix}$$

## What Good Are Complex Exponentials? (contd.)


### 4. Complex products provide 2D rotation operations.

$$e^{i\phi} \cdot z = (\cos\phi + i \sin\phi) \cdot (x + iy) = x \cos\phi - y \sin\phi + i (x \sin\phi + y \cos\phi)$$


$$\mathbf{R}_{+\phi} \cdot \mathbf{r} = (x \cos\phi - y \sin\phi) \hat{\mathbf{e}}_x + (x \sin\phi + y \cos\phi) \hat{\mathbf{e}}_y$$

$$\begin{pmatrix} \cos\phi & -\sin\phi \\ \sin\phi & \cos\phi \end{pmatrix} \cdot \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \cos\phi - y \sin\phi \\ x \sin\phi + y \cos\phi \end{pmatrix}$$

$e^{i\phi}$  acts on this:  $z = re^{i\theta}$



to give this:  $e^{i\phi} e^{i\theta} z = re^{i\phi} e^{i\theta}$



## What Good Are Complex Exponentials? (contd.)

### 4. Complex products provide 2D rotation operations.

$$e^{i\phi} \cdot z = (\cos\phi + i \sin\phi) \cdot (x + iy) = x \cos\phi - y \sin\phi + i(x \sin\phi + y \cos\phi)$$

$$\mathbf{R}_{+\phi} \cdot \mathbf{r} = (x \cos\phi - y \sin\phi) \hat{\mathbf{e}}_x + (x \sin\phi + y \cos\phi) \hat{\mathbf{e}}_y$$
$$\begin{pmatrix} \cos\phi & -\sin\phi \\ \sin\phi & \cos\phi \end{pmatrix} \cdot \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \cos\phi - y \sin\phi \\ x \sin\phi + y \cos\phi \end{pmatrix}$$

### 5. Complex products provide 2D “dot”(•) and “cross”(×) products.

Two complex numbers  $A = A_x + iA_y$  and  $B = B_x + iB_y$  and their “star” (\*)-product  $A * B$ .

$$A * B = (A_x + iA_y)^* (B_x + iB_y) = (A_x - iA_y)(B_x + iB_y)$$
$$= (A_x B_x + A_y B_y) + i(A_x B_y - A_y B_x) = \mathbf{A} \cdot \mathbf{B} + i |\mathbf{A} \times \mathbf{B}|_{Z \perp(x,y)}$$

Real part is scalar or “dot”(•) product  $\mathbf{A} \cdot \mathbf{B}$ .

Imaginary part is vector or “cross”(×) product, but just the Z-component *normal* to xy-plane.

Rewrite  $A * B$  in polar form.

$$A * B = (|A| e^{i\theta_A})^* (|B| e^{i\theta_B}) = |A| e^{-i\theta_A} |B| e^{i\theta_B} = |A| |B| e^{i(\theta_B - \theta_A)}$$
$$= |A| |B| \cos(\theta_B - \theta_A) + i |A| |B| \sin(\theta_B - \theta_A) = \mathbf{A} \cdot \mathbf{B} + i |\mathbf{A} \times \mathbf{B}|_{Z \perp(x,y)}$$

## What Good Are Complex Exponentials? (contd.)

### 4. Complex products provide 2D rotation operations.

$$e^{i\phi} \cdot z = (\cos\phi + i \sin\phi) \cdot (x + iy) = x \cos\phi - y \sin\phi + i(x \sin\phi + y \cos\phi)$$

$$\mathbf{R}_{+\phi} \cdot \mathbf{r} = (x \cos\phi - y \sin\phi) \hat{\mathbf{e}}_x + (x \sin\phi + y \cos\phi) \hat{\mathbf{e}}_y$$

$$\begin{pmatrix} \cos\phi & -\sin\phi \\ \sin\phi & \cos\phi \end{pmatrix} \cdot \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \cos\phi - y \sin\phi \\ x \sin\phi + y \cos\phi \end{pmatrix}$$

### 5. Complex products provide 2D “dot”(•) and “cross”(×) products.

Two complex numbers  $A = A_x + iA_y$  and  $B = B_x + iB_y$  and their “star” (\*)-product  $A * B$ .

$$A * B = (A_x + iA_y)^* (B_x + iB_y) = (A_x - iA_y)(B_x + iB_y)$$

$$= (A_x B_x + A_y B_y) + i(A_x B_y - A_y B_x) = \mathbf{A} \cdot \mathbf{B} + i |\mathbf{A} \times \mathbf{B}|_{Z \perp(x,y)}$$

Real part is scalar or “dot”(•) product  $\mathbf{A} \cdot \mathbf{B}$ .

Imaginary part is vector or “cross”(×) product, but just the Z-component normal to xy-plane.

Rewrite  $A * B$  in polar form.

$$A * B = (|A| e^{i\theta_A})^* (|B| e^{i\theta_B}) = |A| e^{-i\theta_A} |B| e^{i\theta_B} = |A| |B| e^{i(\theta_B - \theta_A)}$$

$$= |A| |B| \cos(\theta_B - \theta_A) + i |A| |B| \sin(\theta_B - \theta_A) = \mathbf{A} \cdot \mathbf{B} + i |\mathbf{A} \times \mathbf{B}|_{Z \perp(x,y)}$$

$$\mathbf{A} \cdot \mathbf{B} = |A| |B| \cos(\theta_B - \theta_A)$$

$$= |A| \cos\theta_A |B| \cos\theta_B + |A| \sin\theta_A |B| \sin\theta_B$$

$$= A_x B_x + A_y B_y$$

$$|\mathbf{A} \times \mathbf{B}| = |A| |B| \sin(\theta_B - \theta_A)$$

$$= |A| \cos\theta_A |B| \sin\theta_B - |A| \sin\theta_A |B| \cos\theta_B$$

$$= A_x B_y - A_y B_x$$



## *What Good are complex variables?*

*Easy 2D vector calculus*

*Easy 2D vector derivatives*

*Easy 2D source-free field theory*

*Easy 2D vector field-potential theory*



## What Good Are Complex Exponentials? (contd.)

### 6. Complex derivative contains “divergence” ( $\nabla \cdot \mathbf{F}$ ) and “curl” ( $\nabla \times \mathbf{F}$ ) of 2D vector field

Relation of  $(z, z^*)$  to  $(x = \operatorname{Re}z, y = \operatorname{Im}z)$  defines a  $z$ -derivative  $\frac{df}{dz}$  and “star”  $z^*$ -derivative.  $\frac{df}{dz^*}$

$$z = x + iy$$

$$x = \frac{1}{2}(z + z^*)$$

$$z^* = x - iy$$

$$y = \frac{1}{2i}(z - z^*)$$

Applying  
chain-rule

$$\frac{df}{dz} = \frac{\partial x}{\partial z} \frac{\partial f}{\partial x} + \frac{\partial y}{\partial z} \frac{\partial f}{\partial y} = \frac{1}{2} \frac{\partial f}{\partial x} - \frac{i}{2} \frac{\partial f}{\partial y}$$

$$\frac{d}{dz} = \frac{1}{2} \frac{\partial}{\partial x} - \frac{i}{2} \frac{\partial}{\partial y}$$

$$\frac{df}{dz^*} = \frac{\partial x}{\partial z^*} \frac{\partial f}{\partial x} + \frac{\partial y}{\partial z^*} \frac{\partial f}{\partial y} = \frac{1}{2} \frac{\partial f}{\partial x} + \frac{i}{2} \frac{\partial f}{\partial y}$$

$$\frac{d}{dz^*} = \frac{1}{2} \frac{\partial}{\partial x} + \frac{i}{2} \frac{\partial}{\partial y}$$

## What Good Are Complex Exponentials? (contd.)

### 6. Complex derivative contains “divergence” ( $\nabla \cdot \mathbf{F}$ ) and “curl” ( $\nabla \times \mathbf{F}$ ) of 2D vector field

Relation of  $(z, z^*)$  to  $(x = \text{Re}z, y = \text{Im}z)$  defines a  $z$ -derivative  $\frac{df}{dz}$  and “star”  $z^*$ -derivative.  $\frac{df}{dz^*}$

$$z = x + iy$$

$$x = \frac{1}{2}(z + z^*)$$

$$z^* = x - iy$$

$$y = \frac{1}{2i}(z - z^*)$$

$$\frac{df}{dz} = \frac{\partial x}{\partial z} \frac{\partial f}{\partial x} + \frac{\partial y}{\partial z} \frac{\partial f}{\partial y} = \frac{1}{2} \frac{\partial f}{\partial x} - \frac{i}{2} \frac{\partial f}{\partial y}$$

$$\frac{d}{dz} = \frac{1}{2} \frac{\partial}{\partial x} - \frac{i}{2} \frac{\partial}{\partial y}$$

$$\frac{df}{dz^*} = \frac{\partial x}{\partial z^*} \frac{\partial f}{\partial x} + \frac{\partial y}{\partial z^*} \frac{\partial f}{\partial y} = \frac{1}{2} \frac{\partial f}{\partial x} + \frac{i}{2} \frac{\partial f}{\partial y}$$

$$\frac{d}{dz^*} = \frac{1}{2} \frac{\partial}{\partial x} + \frac{i}{2} \frac{\partial}{\partial y}$$

$$\frac{d}{dz} = \frac{1}{2} \frac{\partial}{\partial x} - \frac{i}{2} \frac{\partial}{\partial y}$$

Derivative chain-rule shows real part of  $\frac{df}{dz}$  has 2D divergence  $\nabla \cdot \mathbf{f}$  and imaginary part has curl  $\nabla \times \mathbf{f}$ .

$$\frac{df}{dz} = \frac{d}{dz} (f_x + i f_y) = \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) (f_x + i f_y) = \frac{1}{2} \left( \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial f_y}{\partial x} - \frac{\partial f_x}{\partial y} \right) = \frac{1}{2} \nabla \cdot \mathbf{f} + \frac{i}{2} |\nabla \times \mathbf{f}|_{Z \perp (x,y)}$$

## What Good Are Complex Exponentials? (contd.)

### 6. Complex derivative contains “divergence” ( $\nabla \cdot \mathbf{F}$ ) and “curl” ( $\nabla \times \mathbf{F}$ ) of 2D vector field

Relation of  $(z, z^*)$  to  $(x = \text{Re}z, y = \text{Im}z)$  defines a  $z$ -derivative  $\frac{df}{dz}$  and “star”  $z^*$ -derivative.  $\frac{df}{dz^*}$

$$z = x + iy$$

$$x = \frac{1}{2}(z + z^*)$$

$$z^* = x - iy$$

$$y = \frac{1}{2i}(z - z^*)$$

$$\frac{df}{dz} = \frac{\partial x}{\partial z} \frac{\partial f}{\partial x} + \frac{\partial y}{\partial z} \frac{\partial f}{\partial y} = \frac{1}{2} \frac{\partial f}{\partial x} - \frac{i}{2} \frac{\partial f}{\partial y}$$

$$\frac{d}{dz} = \frac{1}{2} \frac{\partial}{\partial x} - \frac{i}{2} \frac{\partial}{\partial y}$$

$$\frac{df}{dz^*} = \frac{\partial x}{\partial z^*} \frac{\partial f}{\partial x} + \frac{\partial y}{\partial z^*} \frac{\partial f}{\partial y} = \frac{1}{2} \frac{\partial f}{\partial x} + \frac{i}{2} \frac{\partial f}{\partial y}$$

$$\frac{d}{dz^*} = \frac{1}{2} \frac{\partial}{\partial x} + \frac{i}{2} \frac{\partial}{\partial y}$$

$$\frac{d}{dz} = \frac{1}{2} \frac{\partial}{\partial x} - \frac{i}{2} \frac{\partial}{\partial y}$$

Derivative chain-rule shows real part of  $\frac{df}{dz}$  has 2D divergence  $\nabla \cdot \mathbf{f}$  and imaginary part has curl  $\nabla \times \mathbf{f}$ .

$$\frac{df}{dz} = \frac{d}{dz} (f_x + i f_y) = \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) (f_x + i f_y) = \frac{1}{2} \left( \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial f_y}{\partial x} - \frac{\partial f_x}{\partial y} \right) = \frac{1}{2} \nabla \cdot \mathbf{f} + \frac{i}{2} |\nabla \times \mathbf{f}|_{Z \perp (x,y)}$$

### 7. Invent source-free 2D vector fields [ $\nabla \cdot \mathbf{F} = 0$ and $\nabla \times \mathbf{F} = 0$ ]

We can invent *source-free 2D vector fields* that are both *zero-divergence* and *zero-curl*.

Take any function  $f(z)$ , conjugate it (change all  $i$ 's to  $-i$ ) to give  $f^*(z^*)$  for which  $\frac{df^*}{dz^*} = 0$ .

## What Good Are Complex Exponentials? (contd.)

### 6. Complex derivative contains “divergence” ( $\nabla \cdot \mathbf{F}$ ) and “curl” ( $\nabla \times \mathbf{F}$ ) of 2D vector field

Relation of  $(z, z^*)$  to  $(x = \text{Re}z, y = \text{Im}z)$  defines a  $z$ -derivative  $\frac{df}{dz}$  and “star”  $z^*$ -derivative.  $\frac{df}{dz^*}$

$$z = x + iy$$

$$x = \frac{1}{2}(z + z^*)$$

$$z^* = x - iy$$

$$y = \frac{1}{2i}(z - z^*)$$

$$\frac{df}{dz} = \frac{\partial x}{\partial z} \frac{\partial f}{\partial x} + \frac{\partial y}{\partial z} \frac{\partial f}{\partial y} = \frac{1}{2} \frac{\partial f}{\partial x} - \frac{i}{2} \frac{\partial f}{\partial y}$$

$$\frac{d}{dz} = \frac{1}{2} \frac{\partial}{\partial x} - \frac{i}{2} \frac{\partial}{\partial y}$$

$$\frac{df}{dz^*} = \frac{\partial x}{\partial z^*} \frac{\partial f}{\partial x} + \frac{\partial y}{\partial z^*} \frac{\partial f}{\partial y} = \frac{1}{2} \frac{\partial f}{\partial x} + \frac{i}{2} \frac{\partial f}{\partial y}$$

$$\frac{d}{dz^*} = \frac{1}{2} \frac{\partial}{\partial x} + \frac{i}{2} \frac{\partial}{\partial y}$$

$$\frac{d}{dz} = \frac{1}{2} \frac{\partial}{\partial x} - \frac{i}{2} \frac{\partial}{\partial y}$$

Derivative chain-rule shows real part of  $\frac{df}{dz}$  has 2D divergence  $\nabla \cdot \mathbf{f}$  and imaginary part has curl  $\nabla \times \mathbf{f}$ .

$$\frac{df}{dz} = \frac{d}{dz} (f_x + i f_y) = \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) (f_x + i f_y) = \frac{1}{2} \left( \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial f_y}{\partial x} - \frac{\partial f_x}{\partial y} \right) = \frac{1}{2} \nabla \cdot \mathbf{f} + \frac{i}{2} |\nabla \times \mathbf{f}|_{Z \perp(x,y)}$$

### 7. Invent source-free 2D vector fields [ $\nabla \cdot \mathbf{F} = 0$ and $\nabla \times \mathbf{F} = 0$ ]

We can invent *source-free 2D vector fields* that are both *zero-divergence* and *zero-curl*.

Take any function  $f(z)$ , conjugate it (change all  $i$ 's to  $-i$ ) to give  $f^*(z^*)$  for which  $\frac{df^*}{dz^*} = 0$ .

For example: if  $f(z) = a \cdot z$  then  $f^*(z^*) = a \cdot z^* = a(x - iy)$  is not function of  $z$  so it has zero  $z$ -derivative.

$\mathbf{F} = (F_x, F_y) = (f_x^*, f_y^*) = (a \cdot x, -a \cdot y)$  has *zero divergence*:  $\nabla \cdot \mathbf{F} = 0$  and has *zero curl*:  $|\nabla \times \mathbf{F}| = 0$ .

$$\nabla \cdot \mathbf{F} = \frac{\partial F_x}{\partial x} + \frac{\partial F_y}{\partial y} = \frac{\partial(ax)}{\partial x} + \frac{\partial(-ay)}{\partial y} = 0 \quad |\nabla \times \mathbf{F}|_{Z \perp(x,y)} = \frac{\partial F_y}{\partial x} - \frac{\partial F_x}{\partial y} = \frac{\partial(-ay)}{\partial x} - \frac{\partial(ax)}{\partial y} = 0$$

A *DFL* field  $\mathbf{F}$  (*Divergence-Free-Laminar*)

## What Good Are Complex Exponentials? (contd.)


### 7. Invent source-free 2D vector fields [ $\nabla \cdot \mathbf{F} = 0$ and $\nabla \times \mathbf{F} = 0$ ]

We can invent *source-free 2D vector fields* that are both *zero-divergence* and *zero-curl*. Take any function  $f(z)$ , conjugate it (change all  $i$ 's to  $-i$ ) to give  $f^*(z^*)$  for which

For example: if  $f(z) = a \cdot z$  then  $f^*(z^*) = a \cdot z^* = a(x - iy)$  is not function of  $z$  so it has zero  $z$ -derivative.

$\mathbf{F} = (F_x, F_y) = (f_x^*, f_y^*) = (a \cdot x, -a \cdot y)$  has *zero divergence*:  $\nabla \cdot \mathbf{F} = 0$  and has *zero curl*:  $|\nabla \times \mathbf{F}| = 0$ .

$$\nabla \cdot \mathbf{F} = \frac{\partial F_x}{\partial x} + \frac{\partial F_y}{\partial y} = \frac{\partial(ax)}{\partial x} + \frac{\partial(-ay)}{\partial y} = 0 \quad |\nabla \times \mathbf{F}|_{z \perp (x,y)} = \frac{\partial F_y}{\partial x} - \frac{\partial F_x}{\partial y} = \frac{\partial(-ay)}{\partial x} - \frac{\partial(ax)}{\partial y} = 0$$



precursor to  
Unit 1  
Fig. 10.7

$\mathbf{F} = (f_x^*, f_y^*) = (a \cdot x, -a \cdot y)$  is a *divergence-free laminar (DFL)* field.

## *What Good are complex variables?*

*Easy 2D vector calculus*

*Easy 2D vector derivatives*

*Easy 2D source-free field theory*



*Easy 2D vector field-potential theory*

## What Good Are Complex Exponentials? (contd.)

8. Complex potential  $\phi$  contains “scalar” ( $\mathbf{F}=\nabla\Phi$ ) and “vector” ( $\mathbf{F}=\nabla\times\mathbf{A}$ ) potentials

Any *DFL* field  $\mathbf{F}$  is a gradient of a *scalar potential field*  $\Phi$  or a curl of a *vector potential field*  $\mathbf{A}$ .

$$\mathbf{F} = \nabla\Phi$$

$$\mathbf{F} = \nabla\times\mathbf{A}$$

A *complex potential*  $\phi(z) = \Phi(x,y) + i\mathbf{A}(x,y)$  exists whose  $z$ -derivative is  $f(z) = d\phi/dz$ .

Its complex conjugate  $\phi^*(z^*) = \Phi(x,y) - i\mathbf{A}(x,y)$  has  $z^*$ -derivative  $f^*(z^*) = d\phi^*/dz^*$  giving *DFL* field  $\mathbf{F}$ .



## What Good Are Complex Exponentials? (contd.)

8. Complex potential  $\phi$  contains “scalar” ( $\mathbf{F}=\nabla\Phi$ ) and “vector” ( $\mathbf{F}=\nabla\times\mathbf{A}$ ) potentials

Any *DFL* field  $\mathbf{F}$  is a gradient of a *scalar potential field*  $\Phi$  or a curl of a *vector potential field*  $\mathbf{A}$ .

$$\mathbf{F} = \nabla\Phi$$

$$\mathbf{F} = \nabla\times\mathbf{A}$$

A *complex potential*  $\phi(z) = \Phi(x,y) + i\mathbf{A}(x,y)$  exists whose  $z$ -derivative is  $f(z) = d\phi/dz$ .

Its complex conjugate  $\phi^*(z^*) = \Phi(x,y) - i\mathbf{A}(x,y)$  has  $z^*$ -derivative  $f^*(z^*) = d\phi^*/dz^*$  giving *DFL* field  $\mathbf{F}$ .

To find  $\phi = \Phi + i\mathbf{A}$  integrate  $f(z) = a \cdot z$  to get  $\phi$  and isolate real ( $\text{Re } \phi = \Phi$ ) and imaginary ( $\text{Im } \phi = \mathbf{A}$ ) parts.

## What Good Are Complex Exponentials? (contd.)

8. Complex potential  $\phi$  contains “scalar” ( $\mathbf{F}=\nabla\Phi$ ) and “vector” ( $\mathbf{F}=\nabla\times\mathbf{A}$ ) potentials

Any *DFL* field  $\mathbf{F}$  is a gradient of a *scalar potential field*  $\Phi$  or a curl of a *vector potential field*  $\mathbf{A}$ .

$$\mathbf{F} = \nabla\Phi$$

$$\mathbf{F} = \nabla\times\mathbf{A}$$

A *complex potential*  $\phi(z)=\Phi(x,y)+i\mathbf{A}(x,y)$  exists whose  $z$ -derivative is  $f(z)=d\phi/dz$ .

Its complex conjugate  $\phi^*(z^*)=\Phi(x,y)-i\mathbf{A}(x,y)$  has  $z^*$ -derivative  $f^*(z^*)=d\phi^*/dz^*$  giving *DFL* field  $\mathbf{F}$ .

To find  $\phi=\Phi+i\mathbf{A}$  integrate  $f(z)=a\cdot z$  to get  $\phi$  and isolate real ( $\text{Re } \phi =\Phi$ ) and imaginary ( $\text{Im } \phi =\mathbf{A}$ ) parts.

$$f(z)=\frac{d\phi}{dz} \Rightarrow \phi = \Phi + i \mathbf{A} = \int f \cdot dz = \int az \cdot dz = \frac{1}{2} az^2$$

## What Good Are Complex Exponentials? (contd.)

8. Complex potential  $\phi$  contains “scalar” ( $\mathbf{F}=\nabla\Phi$ ) and “vector” ( $\mathbf{F}=\nabla\times\mathbf{A}$ ) potentials

Any *DFL* field  $\mathbf{F}$  is a gradient of a *scalar potential field*  $\Phi$  or a curl of a *vector potential field*  $\mathbf{A}$ .

$$\mathbf{F} = \nabla\Phi$$

$$\mathbf{F} = \nabla\times\mathbf{A}$$

A *complex potential*  $\phi(z)=\Phi(x,y)+i\mathbf{A}(x,y)$  exists whose  $z$ -derivative is  $f(z)=d\phi/dz$ .

Its complex conjugate  $\phi^*(z^*)=\Phi(x,y)-i\mathbf{A}(x,y)$  has  $z^*$ -derivative  $f^*(z^*)=d\phi^*/dz^*$  giving *DFL* field  $\mathbf{F}$ .

To find  $\phi=\Phi+i\mathbf{A}$  integrate  $f(z)=a\cdot z$  to get  $\phi$  and isolate real ( $\text{Re } \phi =\Phi$ ) and imaginary ( $\text{Im } \phi =\mathbf{A}$ ) parts.

$$f(z)=\frac{d\phi}{dz} \Rightarrow \phi = \underbrace{\Phi}_{\frac{1}{2} a(x^2 - y^2)} + i \underbrace{\mathbf{A}}_{axy} = \int f \cdot dz = \int az \cdot dz = \frac{1}{2} az^2 = \frac{1}{2} a(x+iy)^2$$

## What Good Are Complex Exponentials? (contd.)

### 8. Complex potential $\phi$ contains “scalar” ( $\mathbf{F}=\nabla\Phi$ ) and “vector” ( $\mathbf{F}=\nabla\times\mathbf{A}$ ) potentials

Any *DFL* field  $\mathbf{F}$  is a gradient of a *scalar potential field*  $\Phi$  or a curl of a *vector potential field*  $\mathbf{A}$ .

$$\mathbf{F} = \nabla\Phi \qquad \mathbf{F} = \nabla\times\mathbf{A}$$


A *complex potential*  $\phi(z) = \Phi(x,y) + i\mathbf{A}(x,y)$  exists whose  $z$ -derivative is  $f(z) = d\phi/dz$ .

Its complex conjugate  $\phi^*(z^*) = \Phi(x,y) - i\mathbf{A}(x,y)$  has  $z^*$ -derivative  $f^*(z^*) = d\phi^*/dz^*$  giving *DFL* field  $\mathbf{F}$ .

To find  $\phi = \Phi + i\mathbf{A}$  integrate  $f(z) = a \cdot z$  to get  $\phi$  and isolate real ( $\text{Re } \phi = \Phi$ ) and imaginary ( $\text{Im } \phi = \mathbf{A}$ ) parts.

$$f(z) = \frac{d\phi}{dz} \Rightarrow \phi = \underbrace{\Phi}_{\frac{1}{2} a(x^2 - y^2)} + i \underbrace{\mathbf{A}}_{axy} = \int f \cdot dz = \int az \cdot dz = \frac{1}{2} az^2 = \frac{1}{2} a(x + iy)^2$$

Unit 1  
Fig. 10.7



## What Good Are Complex Exponentials? (contd.)

### 8. Complex potential $\phi$ contains “scalar” ( $\mathbf{F}=\nabla\Phi$ ) and “vector” ( $\mathbf{F}=\nabla\times\mathbf{A}$ ) potentials

Any *DFL* field  $\mathbf{F}$  is a gradient of a *scalar potential field*  $\Phi$  or a curl of a *vector potential field*  $\mathbf{A}$ .

$$\mathbf{F} = \nabla\Phi \qquad \mathbf{F} = \nabla\times\mathbf{A}$$


A *complex potential*  $\phi(z) = \Phi(x,y) + i\mathbf{A}(x,y)$  exists whose  $z$ -derivative is  $f(z) = d\phi/dz$ .

Its complex conjugate  $\phi^*(z^*) = \Phi(x,y) - i\mathbf{A}(x,y)$  has  $z^*$ -derivative  $f^*(z^*) = d\phi^*/dz^*$  giving *DFL* field  $\mathbf{F}$ .

To find  $\phi = \Phi + i\mathbf{A}$  integrate  $f(z) = a \cdot z$  to get  $\phi$  and isolate real ( $\text{Re } \phi = \Phi$ ) and imaginary ( $\text{Im } \phi = \mathbf{A}$ ) parts.

$$f(z) = \frac{d\phi}{dz} \Rightarrow \phi = \underbrace{\Phi}_{\frac{1}{2}a(x^2 - y^2)} + i \underbrace{\mathbf{A}}_{axy} = \int f \cdot dz = \int az \cdot dz = \frac{1}{2} az^2 = \frac{1}{2} a(x + iy)^2$$

*BONUS!*  
Get a free  
coordinate  
system!



Unit 1  
Fig. 10.7

Field:  
 $f^*(z^*) = z^* = x - iy$   
 $\mathbf{F}(x,y) = (x, -y)$   
Potential:  
 $\phi(z) = z^2$   
 $= x^2 - y^2 + i2xy$   
 $= \Phi + i\mathbf{A}$

The  $(\Phi, \mathbf{A})$  grid is a GCC coordinate system\*:

$$q^1 = \Phi = (x^2 - y^2)/2 = \text{const.}$$

$$q^2 = \mathbf{A} = (xy) = \text{const.}$$

\*Actually it's OCC.


# *What Good are complex variables?*

*Easy 2D vector calculus*

*Easy 2D vector derivatives*

*Easy 2D source-free field theory*

 *Easy 2D vector field-potential theory*

 The *half-n'-half* results: (Riemann-Cauchy Derivative Relations)

## What Good Are Complex Exponentials? (contd.)

8. (contd.) Complex potential  $\phi$  contains “scalar” ( $\mathbf{F} = \nabla \Phi$ ) and “vector” ( $\mathbf{F} = \nabla \times \mathbf{A}$ ) potentials  
 ...and either one (or *half-n'-half!*) works just as well.

Derivative  $\frac{d\phi^*}{dz^*}$  has 2D gradient  $\nabla \Phi = \begin{pmatrix} \frac{\partial \Phi}{\partial x} \\ \frac{\partial \Phi}{\partial y} \end{pmatrix}$  of scalar  $\Phi$  and curl  $\nabla \times \mathbf{A} = \begin{pmatrix} \frac{\partial \mathbf{A}}{\partial y} \\ -\frac{\partial \mathbf{A}}{\partial x} \end{pmatrix}$  of vector  $\mathbf{A}$  (and they're equal!)

$$f(z) = \frac{d\phi}{dz} \Rightarrow$$

$$\frac{d}{dz^*} \phi^* = \frac{d}{dz^*} (\Phi - i\mathbf{A}) = \frac{1}{2} \left( \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right) (\Phi - i\mathbf{A}) = \frac{1}{2} \left( \frac{\partial \Phi}{\partial x} + i \frac{\partial \Phi}{\partial y} \right) + \frac{1}{2} \left( \frac{\partial \mathbf{A}}{\partial y} - i \frac{\partial \mathbf{A}}{\partial x} \right) = \frac{1}{2} \nabla \Phi + \frac{1}{2} \nabla \times \mathbf{A}$$

$$\frac{d}{dz} = \frac{1}{2} \frac{\partial}{\partial x} - \frac{i}{2} \frac{\partial}{\partial y}$$

$$\frac{d}{dz^*} = \frac{1}{2} \frac{\partial}{\partial x} + \frac{i}{2} \frac{\partial}{\partial y}$$



## What Good Are Complex Exponentials? (contd.)

8. (contd.) Complex potential  $\phi$  contains “scalar” ( $\mathbf{F} = \nabla \Phi$ ) and “vector” ( $\mathbf{F} = \nabla \times \mathbf{A}$ ) potentials  
...and either one (or *half-n'-half!*) works just as well.

Derivative  $\frac{d\phi^*}{dz^*}$  has 2D gradient  $\nabla \Phi = \begin{pmatrix} \frac{\partial \Phi}{\partial x} \\ \frac{\partial \Phi}{\partial y} \end{pmatrix}$  of scalar  $\Phi$  and curl  $\nabla \times \mathbf{A} = \begin{pmatrix} \frac{\partial \mathbf{A}}{\partial y} \\ -\frac{\partial \mathbf{A}}{\partial x} \end{pmatrix}$  of vector  $\mathbf{A}$  (and they're equal!)

$$f(z) = \frac{d\phi}{dz} \Rightarrow$$

$$\frac{d}{dz^*} \phi^* = \frac{d}{dz^*} (\Phi - i\mathbf{A}) = \frac{1}{2} \left( \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right) (\Phi - i\mathbf{A}) = \frac{1}{2} \left( \frac{\partial \Phi}{\partial x} + i \frac{\partial \Phi}{\partial y} \right) + \frac{1}{2} \left( \frac{\partial \mathbf{A}}{\partial y} - i \frac{\partial \mathbf{A}}{\partial x} \right) = \frac{1}{2} \nabla \Phi + \frac{1}{2} \nabla \times \mathbf{A}$$

Note, *mathematician definition* of force field  $\mathbf{F} = +\nabla \Phi$  replaces usual physicist's definition  $\mathbf{F} = -\nabla \Phi$



## What Good Are Complex Exponentials? (contd.)

8. (contd.) Complex potential  $\phi$  contains “scalar” ( $\mathbf{F} = \nabla \Phi$ ) and “vector” ( $\mathbf{F} = \nabla \times \mathbf{A}$ ) potentials  
 ...and either one (or *half-n'-half!*) works just as well.

Derivative  $\frac{d\phi^*}{dz^*}$  has 2D gradient  $\nabla \Phi = \begin{pmatrix} \frac{\partial \Phi}{\partial x} \\ \frac{\partial \Phi}{\partial y} \end{pmatrix}$  of scalar  $\Phi$  and curl  $\nabla \times \mathbf{A} = \begin{pmatrix} \frac{\partial \mathbf{A}}{\partial y} \\ -\frac{\partial \mathbf{A}}{\partial x} \end{pmatrix}$  of vector  $\mathbf{A}$  (and they're equal!)

$$f(z) = \frac{d\phi}{dz} \Rightarrow$$

$$\frac{d}{dz^*} \phi^* = \frac{d}{dz^*} (\Phi - i\mathbf{A}) = \frac{1}{2} \left( \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right) (\Phi - i\mathbf{A}) = \frac{1}{2} \left( \frac{\partial \Phi}{\partial x} + i \frac{\partial \Phi}{\partial y} \right) + \frac{1}{2} \left( \frac{\partial \mathbf{A}}{\partial y} - i \frac{\partial \mathbf{A}}{\partial x} \right) = \frac{1}{2} \nabla \Phi + \frac{1}{2} \nabla \times \mathbf{A}$$

Note, *mathematician definition* of force field  $\mathbf{F} = +\nabla \Phi$  replaces usual physicist's definition  $\mathbf{F} = -\nabla \Phi$

Given  $\phi$ :

$$\begin{aligned} \phi &= \Phi + i\mathbf{A} \\ &= \frac{1}{2} a(x^2 - y^2) + i axy \end{aligned}$$

*The half-n'-half result*

find:

$$\nabla \Phi = \begin{pmatrix} \frac{\partial \Phi}{\partial x} \\ \frac{\partial \Phi}{\partial y} \end{pmatrix} = \begin{pmatrix} \frac{\partial}{\partial x} \frac{a}{2} (x^2 - y^2) \\ \frac{\partial}{\partial y} \frac{a}{2} (x^2 - y^2) \end{pmatrix} = \begin{pmatrix} ax \\ -ay \end{pmatrix} = \mathbf{F}$$

or find:

$$\nabla \times \mathbf{A} = \begin{pmatrix} \frac{\partial \mathbf{A}}{\partial y} \\ -\frac{\partial \mathbf{A}}{\partial x} \end{pmatrix} = \begin{pmatrix} \frac{\partial}{\partial y} axy \\ -\frac{\partial}{\partial x} axy \end{pmatrix} = \begin{pmatrix} ax \\ -ay \end{pmatrix} = \mathbf{F}$$

## What Good Are Complex Exponentials? (contd.)

8. (contd.) Complex potential  $\phi$  contains “scalar” ( $\mathbf{F}=\nabla\Phi$ ) and “vector” ( $\mathbf{F}=\nabla\times\mathbf{A}$ ) potentials  
 ...and either one (or *half-n'-half!*) works just as well.

Derivative  $\frac{d\phi^*}{dz^*}$  has 2D gradient  $\nabla\Phi = \begin{pmatrix} \frac{\partial\Phi}{\partial x} \\ \frac{\partial\Phi}{\partial y} \end{pmatrix}$  of scalar  $\Phi$  and curl  $\nabla\times\mathbf{A} = \begin{pmatrix} \frac{\partial\mathbf{A}}{\partial y} \\ -\frac{\partial\mathbf{A}}{\partial x} \end{pmatrix}$  of vector  $\mathbf{A}$  (and they're equal!)

$$f(z) = \frac{d\phi}{dz} \Rightarrow$$


$$\frac{d}{dz^*} \phi^* = \frac{d}{dz^*} (\Phi - i\mathbf{A}) = \frac{1}{2} \left( \frac{\partial}{\partial x} + i\frac{\partial}{\partial y} \right) (\Phi - i\mathbf{A}) = \frac{1}{2} \left( \frac{\partial\Phi}{\partial x} + i\frac{\partial\Phi}{\partial y} \right) + \frac{1}{2} \left( \frac{\partial\mathbf{A}}{\partial y} - i\frac{\partial\mathbf{A}}{\partial x} \right) = \frac{1}{2} \nabla\Phi + \frac{1}{2} \nabla\times\mathbf{A}$$

Note, *mathematician definition* of force field  $\mathbf{F} = +\nabla\Phi$  replaces usual physicist's definition  $\mathbf{F} = -\nabla\Phi$

Given  $\phi$ :  $\phi = \Phi + i\mathbf{A} = \frac{1}{2} a(x^2 - y^2) + i axy$  The *half-n'-half* result

find:  $\nabla\Phi = \begin{pmatrix} \frac{\partial\Phi}{\partial x} \\ \frac{\partial\Phi}{\partial y} \end{pmatrix} = \begin{pmatrix} \frac{\partial}{\partial x} \frac{a}{2}(x^2 - y^2) \\ \frac{\partial}{\partial y} \frac{a}{2}(x^2 - y^2) \end{pmatrix} = \begin{pmatrix} ax \\ -ay \end{pmatrix} = \mathbf{F}$  or find:  $\nabla\times\mathbf{A} = \begin{pmatrix} \frac{\partial\mathbf{A}}{\partial y} \\ -\frac{\partial\mathbf{A}}{\partial x} \end{pmatrix} = \begin{pmatrix} \frac{\partial}{\partial y} axy \\ -\frac{\partial}{\partial x} axy \end{pmatrix} = \begin{pmatrix} ax \\ -ay \end{pmatrix} = \mathbf{F}$

Scalar *static potential lines*  $\Phi = \text{const.}$  and vector *flux potential lines*  $\mathbf{A} = \text{const.}$  define *DFL field-net*.



# What Good Are Complex Exponentials? (contd.)

8. (contd.) Complex potential  $\phi$  contains “scalar” ( $\mathbf{F}=\nabla\Phi$ ) and “vector” ( $\mathbf{F}=\nabla\times\mathbf{A}$ ) potentials  
 ...and either one (or *half-n'-half!*) works just as well.

Derivative  $\frac{d\phi^*}{dz^*}$  has 2D gradient  $\nabla\Phi = \begin{pmatrix} \frac{\partial\Phi}{\partial x} \\ \frac{\partial\Phi}{\partial y} \end{pmatrix}$  of scalar  $\Phi$  and curl  $\nabla\times\mathbf{A} = \begin{pmatrix} \frac{\partial\mathbf{A}}{\partial y} \\ -\frac{\partial\mathbf{A}}{\partial x} \end{pmatrix}$  of vector  $\mathbf{A}$  (and they're equal!)

The *half-n'-half* result

$$\frac{d}{dz^*} \phi^* = \frac{d}{dz^*} (\Phi - i\mathbf{A}) = \frac{1}{2} \left( \frac{\partial}{\partial x} + i\frac{\partial}{\partial y} \right) (\Phi - i\mathbf{A}) = \frac{1}{2} \left( \frac{\partial\Phi}{\partial x} + i\frac{\partial\Phi}{\partial y} \right) + \frac{1}{2} \left( \frac{\partial\mathbf{A}}{\partial y} - i\frac{\partial\mathbf{A}}{\partial x} \right) = \frac{1}{2} \nabla\Phi + \frac{1}{2} \nabla\times\mathbf{A}$$

Note, *mathematician definition* of force field  $\mathbf{F} = +\nabla\Phi$  replaces usual physicist's definition  $\mathbf{F} = -\nabla\Phi$

Given  $\phi$ :

$$\phi = \Phi + i\mathbf{A} = \frac{1}{2} a(x^2 - y^2) + i axy$$

The *half-n'-half* result


find:

$$\nabla\Phi = \begin{pmatrix} \frac{\partial\Phi}{\partial x} \\ \frac{\partial\Phi}{\partial y} \end{pmatrix} = \begin{pmatrix} \frac{\partial}{\partial x} \frac{a}{2}(x^2 - y^2) \\ \frac{\partial}{\partial y} \frac{a}{2}(x^2 - y^2) \end{pmatrix} = \begin{pmatrix} ax \\ -ay \end{pmatrix} = \mathbf{F}$$

or find:

$$\nabla\times\mathbf{A} = \begin{pmatrix} \frac{\partial\mathbf{A}}{\partial y} \\ -\frac{\partial\mathbf{A}}{\partial x} \end{pmatrix} = \begin{pmatrix} \frac{\partial}{\partial y} axy \\ -\frac{\partial}{\partial x} axy \end{pmatrix} = \begin{pmatrix} ax \\ -ay \end{pmatrix} = \mathbf{F}$$

Scalar *static potential lines*  $\Phi = \text{const.}$  and vector *flux potential lines*  $\mathbf{A} = \text{const.}$  define *DFL field-net*.



The *half-n'-half* results

are called

*Riemann-Cauchy  
 Derivative Relations*

$$\frac{\partial\Phi}{\partial x} = \frac{\partial\mathbf{A}}{\partial y} \quad \text{is:} \quad \frac{\partial\text{Re}f(z)}{\partial x} = \frac{\partial\text{Im}f(z)}{\partial y}$$

$$\frac{\partial\Phi}{\partial y} = -\frac{\partial\mathbf{A}}{\partial x} \quad \text{is:} \quad \frac{\partial\text{Re}f(z)}{\partial y} = -\frac{\partial\text{Im}f(z)}{\partial x}$$

→ *4. Riemann-Cauchy conditions* *What's analytic? (...and what's not?)*

*Review  $(z, z^*)$  to  $(x, y)$  transformation relations*

$$z = x + iy$$

$$x = \frac{1}{2} (z + z^*)$$

$$\frac{df}{dz} = \frac{\partial x}{\partial z} \frac{\partial f}{\partial x} + \frac{\partial y}{\partial z} \frac{\partial f}{\partial y} = \frac{1}{2} \frac{\partial f}{\partial x} + \frac{1}{2i} \frac{\partial f}{\partial y} = \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) f$$

$$z^* = x - iy$$

$$y = \frac{1}{2i} (z - z^*)$$

$$\frac{df}{dz^*} = \frac{\partial x}{\partial z^*} \frac{\partial f}{\partial x} + \frac{\partial y}{\partial z^*} \frac{\partial f}{\partial y} = \frac{1}{2} \frac{\partial f}{\partial x} - \frac{1}{2i} \frac{\partial f}{\partial y} = \frac{1}{2} \left( \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right) f$$

*Criteria for a field function  $f = f_x(x, y) + i f_y(x, y)$  to be an **analytic function  $f(z)$**  of  $z = x + iy$ :*

*First,  $f(z)$  must not be a function of  $z^* = x - iy$ , that is:  $\frac{df}{dz^*} = 0$*

*This implies  $f(z)$  satisfies differential equations known as the **Riemann-Cauchy conditions***

$$\frac{df}{dz^*} = 0 = \frac{1}{2} \left( \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right) (f_x + i f_y) = \frac{1}{2} \left( \frac{\partial f_x}{\partial x} - \frac{\partial f_y}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial f_y}{\partial x} + \frac{\partial f_x}{\partial y} \right) \text{ implies: } \frac{\partial f_x}{\partial x} = \frac{\partial f_y}{\partial y} \quad \text{and:} \quad \frac{\partial f_y}{\partial x} = -\frac{\partial f_x}{\partial y}$$

$$\frac{df}{dz} = \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) (f_x + i f_y) = \frac{1}{2} \left( \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial f_y}{\partial x} - \frac{\partial f_x}{\partial y} \right) = \frac{\partial f_x}{\partial x} + i \frac{\partial f_y}{\partial x} = \frac{\partial f_y}{\partial y} - i \frac{\partial f_x}{\partial y} = \frac{\partial}{\partial x} (f_x + i f_y) = \frac{\partial}{\partial iy} (f_x + i f_y)$$

*Review (z,z\*) to (x,y) transformation relations*

$$z = x + iy$$

$$x = \frac{1}{2} (z + z^*)$$

$$\frac{df}{dz} = \frac{\partial x}{\partial z} \frac{\partial f}{\partial x} + \frac{\partial y}{\partial z} \frac{\partial f}{\partial y} = \frac{1}{2} \frac{\partial f}{\partial x} + \frac{1}{2i} \frac{\partial f}{\partial y} = \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) f$$

$$z^* = x - iy$$

$$y = \frac{1}{2i} (z - z^*)$$

$$\frac{df}{dz^*} = \frac{\partial x}{\partial z^*} \frac{\partial f}{\partial x} + \frac{\partial y}{\partial z^*} \frac{\partial f}{\partial y} = \frac{1}{2} \frac{\partial f}{\partial x} - \frac{1}{2i} \frac{\partial f}{\partial y} = \frac{1}{2} \left( \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right) f$$

*Criteria for a field function  $f = f_x(x,y) + i f_y(x,y)$  to be an **analytic function  $f(z)$**  of  $z=x+iy$ :*

*First,  $f(z)$  must not be a function of  $z^*=x-iy$ , that is:  $\frac{df}{dz^*} = 0$*

*This implies  $f(z)$  satisfies differential equations known as the **Riemann-Cauchy conditions***

$$\frac{df}{dz^*} = 0 = \frac{1}{2} \left( \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right) (f_x + i f_y) = \frac{1}{2} \left( \frac{\partial f_x}{\partial x} - \frac{\partial f_y}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial f_y}{\partial x} + \frac{\partial f_x}{\partial y} \right) \text{ implies: } \frac{\partial f_x}{\partial x} = \frac{\partial f_y}{\partial y} \quad \text{and:} \quad \frac{\partial f_y}{\partial x} = -\frac{\partial f_x}{\partial y}$$

$$\frac{df}{dz} = \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) (f_x + i f_y) = \frac{1}{2} \left( \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial f_y}{\partial x} - \frac{\partial f_x}{\partial y} \right) = \frac{\partial f_x}{\partial x} + i \frac{\partial f_y}{\partial x} = \frac{\partial f_y}{\partial y} - i \frac{\partial f_x}{\partial y} = \frac{\partial}{\partial x} (f_x + i f_y) = \frac{\partial}{\partial iy} (f_x + i f_y)$$

*Criteria for a field function  $f = f_x(x,y) + i f_y(x,y)$  to be an **analytic function  $f(z^*)$**  of  $z^*=x-iy$ :*

*First,  $f(z^*)$  must not be a function of  $z=x+iy$ , that is:  $\frac{df}{dz} = 0$*

*This implies  $f(z^*)$  satisfies differential equations we call **Anti-Riemann-Cauchy conditions***

$$\frac{df}{dz} = 0 = \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) (f_x + i f_y) = \frac{1}{2} \left( \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial f_y}{\partial x} - \frac{\partial f_x}{\partial y} \right) = \text{implies: } \frac{\partial f_x}{\partial x} = -\frac{\partial f_y}{\partial y} \quad \text{and:} \quad \frac{\partial f_y}{\partial x} = \frac{\partial f_x}{\partial y}$$

$$\frac{df}{dz^*} = \frac{1}{2} \left( \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right) (f_x + i f_y) = \frac{1}{2} \left( \frac{\partial f_x}{\partial x} - \frac{\partial f_y}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial f_y}{\partial x} + \frac{\partial f_x}{\partial y} \right) = \frac{\partial f_x}{\partial x} + i \frac{\partial f_y}{\partial x} = -\frac{\partial f_y}{\partial y} + i \frac{\partial f_x}{\partial y} = \frac{\partial}{\partial x} (f_x + i f_y) = -\frac{\partial}{\partial iy} (f_x + i f_y)$$

*What's analytic? (...and what's not?)*

Example: Is  $f(x,y) = 2x + iy$  an analytic function of  $z=x+iy$ ?

## *What's analytic? (...and what's not?)*

Example: Q: Is  $f(x,y) = 2x + i4y$  an analytic function of  $z=x+iy$ ?

Well, test it using definitions:  $z = x + iy$  and:  $z^* = x - iy$   
or:  $x = (z+z^*)/2$  and:  $y = -i(z-z^*)/2$



## *What's analytic? (...and what's not?)*

Example: Q: Is  $f(x,y) = 2x + i4y$  an analytic function of  $z=x+iy$ ?

Well, test it using definitions:  $z = x + iy$                       and:  $z^* = x - iy$   
    or:  $x = (z+z^*)/2$                       and:  $y = -i(z-z^*)/2$

$$f(x,y) = 2x + i4y = 2 (z+z^*)/2 + i4(-i(z-z^*)/2)$$

## *What's analytic? (...and what's not?)*

Example: Q: Is  $f(x,y) = 2x + i4y$  an analytic function of  $z=x+iy$ ?

Well, test it using definitions:  $z = x + iy$                       and:  $z^* = x - iy$   
or:  $x = (z+z^*)/2$                       and:  $y = -i(z-z^*)/2$

$$\begin{aligned} f(x,y) = 2x + i4y &= 2 \frac{(z+z^*)}{2} + i4 \frac{-i(z-z^*)}{2} \\ &= z+z^* + (2z-2z^*) \end{aligned}$$

## *What's analytic? (...and what's not?)*

Example: Q: Is  $f(x,y) = 2x + i4y$  an analytic function of  $z=x+iy$ ?

Well, test it using definitions:  $z = x + iy$                       and:  $z^* = x - iy$   
or:  $x = (z+z^*)/2$                       and:  $y = -i(z-z^*)/2$

$$\begin{aligned}f(x,y) = 2x + i4y &= 2 \frac{(z+z^*)}{2} + i4 \frac{-i(z-z^*)}{2} \\&= z+z^* + (2z-2z^*) \\&= 3z-z^*\end{aligned}$$

## *What's analytic? (...and what's not?)*

Example: Q: Is  $f(x,y) = 2x + i4y$  an analytic function of  $z=x+iy$ ?

Well, test it using definitions:  $z = x + iy$                       and:  $z^* = x - iy$   
or:  $x = (z+z^*)/2$                       and:  $y = -i(z-z^*)/2$

$$\begin{aligned}f(x,y) = 2x + i4y &= 2 \frac{(z+z^*)}{2} + i4 \frac{-i(z-z^*)}{2} \\ &= z+z^* + (2z-2z^*) \\ &= 3z-z^*\end{aligned}$$

A: ***NO!***    *It's a function of  $z$  and  $z^*$  so not analytic for either.*

## What's analytic? (...and what's not?)

Example: Q: Is  $f(x,y) = 2x + i4y$  an analytic function of  $z=x+iy$ ?

Well, test it using definitions:  $z = x + iy$  and:  $z^* = x - iy$   
or:  $x = (z+z^*)/2$  and:  $y = -i(z-z^*)/2$

$$\begin{aligned} f(x,y) = 2x + i4y &= 2 \frac{(z+z^*)}{2} + i4 \frac{-i(z-z^*)}{2} \\ &= z+z^* + (2z-2z^*) \\ &= 3z-z^* \end{aligned}$$

A: **NO!** It's a function of  $z$  and  $z^*$  so not analytic for either.

Example 2: Q: Is  $r(x,y) = x^2 + y^2$  an analytic function of  $z=x+iy$ ?

A: **NO!**  $r(x,y)=z^*z$  is a function of  $z$  and  $z^*$  so not analytic for either.

## What's analytic? (...and what's not?)

Example: Q: Is  $f(x,y) = 2x + i4y$  an analytic function of  $z=x+iy$ ?

Well, test it using definitions:  $z = x + iy$                       and:  $z^* = x - iy$   
or:  $x = (z+z^*)/2$                       and:  $y = -i(z-z^*)/2$

$$\begin{aligned} f(x,y) = 2x + i4y &= 2 \frac{(z+z^*)}{2} + i4 \frac{-i(z-z^*)}{2} \\ &= z+z^* + (2z-2z^*) \\ &= 3z-z^* \end{aligned}$$

A: **NO!** It's a function of  $z$  and  $z^*$  so not analytic for either.

Example 2: Q: Is  $r(x,y) = x^2 + y^2$  an analytic function of  $z=x+iy$ ?

A: **NO!**  $r(x,y)=z^*z$  is a function of  $z$  and  $z^*$  so not analytic for either.

Example 3: Q: Is  $s(x,y) = x^2-y^2 + 2ixy$  an analytic function of  $z=x+iy$ ?

A: **YES!**  $s(x,y)=(x+iy)^2 = z^2$  is analytic function of  $z$ . (Yay!)

## *4. Riemann-Cauchy conditions What's analytic? (...and what's not?)*

 *Easy 2D circulation and flux integrals*

*Easy 2D curvilinear coordinate discovery*

*Easy 2D monopole, dipole, and  $2^n$ -pole analysis*

*Easy  $2^n$ -multipole field and potential expansion*

*Easy stereo-projection visualization*

## What Good Are Complex Exponentials? (contd.)

9. Complex integrals  $\int f(z)dz$  count 2D “circulation” ( $\int \mathbf{F} \cdot d\mathbf{r}$ ) and “flux” ( $\int \mathbf{F} \times d\mathbf{r}$ )

Integral of  $f(z)$  between point  $z_1$  and point  $z_2$  is potential difference  $\Delta\phi = \phi(z_2) - \phi(z_1)$

$$\Delta\phi = \phi(z_2) - \phi(z_1) = \int_{z_1}^{z_2} f(z)dz = \underbrace{\Phi(x_2, y_2) - \Phi(x_1, y_1)}_{\Delta\Phi} + i \underbrace{[\mathbf{A}(x_2, y_2) - \mathbf{A}(x_1, y_1)]}_{\Delta\mathbf{A}}$$

$\Delta\phi = \qquad \qquad \Delta\Phi \qquad \qquad + i \qquad \qquad \Delta\mathbf{A}$

In *DFL*-field  $\mathbf{F}$ ,  $\Delta\phi$  is independent of the integration path  $z(t)$  connecting  $z_1$  and  $z_2$ .



## What Good Are Complex Exponentials? (contd.)

9. Complex integrals  $\int f(z)dz$  count 2D “circulation” ( $\int \mathbf{F} \cdot d\mathbf{r}$ ) and “flux” ( $\int \mathbf{F} \times d\mathbf{r}$ )

Integral of  $f(z)$  between point  $z_1$  and point  $z_2$  is potential difference  $\Delta\phi = \phi(z_2) - \phi(z_1)$

$$\Delta\phi = \phi(z_2) - \phi(z_1) = \int_{z_1}^{z_2} f(z)dz = \underbrace{\Phi(x_2, y_2) - \Phi(x_1, y_1)}_{\Delta\Phi} + i \underbrace{[\mathbf{A}(x_2, y_2) - \mathbf{A}(x_1, y_1)]}_{\Delta\mathbf{A}}$$

$$\Delta\phi = \Delta\Phi + i \Delta\mathbf{A}$$

In *DFL*-field  $\mathbf{F}$ ,  $\Delta\phi$  is independent of the integration path  $z(t)$  connecting  $z_1$  and  $z_2$ .

$$\begin{aligned} \int f(z) dz &= \int \left( f^*(z^*) \right)^* dz = \int \left( f^*(z^*) \right)^* (dx + i dy) = \int \left( f_x^* + i f_y^* \right)^* (dx + i dy) = \int \left( f_x^* - i f_y^* \right) (dx + i dy) \\ &= \int (f_x^* dx + f_y^* dy) + i \int (f_x^* dy - f_y^* dx) \\ &= \int \mathbf{F} \cdot d\mathbf{r} + i \int \mathbf{F} \times d\mathbf{r} \cdot \hat{\mathbf{e}}_z \\ &= \int \mathbf{F} \cdot d\mathbf{r} + i \int \mathbf{F} \cdot d\mathbf{r} \times \hat{\mathbf{e}}_z \\ &= \int \mathbf{F} \cdot d\mathbf{r} + i \int \mathbf{F} \cdot d\mathbf{S} \quad \text{where: } d\mathbf{S} = d\mathbf{r} \times \hat{\mathbf{e}}_z \end{aligned}$$

## What Good Are Complex Exponentials? (contd.)

### 9. Complex integrals $\int f(z)dz$ count 2D “circulation” ( $\int \mathbf{F} \cdot d\mathbf{r}$ ) and “flux” ( $\int \mathbf{F} \times d\mathbf{r}$ )

Integral of  $f(z)$  between point  $z_1$  and point  $z_2$  is potential difference  $\Delta\phi = \phi(z_2) - \phi(z_1)$



$$\Delta\phi = \phi(z_2) - \phi(z_1) = \int_{z_1}^{z_2} f(z)dz = \underbrace{\Phi(x_2, y_2) - \Phi(x_1, y_1)}_{\Delta\Phi} + i \underbrace{[\mathbf{A}(x_2, y_2) - \mathbf{A}(x_1, y_1)]}_{\Delta\mathbf{A}}$$

$$\Delta\phi = \Delta\Phi + i \Delta\mathbf{A}$$

In *DFL*-field  $\mathbf{F}$ ,  $\Delta\phi$  is independent of the integration path  $z(t)$  connecting  $z_1$  and  $z_2$ .

$$\begin{aligned} \int f(z)dz &= \int \left( f^*(z^*) \right)^* dz = \int \left( f^*(z^*) \right)^* (dx + i dy) = \int \left( f_x^* + i f_y^* \right)^* (dx + i dy) = \int \left( f_x^* - i f_y^* \right) (dx + i dy) \\ &= \int (f_x^* dx + f_y^* dy) + i \int (f_x^* dy - f_y^* dx) \\ &= \int \mathbf{F} \cdot d\mathbf{r} + i \int \mathbf{F} \times d\mathbf{r} \cdot \hat{\mathbf{e}}_z \\ &= \int \mathbf{F} \cdot d\mathbf{r} + i \int \mathbf{F} \cdot d\mathbf{r} \times \hat{\mathbf{e}}_z \\ &= \boxed{\int \mathbf{F} \cdot d\mathbf{r}} + i \boxed{\int \mathbf{F} \cdot d\mathbf{S}} \quad \text{where: } d\mathbf{S} = d\mathbf{r} \times \hat{\mathbf{e}}_z \end{aligned}$$

**Real part**  $\int_1^2 \mathbf{F} \cdot d\mathbf{r} = \Delta\Phi$   
 sums  $\mathbf{F}$  projections *along* path  $d\mathbf{r}$  that is, *circulation* on path to get  $\Delta\Phi$ .




**Imaginary part**  $\int_1^2 \mathbf{F} \cdot d\mathbf{S} = \Delta\mathbf{A}$   
 sums  $\mathbf{F}$  projection *across* path  $d\mathbf{r}$  that is, *flux* thru surface elements  $d\mathbf{S} = d\mathbf{r} \times \mathbf{e}_z$  normal to  $d\mathbf{r}$  to get  $\Delta\mathbf{A}$ .

Here the scalar potential  $\Phi=(x^2-y^2)/2$  is stereo-plotted vs.  $(x,y)$

The  $\Phi=(x^2-y^2)/2=const.$  curves are topography lines

The  $A=(xy)=const.$  curves are streamlines normal to topography lines



## *4. Riemann-Cauchy conditions What's analytic? (...and what's not?)*

*Easy 2D circulation and flux integrals*

 *Easy 2D curvilinear coordinate discovery*

*Easy 2D monopole, dipole, and  $2^n$ -pole analysis*

*Easy  $2^n$ -multipole field and potential expansion*

*Easy stereo-projection visualization*

## What Good Are Complex Exponentials? (contd.)


### 10. Complex potentials define 2D Orthogonal Curvilinear Coordinates (OCC) of field

The  $(\Phi, A)$  grid is a GCC coordinate system\*:

$$q^1 = \Phi = (x^2 - y^2)/2 = \text{const.}$$

$$q^2 = A = (xy) = \text{const.}$$

\*Actually it's OCC.



$$\begin{aligned}
 \text{Kajobian} &= \begin{pmatrix} \frac{\partial q^1}{\partial x} & \frac{\partial q^1}{\partial y} \\ \frac{\partial q^2}{\partial x} & \frac{\partial q^2}{\partial y} \end{pmatrix} = \begin{pmatrix} \frac{\partial \Phi}{\partial x} & \frac{\partial \Phi}{\partial y} \\ \frac{\partial A}{\partial x} & \frac{\partial A}{\partial y} \end{pmatrix} = \begin{pmatrix} x & -y \\ y & x \end{pmatrix} \begin{matrix} \leftarrow \mathbf{E}^\Phi \\ \leftarrow \mathbf{E}^A \end{matrix} & \quad \text{Jacobian} = \begin{pmatrix} \frac{\partial x}{\partial q^1} & \frac{\partial x}{\partial q^2} \\ \frac{\partial y}{\partial q^1} & \frac{\partial y}{\partial q^2} \end{pmatrix} = \begin{pmatrix} \frac{\partial x}{\partial \Phi} & \frac{\partial x}{\partial A} \\ \frac{\partial y}{\partial \Phi} & \frac{\partial y}{\partial A} \end{pmatrix} = \frac{1}{r^2} \begin{pmatrix} x & y \\ -y & x \end{pmatrix} \\
 & & & & & & & \begin{matrix} \uparrow & \uparrow \\ \mathbf{E}_\Phi & \mathbf{E}_A \end{matrix} & & \begin{matrix} \uparrow & \uparrow \\ \mathbf{E}_\Phi & \mathbf{E}_A \end{matrix}
 \end{aligned}$$

$$\text{Metric tensor} = \begin{pmatrix} g_{\Phi\Phi} & g_{\Phi A} \\ g_{A\Phi} & g_{AA} \end{pmatrix} = \begin{pmatrix} \mathbf{E}_\Phi \cdot \mathbf{E}_\Phi & \mathbf{E}_\Phi \cdot \mathbf{E}_A \\ \mathbf{E}_A \cdot \mathbf{E}_\Phi & \mathbf{E}_A \cdot \mathbf{E}_A \end{pmatrix} = \begin{pmatrix} r^2 & 0 \\ 0 & r^2 \end{pmatrix} \text{ where: } r^2 = x^2 + y^2$$

# What Good Are Complex Exponentials? (contd.)


## 10. Complex potentials define 2D Orthogonal Curvilinear Coordinates (OCC) of field

The  $(\Phi, \mathbf{A})$  grid is a GCC coordinate system\*:

$$q^1 = \Phi = (x^2 - y^2)/2 = \text{const.}$$

$$q^2 = \mathbf{A} = (xy) = \text{const.}$$

\*Actually it's OCC.



$$\text{Kajobian} = \begin{pmatrix} \frac{\partial q^1}{\partial x} & \frac{\partial q^1}{\partial y} \\ \frac{\partial q^2}{\partial x} & \frac{\partial q^2}{\partial y} \end{pmatrix} = \begin{pmatrix} \frac{\partial \Phi}{\partial x} & \frac{\partial \Phi}{\partial y} \\ \frac{\partial \mathbf{A}}{\partial x} & \frac{\partial \mathbf{A}}{\partial y} \end{pmatrix} = \begin{pmatrix} x & -y \\ y & x \end{pmatrix} \leftarrow \begin{matrix} \mathbf{E}^\Phi \\ \mathbf{E}^{\mathbf{A}} \end{matrix}$$

$$\text{Jacobian} = \begin{pmatrix} \frac{\partial x}{\partial q^1} & \frac{\partial x}{\partial q^2} \\ \frac{\partial y}{\partial q^1} & \frac{\partial y}{\partial q^2} \end{pmatrix} = \begin{pmatrix} \frac{\partial \Phi}{\partial \Phi} & \frac{\partial \Phi}{\partial \mathbf{A}} \\ \frac{\partial \mathbf{A}}{\partial \Phi} & \frac{\partial \mathbf{A}}{\partial \mathbf{A}} \end{pmatrix} = \frac{1}{r^2} \begin{pmatrix} x & y \\ -y & x \end{pmatrix}$$

$$\text{Metric tensor} = \begin{pmatrix} g_{\Phi\Phi} & g_{\Phi\mathbf{A}} \\ g_{\mathbf{A}\Phi} & g_{\mathbf{A}\mathbf{A}} \end{pmatrix} = \begin{pmatrix} \mathbf{E}_\Phi \cdot \mathbf{E}_\Phi & \mathbf{E}_\Phi \cdot \mathbf{E}_{\mathbf{A}} \\ \mathbf{E}_{\mathbf{A}} \cdot \mathbf{E}_\Phi & \mathbf{E}_{\mathbf{A}} \cdot \mathbf{E}_{\mathbf{A}} \end{pmatrix} = \begin{pmatrix} r^2 & 0 \\ 0 & r^2 \end{pmatrix} \text{ where: } r^2 = x^2 + y^2$$

Riemann-Cauchy Derivative Relations make coordinates orthogonal

$$\nabla \Phi = \begin{pmatrix} \frac{\partial \Phi}{\partial x} \\ \frac{\partial \Phi}{\partial y} \end{pmatrix} = \begin{pmatrix} \frac{\partial}{\partial x} \frac{a}{2} (x^2 - y^2) \\ \frac{\partial}{\partial y} \frac{a}{2} (x^2 - y^2) \end{pmatrix} = \begin{pmatrix} ax \\ -ay \end{pmatrix} = \mathbf{F}$$

The half-n'-half results assure

$$\mathbf{E}_\Phi \cdot \mathbf{E}_{\mathbf{A}} = \frac{\partial \Phi}{\partial x} \frac{\partial \mathbf{A}}{\partial x} + \frac{\partial \Phi}{\partial y} \frac{\partial \mathbf{A}}{\partial y}$$

$$= -\frac{\partial \Phi}{\partial x} \frac{\partial \Phi}{\partial y} + \frac{\partial \Phi}{\partial y} \frac{\partial \Phi}{\partial x} = 0$$

$$\nabla \times \mathbf{A} = \begin{pmatrix} \frac{\partial \mathbf{A}}{\partial y} \\ -\frac{\partial \mathbf{A}}{\partial x} \end{pmatrix} = \begin{pmatrix} \frac{\partial}{\partial y} axy \\ -\frac{\partial}{\partial x} axy \end{pmatrix} = \begin{pmatrix} ax \\ -ay \end{pmatrix} = \mathbf{F}$$



# What Good Are Complex Exponentials? (contd.)


## 10. Complex potentials define 2D Orthogonal Curvilinear Coordinates (OCC) of field

The  $(\Phi, A)$  grid is a GCC coordinate system\*:

$$q^1 = \Phi = (x^2 - y^2)/2 = \text{const.}$$

$$q^2 = A = (xy) = \text{const.}$$

\*Actually it's OCC.



$$\text{Kajobian} = \begin{pmatrix} \frac{\partial q^1}{\partial x} & \frac{\partial q^1}{\partial y} \\ \frac{\partial q^2}{\partial x} & \frac{\partial q^2}{\partial y} \end{pmatrix} = \begin{pmatrix} \frac{\partial \Phi}{\partial x} & \frac{\partial \Phi}{\partial y} \\ \frac{\partial A}{\partial x} & \frac{\partial A}{\partial y} \end{pmatrix} = \begin{pmatrix} x & -y \\ y & x \end{pmatrix} \leftarrow \begin{matrix} \mathbf{E}^\Phi \\ \mathbf{E}^A \end{matrix}$$

$$\text{Jacobian} = \begin{pmatrix} \frac{\partial x}{\partial q^1} & \frac{\partial x}{\partial q^2} \\ \frac{\partial y}{\partial q^1} & \frac{\partial y}{\partial q^2} \end{pmatrix} = \begin{pmatrix} \frac{\partial \Phi}{\partial \Phi} & \frac{\partial \Phi}{\partial A} \\ \frac{\partial A}{\partial \Phi} & \frac{\partial A}{\partial A} \end{pmatrix} = \frac{1}{r^2} \begin{pmatrix} x & y \\ -y & x \end{pmatrix}$$

$$\text{Metric tensor} = \begin{pmatrix} g_{\Phi\Phi} & g_{\Phi A} \\ g_{A\Phi} & g_{AA} \end{pmatrix} = \begin{pmatrix} \mathbf{E}_\Phi \cdot \mathbf{E}_\Phi & \mathbf{E}_\Phi \cdot \mathbf{E}_A \\ \mathbf{E}_A \cdot \mathbf{E}_\Phi & \mathbf{E}_A \cdot \mathbf{E}_A \end{pmatrix} = \begin{pmatrix} r^2 & 0 \\ 0 & r^2 \end{pmatrix} \text{ where: } r^2 = x^2 + y^2$$

Riemann-Cauchy Derivative Relations make coordinates orthogonal

$$\nabla \Phi = \begin{pmatrix} \frac{\partial \Phi}{\partial x} \\ \frac{\partial \Phi}{\partial y} \end{pmatrix} = \begin{pmatrix} \frac{\partial}{\partial x} \frac{a}{2} (x^2 - y^2) \\ \frac{\partial}{\partial y} \frac{a}{2} (x^2 - y^2) \end{pmatrix} = \begin{pmatrix} ax \\ -ay \end{pmatrix} = \mathbf{F}$$

The half-n'-half results assure

$$\mathbf{E}_\Phi \cdot \mathbf{E}_A = \frac{\partial \Phi}{\partial x} \frac{\partial A}{\partial x} + \frac{\partial \Phi}{\partial y} \frac{\partial A}{\partial y}$$

$$= -\frac{\partial \Phi}{\partial x} \frac{\partial \Phi}{\partial y} + \frac{\partial \Phi}{\partial y} \frac{\partial \Phi}{\partial x} = 0$$

$$\nabla \times \mathbf{A} = \begin{pmatrix} \frac{\partial A}{\partial y} \\ -\frac{\partial A}{\partial x} \end{pmatrix} = \begin{pmatrix} \frac{\partial}{\partial y} axy \\ -\frac{\partial}{\partial x} axy \end{pmatrix} = \begin{pmatrix} ax \\ -ay \end{pmatrix} = \mathbf{F}$$

Zero divergence requirement:  $0 = \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} = \frac{\partial}{\partial x} \frac{\partial \Phi}{\partial x} + \frac{\partial}{\partial y} \frac{\partial \Phi}{\partial y} = \frac{\partial^2 \Phi}{\partial x^2} + \frac{\partial^2 \Phi}{\partial y^2} = 0$  potential  $\Phi$  obeys Laplace equation

# What Good Are Complex Exponentials? (contd.)


## 10. Complex potentials define 2D Orthogonal Curvilinear Coordinates (OCC) of field

The  $(\Phi, A)$  grid is a GCC coordinate system\*:

$$q^1 = \Phi = (x^2 - y^2)/2 = \text{const.}$$

$$q^2 = A = (xy) = \text{const.}$$

\*Actually it's OCC.



$$Kajobian = \begin{pmatrix} \frac{\partial q^1}{\partial x} & \frac{\partial q^1}{\partial y} \\ \frac{\partial q^2}{\partial x} & \frac{\partial q^2}{\partial y} \end{pmatrix} = \begin{pmatrix} \frac{\partial \Phi}{\partial x} & \frac{\partial \Phi}{\partial y} \\ \frac{\partial A}{\partial x} & \frac{\partial A}{\partial y} \end{pmatrix} = \begin{pmatrix} x & -y \\ y & x \end{pmatrix} \leftarrow \begin{matrix} \mathbf{E}^\Phi \\ \mathbf{E}^A \end{matrix}$$

$$Jacobian = \begin{pmatrix} \frac{\partial x}{\partial q^1} & \frac{\partial x}{\partial q^2} \\ \frac{\partial y}{\partial q^1} & \frac{\partial y}{\partial q^2} \end{pmatrix} = \begin{pmatrix} \frac{\partial x}{\partial \Phi} & \frac{\partial x}{\partial A} \\ \frac{\partial y}{\partial \Phi} & \frac{\partial y}{\partial A} \end{pmatrix} = \frac{1}{r^2} \begin{pmatrix} x & y \\ -y & x \end{pmatrix}$$

$$Metric\ tensor = \begin{pmatrix} g_{\Phi\Phi} & g_{\Phi A} \\ g_{A\Phi} & g_{AA} \end{pmatrix} = \begin{pmatrix} \mathbf{E}_\Phi \cdot \mathbf{E}_\Phi & \mathbf{E}_\Phi \cdot \mathbf{E}_A \\ \mathbf{E}_A \cdot \mathbf{E}_\Phi & \mathbf{E}_A \cdot \mathbf{E}_A \end{pmatrix} = \begin{pmatrix} r^2 & 0 \\ 0 & r^2 \end{pmatrix} \text{ where: } r^2 = x^2 + y^2$$

Riemann-Cauchy Derivative Relations make coordinates orthogonal

$$\nabla \Phi = \begin{pmatrix} \frac{\partial \Phi}{\partial x} \\ \frac{\partial \Phi}{\partial y} \end{pmatrix} = \begin{pmatrix} \frac{\partial}{\partial x} \frac{a}{2} (x^2 - y^2) \\ \frac{\partial}{\partial y} \frac{a}{2} (x^2 - y^2) \end{pmatrix} = \begin{pmatrix} ax \\ -ay \end{pmatrix} = \mathbf{F}$$

The half-n'-half results assure

$$\mathbf{E}_\Phi \cdot \mathbf{E}_A = \frac{\partial \Phi}{\partial x} \frac{\partial A}{\partial x} + \frac{\partial \Phi}{\partial y} \frac{\partial A}{\partial y}$$

$$= -\frac{\partial \Phi}{\partial x} \frac{\partial \Phi}{\partial y} + \frac{\partial \Phi}{\partial y} \frac{\partial \Phi}{\partial x} = 0$$

$$\nabla \times \mathbf{A} = \begin{pmatrix} \frac{\partial A}{\partial y} \\ -\frac{\partial A}{\partial x} \end{pmatrix} = \begin{pmatrix} \frac{\partial}{\partial y} axy \\ -\frac{\partial}{\partial x} axy \end{pmatrix} = \begin{pmatrix} ax \\ -ay \end{pmatrix} = \mathbf{F}$$

or Riemann-Cauchy

Zero divergence requirement:  $0 = \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} = \frac{\partial}{\partial x} \frac{\partial \Phi}{\partial x} + \frac{\partial}{\partial y} \frac{\partial \Phi}{\partial y} = \frac{\partial^2 \Phi}{\partial x^2} + \frac{\partial^2 \Phi}{\partial y^2} = 0$  and so does  $A$  potential  $\Phi$  obeys Laplace equation



## *4. Riemann-Cauchy conditions What's analytic? (...and what's not?)*

*Easy 2D circulation and flux integrals*

*Easy 2D curvilinear coordinate discovery*

 *Easy 2D monopole, dipole, and  $2^n$ -pole analysis*

*Easy  $2^n$ -multipole field and potential expansion*

*Easy stereo-projection visualization*

## What Good Are Complex Exponentials? (contd.)

### 11. Complex integrals define 2D *monopole* fields and potentials

Of all power-law fields  $f(z)=az^n$  one lacks a power-law potential  $\phi(z)=\frac{a}{n+1}z^{n+1}$ . It is the  $n = -1$  case.

Unit *monopole* field:  $f(z)=\frac{1}{z}=z^{-1}$

$f(z)=\frac{a}{z}=az^{-1}$  Source- $a$  *monopole*

It has a *logarithmic potential*  $\phi(z)=a\cdot\ln(z)=a\cdot\ln(x+iy)$ .

## What Good Are Complex Exponentials? (contd.)

### 11. Complex integrals define 2D *monopole* fields and potentials

Of all power-law fields  $f(z)=az^n$  one lacks a power-law potential  $\phi(z)=\frac{a}{n+1}z^{n+1}$ . It is the  $n = -1$  case.

Unit *monopole* field:  $f(z)=\frac{1}{z}=z^{-1}$        $f(z)=\frac{a}{z}=az^{-1}$  Source- $a$  *monopole*

It has a *logarithmic potential*  $\phi(z)=a\cdot\ln(z)=a\cdot\ln(x+iy)$ .

$$\phi(z) = \Phi + i\mathbf{A} = \int f(z)dz = \int \frac{a}{z} dz = a \ln(z)$$

## What Good Are Complex Exponentials? (contd.)

### 11. Complex integrals define 2D *monopole* fields and potentials

Of all power-law fields  $f(z)=az^n$  one lacks a power-law potential  $\phi(z)=\frac{a}{n+1}z^{n+1}$ . It is the  $n = -1$  case.

$$\text{Unit monopole field: } f(z)=\frac{1}{z}=z^{-1} \qquad f(z)=\frac{a}{z}=az^{-1} \text{ Source-}a \text{ monopole}$$

It has a *logarithmic potential*  $\phi(z)=a \cdot \ln(z)=a \cdot \ln(x+iy)$ . Note:  $\ln(a \cdot b)=\ln(a)+\ln(b)$ ,  $\ln(e^{i\theta})=i\theta$ , and  $z=re^{i\theta}$ .

$$\begin{aligned} \phi(z) &= \Phi + i\mathbf{A} = \int f(z)dz = \int \frac{a}{z} dz = a \ln(z) = a \ln(re^{i\theta}) \\ &= \underbrace{a \ln(r)} + i \underbrace{a\theta} \end{aligned}$$

## What Good Are Complex Exponentials? (contd.)

### 11. Complex integrals define 2D *monopole* fields and potentials


Of all power-law fields  $f(z)=az^n$  one lacks a power-law potential  $\phi(z)=\frac{a}{n+1}z^{n+1}$ . It is the  $n = -1$  case.

Unit *monopole* field:  $f(z)=\frac{1}{z}=z^{-1}$        $f(z)=\frac{a}{z}=az^{-1}$  Source- $a$  *monopole*

It has a *logarithmic potential*  $\phi(z)=a\cdot\ln(z)=a\cdot\ln(x+iy)$ . Note:  $\ln(a\cdot b)=\ln(a)+\ln(b)$ ,  $\ln(e^{i\theta})=i\theta$ , and  $z=re^{i\theta}$ .

$$\begin{aligned}\phi(z) &= \Phi + i\mathbf{A} = \int f(z)dz = \int \frac{a}{z} dz = a \ln(z) = a \ln(re^{i\theta}) \\ &= \underbrace{a \ln(r)} + i \underbrace{a\theta}\end{aligned}$$

(a) Unit Z-line-flux field  $f(z)=1/z$



Lecture 12 Tue. 10.03  
 May end here

## What Good Are Complex Exponentials? (contd.)

### 11. Complex integrals define 2D *monopole* fields and potentials

Of all power-law fields  $f(z)=az^n$  one lacks a power-law potential  $\phi(z)=\frac{a}{n+1}z^{n+1}$ . It is the  $n = -1$  case.


Unit *monopole* field:  $f(z)=\frac{1}{z}=z^{-1}$        $f(z)=\frac{a}{z}=az^{-1}$  Source- $a$  *monopole*

It has a *logarithmic potential*  $\phi(z)=a \cdot \ln(z)=a \cdot \ln(x+iy)$ . Note:  $\ln(a \cdot b)=\ln(a)+\ln(b)$ ,  $\ln(e^{i\theta})=i\theta$ , and  $z=re^{i\theta}$ .


$$\begin{aligned} \phi(z) &= \Phi + i\mathbf{A} = \int f(z)dz = \int \frac{a}{z} dz = a \ln(z) = a \ln(re^{i\theta}) \\ &= \underbrace{a \ln(r)} + i \underbrace{a\theta} \end{aligned}$$

(a) Unit Z-line-flux field  $f(z)=1/z$

(b) Unit Z-line-vortex field  $f(z)=i/z$



Field:  
 $f^*(z^*)=1/z^*=e^{i\theta}/r$   
 $\mathbf{F}_{(x,y)}=(x,y)/r^2$   
Potential:  
 $\phi(z)=\ln z$   
 $=\ln r+i\theta$   
 $=\Phi+i\mathbf{A}$



Field:  
 $f^*(z^*)=-i/z^*=-ie^{i\theta}/r$   
 $\mathbf{F}_{(x,y)}=(y,-x)/r^2$   
Potential:  
 $\phi(z)=i \ln z$   
 $=\theta+i \ln r$   
 $=\Phi+i\mathbf{A}$

## What Good Are Complex Exponentials? (contd.)

### 11. Complex integrals define 2D *monopole* fields and potentials



Of all power-law fields  $f(z)=az^n$  one lacks a power-law potential  $\phi(z)=\frac{a}{n+1}z^{n+1}$ . It is the  $n = -1$  case.

Unit *monopole* field:  $f(z)=\frac{1}{z}=z^{-1}$        $f(z)=\frac{a}{z}=az^{-1}$  Source- $a$  *monopole*

It has a *logarithmic potential*  $\phi(z)=a\cdot\ln(z)=a\cdot\ln(x+iy)$ . Note:  $\ln(a\cdot b)=\ln(a)+\ln(b)$ ,  $\ln(e^{i\theta})=i\theta$ , and  $z=re^{i\theta}$ .

$$\begin{aligned} \phi(z) &= \underbrace{\Phi}_{= a \ln(r)} + \underbrace{i\mathbf{A}}_{+ i a \theta} = \int f(z)dz = \int \frac{a}{z} dz = a \ln(z) = a \ln(re^{i\theta}) \\ &= a \ln(r) + i a \theta \end{aligned}$$

A *monopole* field is the only power-law field whose integral (potential) depends on *path of integration*.




*path that goes N times around origin ( $r=0$ ) at constant  $r = R$ .*

$$\Delta\phi = \oint f(z)dz = a \oint \frac{dz}{z} = a \int_{\theta=0}^{\theta=2\pi N} \frac{d(Re^{i\theta})}{Re^{i\theta}} = a \int_{\theta=0}^{\theta=2\pi N} id\theta = ai \theta \Big|_0^{2\pi N} = 2a\pi iN$$




(a) Unit Z-line-flux field  $f(z)=1/z$




$$\phi(z) = \underbrace{\Phi}_{\ln(r)} + \underbrace{iA}_{i\theta} = \int f(z)dz = \int \frac{a}{z} dz = a \ln(re^{i\theta})$$

*(For a=1)*






*Each turn around origin adds  $2\pi i$  to vector potential  $iA$*






*(For a=1)*



(a) Unit Z-line-flux field  $f(z)=1/z$



(b) Unit Z-line-vortex field  $f(z)=i/z$




# What Good Are Complex Exponentials? (contd.)

$$f(z) = (0.5 + i0.5)/z = e^{i\pi/4}/z\sqrt{2}$$

“Vortex”

x= -0.82 y= -4.8 Fx= -0.12 Fy= -0.084 Phi = 1.7 A = -0.078




$\phi(z) = A \ln z$   $f(z) = A/z$   $A = 0.5 + i0.5$


$$f(z) = (0.75 + i0.25)/z = e^{i18^\circ}/z\sqrt{n}$$

“Hurricane”

x= -3.6 y= 3.2 Fx= -0.083 Fy= 0.14 Phi = 0.57 A = 2.2



$\phi(z) = A \ln z$   $f(z) = A/z$   $A = 0.75 + i0.25$



## *4. Riemann-Cauchy conditions What's analytic? (...and what's not?)*

*Easy 2D circulation and flux integrals*

*Easy 2D curvilinear coordinate discovery*

 *Easy 2D monopole, dipole, and  $2^n$ -pole analysis*

*Easy  $2^n$ -multipole field and potential expansion*

*Easy stereo-projection visualization*



# What Good Are Complex Exponentials? (2D monopole, dipole, and $2^n$ -pole analysis)

## 12. Complex derivatives give 2D dipole fields

Start with  $f(z)=az^{-1}$ : 2D line *monopole field* and is its *monopole potential*  $\phi(z)=a \ln z$  of source strength  $a$ .

$$f^{1-pole}(z) = \frac{a}{z} = \frac{d\phi^{1-pole}}{dz} \quad \phi^{1-pole}(z) = a \ln z$$

Now let these two line-sources of equal but opposite source constants  $+a$  and  $-a$  be located at  $z=\pm\Delta/2$  separated by a small interval  $\Delta$ . This sum (actually difference) of  $f^{1-pole}$ -fields is called a *dipole field*.

$$f^{dipole}(z) = \frac{a}{z + \frac{\Delta}{2}} - \frac{a}{z - \frac{\Delta}{2}} = \frac{-a \cdot \Delta}{z^2 - \frac{\Delta^2}{4}} \quad \phi^{dipole}(z) = a \ln\left(z - \frac{\Delta}{2}\right) - a \ln\left(z + \frac{\Delta}{2}\right) = a \ln \frac{z - \frac{\Delta}{2}}{z + \frac{\Delta}{2}}$$

*This is like the derivative definition:*


$$\frac{df}{dz} = \frac{f(z + \Delta) - f(z)}{\Delta}$$

*or:*

$$\frac{df}{dz} = \frac{f\left(z + \frac{\Delta}{2}\right) - f\left(z - \frac{\Delta}{2}\right)}{\Delta}$$

*if  $\Delta$  is infinitesimal*

$$(\Delta \rightarrow 0)$$



*So-called “physical dipole” has finite  $\Delta$  (+)(-) separation*

## What Good Are Complex Exponentials? (2D monopole, dipole, and 2<sup>n</sup>-pole analysis)

### 12. Complex derivatives give 2D dipole fields

Start with  $f(z)=az^{-1}$ : 2D line *monopole field* and is its *monopole potential*  $\phi(z)=a \ln z$  of source strength  $a$ .

$$f^{1-pole}(z) = \frac{a}{z} = \frac{d\phi^{1-pole}}{dz} \quad \phi^{1-pole}(z) = a \ln z$$

Now let these two line-sources of equal but opposite source constants  $+a$  and  $-a$  be located at  $z=\pm\Delta/2$  separated by a small interval  $\Delta$ . This sum (actually difference) of  $f^{1-pole}$ -fields is called a *dipole field*.

$$f^{dipole}(z) = \frac{a}{z + \frac{\Delta}{2}} - \frac{a}{z - \frac{\Delta}{2}} = \frac{-a \cdot \Delta}{z^2 - \frac{\Delta^2}{4}} \quad \phi^{dipole}(z) = a \ln\left(z - \frac{\Delta}{2}\right) - a \ln\left(z + \frac{\Delta}{2}\right) = a \ln \frac{z - \frac{\Delta}{2}}{z + \frac{\Delta}{2}}$$

If interval  $\Delta$  is *tiny* and is divided out we get a *point-dipole field*  $f^{2-pole}$  that is the  $z$ -derivative of  $f^{1-pole}$ .

$$f^{2-pole} = \frac{-a}{z^2} = \frac{df^{1-pole}}{dz} = \frac{d\phi^{2-pole}}{dz} \quad \phi^{2-pole} = \frac{a}{z} = \frac{d\phi^{1-pole}}{dz}$$

## What Good Are Complex Exponentials? (2D monopole, dipole, and $2^n$ -pole analysis)

### 12. Complex derivatives give 2D dipole fields

Start with  $f(z)=az^{-1}$ : 2D line *monopole field* and is its *monopole potential*  $\phi(z)=a \ln z$  of source strength  $a$ .


$$f^{1-pole}(z) = \frac{a}{z} = \frac{d\phi^{1-pole}}{dz} \quad \phi^{1-pole}(z) = a \ln z$$

Now let these two line-sources of equal but opposite source constants  $+a$  and  $-a$  be located at  $z=\pm\Delta/2$  separated by a small interval  $\Delta$ . This sum (actually difference) of  $f^{1-pole}$ -fields is called a *dipole field*.

$$f^{dipole}(z) = \frac{a}{z + \frac{\Delta}{2}} - \frac{a}{z - \frac{\Delta}{2}} = \frac{-a \cdot \Delta}{z^2 - \frac{\Delta^2}{4}} \quad \phi^{dipole}(z) = a \ln\left(z - \frac{\Delta}{2}\right) - a \ln\left(z + \frac{\Delta}{2}\right) = a \ln \frac{z - \frac{\Delta}{2}}{z + \frac{\Delta}{2}}$$

If interval  $\Delta$  is *tiny* and is divided out we get a *point-dipole field*  $f^{2-pole}$  that is the  $z$ -derivative of  $f^{1-pole}$ .

$$f^{2-pole} = \frac{-a}{z^2} = \frac{df^{1-pole}}{dz} = \frac{d\phi^{2-pole}}{dz} \quad \phi^{2-pole} = \frac{a}{z} = \frac{d\phi^{1-pole}}{dz}$$



## What Good Are Complex Exponentials? (2D monopole, dipole, and 2<sup>n</sup>-pole analysis)

### 12. Complex derivatives give 2D dipole fields

Start with  $f(z)=az^{-1}$ : 2D line *monopole field* and is its *monopole potential*  $\phi(z)=a \ln z$  of source strength  $a$ .

$$f^{1-pole}(z) = \frac{a}{z} = \frac{d\phi^{1-pole}}{dz} \quad \phi^{1-pole}(z) = a \ln z$$

Now let these two line-sources of equal but opposite source constants  $+a$  and  $-a$  be located at  $z=\pm\Delta/2$  separated by a small interval  $\Delta$ . This sum (actually difference) of  $f^{1-pole}$ -fields is called a *dipole field*.

$$f^{dipole}(z) = \frac{a}{z+\frac{\Delta}{2}} - \frac{a}{z-\frac{\Delta}{2}} = \frac{-a \cdot \Delta}{z^2 - \frac{\Delta^2}{4}} \quad \phi^{dipole}(z) = a \ln\left(z - \frac{\Delta}{2}\right) - a \ln\left(z + \frac{\Delta}{2}\right) = a \ln \frac{z - \frac{\Delta}{2}}{z + \frac{\Delta}{2}}$$

If interval  $\Delta$  is *tiny* and is divided out we get a *point-dipole field*  $f^{2-pole}$  that is the  $z$ -derivative of  $f^{1-pole}$ .


$$f^{2-pole} = \frac{-a}{z^2} = \frac{df^{1-pole}}{dz} = \frac{d\phi^{2-pole}}{dz} \quad \phi^{2-pole} = \frac{a}{z} = \frac{d\phi^{1-pole}}{dz}$$

A *point-dipole potential*  $\phi^{2-pole}$  (whose  $z$ -derivative is  $f^{2-pole}$ ) is a  $z$ -derivative of  $\phi^{1-pole}$ .

$$\begin{aligned} \phi^{2-pole} &= \frac{a}{z} = \frac{a}{x+iy} = \frac{a}{x+iy} \frac{x-iy}{x-iy} = \frac{ax}{x^2+y^2} + i \frac{-ay}{x^2+y^2} = \frac{a}{r} \cos \theta - i \frac{a}{r} \sin \theta \\ &= \Phi^{2-pole} + i \mathbf{A}^{2-pole} \end{aligned}$$


A *point-dipole potential*  $\phi^{2-pole}$  (whose  $z$ -derivative is  $f^{2-pole}$ ) is a  $z$ -derivative of  $\phi^{1-pole}$ .

$$\begin{aligned} \phi^{2-pole} &= \frac{a}{z} = \frac{a}{x+iy} = \frac{a}{x+iy} \frac{x-iy}{x-iy} = \frac{ax}{x^2+y^2} + i \frac{-ay}{x^2+y^2} = \frac{a}{r} \cos \theta - i \frac{a}{r} \sin \theta \\ &= \Phi^{2-pole} + i A^{2-pole} \end{aligned}$$



Scalar potentials

$\Phi = (a/r) \cos \theta = const.$



Vector potentials

$A = (a/r) \sin \theta = const.$



## $2^n$ -pole analysis (quadrupole: $2^2=4$ -pole, octapole: $2^3=8$ -pole, ..., pole dancer,

What if we put a (-)copy of a 2-pole near its original?

Well, the result is 4-pole or *quadrupole* field  $f^{4-pole}$  and potential  $\phi^{4-pole}$ .

Each a z-derivative of  $f^{2-pole}$  and  $\phi^{2-pole}$ .

$$f^{4-pole} = \frac{a}{z^3} = \frac{1}{2} \frac{df^{2-pole}}{dz} = \frac{d\phi^{4-pole}}{dz}$$

$$\phi^{4-pole} = -\frac{a}{2z^2} = \frac{1}{2} \frac{d\phi^{2-pole}}{dz}$$

# $2^n$ -pole analysis (quadrupole: $2^2=4$ -pole, octapole: $2^3=8$ -pole, ..., pole dancer,



What if we put a (-)copy of a 2-pole near its original?

Well, the result is 4-pole or *quadrupole* field  $f^{4-pole}$  and potential  $\phi^{4-pole}$ .

Each a z-derivative of  $f^{2-pole}$  and  $\phi^{2-pole}$ .

$$f^{4-pole} = \frac{a}{z^3} = \frac{1}{2} \frac{df^{2-pole}}{dz} = \frac{d\phi^{4-pole}}{dz}$$

$$\phi^{4-pole} = -\frac{a}{2z^2} = \frac{1}{2} \frac{d\phi^{2-pole}}{dz}$$



Field:  
 $f^*(z^*) = 1/z^3 = e^{i3\theta}/r^3$   
 $\mathbf{F}(x,y) = (\cos 3\theta, \sin 3\theta)/r^3$   
Potential:  
 $-2\phi(z) = 1/z^2$   
 $= (\cos 2\theta)/r^2 + i(\sin 2\theta)/r^2$   
 $= \Phi + iA$

## *4. Riemann-Cauchy conditions What's analytic? (...and what's not?)*

*Easy 2D circulation and flux integrals*

*Easy 2D curvilinear coordinate discovery*

*Easy 2D monopole, dipole, and  $2^n$ -pole analysis*

*Easy  $2^n$ -multipole field and potential expansion*

*Easy stereo-projection visualization*



## $2^n$ -pole analysis: Laurent series (Generalization of Maclaurin-Taylor series)

*Laurent series* or *multipole expansion* of a given complex field function  $f(z)$  around  $z=0$ .

$$\begin{aligned} \frac{d\phi}{dz} = f(z) &= \dots a_{-3}z^{-3} + a_{-2}z^{-2} + \mathbf{a_{-1}z^{-1}} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots \\ &\quad \dots \begin{array}{l} 2^2\text{-pole} \\ \text{(quadrupole)} \\ \text{at } z=0 \end{array} \begin{array}{l} 2^1\text{-pole} \\ \text{(dipole)} \\ \text{at } z=0 \end{array} \begin{array}{l} \mathbf{2^0\text{-pole}} \\ \text{(monopole)} \\ \text{at } z=0 \end{array} \begin{array}{l} 2^1\text{-pole} \\ \text{(dipole)} \\ \text{at } z=\infty \end{array} \begin{array}{l} 2^2\text{-pole} \\ \text{(quadrupole)} \\ \text{at } z=\infty \end{array} \begin{array}{l} 2^3\text{-pole} \\ \text{(octapole)} \\ \text{at } z=\infty \end{array} \begin{array}{l} 2^4\text{-pole} \\ \text{(hexadecapole)} \\ \text{at } z=\infty \end{array} \begin{array}{l} 2^5\text{-pole} \\ \text{at } z=\infty \end{array} \begin{array}{l} 2^6\text{-pole} \\ \text{at } z=\infty \end{array} \dots \\ \int f dz = \\ \phi(z) &= \dots \frac{a_{-3}}{-2}z^{-2} + \frac{a_{-2}}{-1}z^{-1} + \mathbf{a_{-1} \ln z} + a_0z + \frac{a_1}{2}z^2 + \frac{a_2}{3}z^3 + \frac{a_3}{4}z^4 + \frac{a_4}{5}z^5 + \frac{a_5}{6}z^6 + \dots \end{aligned}$$

All field terms  $a_{m-1}z^{m-1}$  except  $\mathbf{1\text{-pole } \frac{a_{-1}}{z}}$  have potential term  $a_{m-1}z^m/m$  of a  $2^m$ -pole.

These are located at  $z=0$  for  $m < 0$  and at  $z=\infty$  for  $m > 0$ .

$$\phi(z) = \dots \begin{array}{l} \text{(octapole)}_0 \\ \frac{a_{-4}}{-3}z^{-3} \end{array} + \begin{array}{l} \text{(quadrupole)}_0 \\ \frac{a_{-3}}{-2}z^{-2} \end{array} + \begin{array}{l} \text{(dipole)}_0 \\ \frac{a_{-2}}{-1}z^{-1} \end{array} + \mathbf{a_{-1} \ln z} + \begin{array}{l} \text{(monopole)} \\ a_0z \end{array} + \begin{array}{l} \text{(dipole)}_\infty \\ \frac{a_1}{2}z^2 \end{array} + \begin{array}{l} \text{(quadrupole)}_\infty \\ \frac{a_2}{3}z^3 \end{array} + \dots$$

## $2^n$ -pole analysis: Laurent series (Generalization of Maclaurin-Taylor series)

*Laurent series* or *multipole expansion* of a given complex field function  $f(z)$  around  $z=0$ .

$$\begin{aligned} \frac{d\phi}{dz} = f(z) &= \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots \\ &\quad \dots \begin{array}{c} 2^2\text{-pole} \\ \text{(quadrupole)} \\ \text{at } z=0 \end{array} + \begin{array}{c} 2^1\text{-pole} \\ \text{(dipole)} \\ \text{at } z=0 \end{array} + \begin{array}{c} 2^0\text{-pole} \\ \text{(monopole)} \\ \text{at } z=0 \end{array} + \begin{array}{c} 2^1\text{-pole} \\ \text{(dipole)} \\ \text{at } z=\infty \end{array} + \begin{array}{c} 2^2\text{-pole} \\ \text{(quadrupole)} \\ \text{at } z=\infty \end{array} + \begin{array}{c} 2^3\text{-pole} \\ \text{(octapole)} \\ \text{at } z=\infty \end{array} + \begin{array}{c} 2^4\text{-pole} \\ \text{(hexadecapole)} \\ \text{at } z=\infty \end{array} + \begin{array}{c} 2^5\text{-pole} \\ \text{at } z=\infty \end{array} + \begin{array}{c} 2^6\text{-pole} \\ \text{at } z=\infty \end{array} \dots \\ \int f dz = \phi(z) &= \dots \frac{a_{-3}}{-2} z^{-2} + \frac{a_{-2}}{-1} z^{-1} + a_{-1} \ln z + a_0 z + \frac{a_1}{2} z^2 + \frac{a_2}{3} z^3 + \frac{a_3}{4} z^4 + \frac{a_4}{5} z^5 + \frac{a_5}{6} z^6 + \dots \end{aligned}$$

All field terms  $a_{m-1}z^{m-1}$  except  $1\text{-pole } \frac{a_{-1}}{z}$  have potential term  $a_{m-1}z^m/m$  of a  $2^m$ -pole.

These are located at  $z=0$  for  $m < 0$  and at  $z=\infty$  for  $m > 0$ .

$$\phi(z) = \dots \begin{array}{c} \text{(octapole)}_0 \\ \frac{a_{-3}}{-2} z^{-2} \end{array} + \begin{array}{c} \text{(quadrupole)}_0 \\ \frac{a_{-2}}{-2} z^{-2} \end{array} + \begin{array}{c} \text{(dipole)}_0 \\ \frac{a_{-1}}{-1} z^{-1} \end{array} + \begin{array}{c} \text{(monopole)} \\ a_{-1} \ln z \end{array} + \begin{array}{c} \text{(dipole)}_\infty \\ a_0 z \end{array} + \begin{array}{c} \text{(quadrupole)}_\infty \\ \frac{a_1}{2} z^2 \end{array} + \begin{array}{c} \text{(octapole)}_\infty \\ \frac{a_2}{3} z^3 \end{array} + \dots$$

$$\phi(w) = \dots \frac{a_{-4}}{-3} w^{-3} + \frac{a_{-3}}{-2} w^{-2} + \frac{a_{-2}}{-1} w^{-1} + a_{-1} \ln w + a_0 w + \frac{a_1}{2} w^2 + \frac{a_2}{3} w^3 + \dots$$

(with  $z=w^{-1}$ )

# $2^n$ -pole analysis: Laurent series (Generalization of Maclaurin-Taylor series)

*Laurent series* or *multipole expansion* of a given complex field function  $f(z)$  around  $z=0$ .

$$\frac{d\phi}{dz} = f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

	...	2 <sup>2</sup> -pole	2 <sup>1</sup> -pole	2 <sup>0</sup> -pole	2 <sup>1</sup> -pole	2 <sup>2</sup> -pole	2 <sup>3</sup> -pole	2 <sup>4</sup> -pole	2 <sup>5</sup> -pole	2 <sup>6</sup> -pole	...
		(quadrupole)	(dipole)	(monopole)	(dipole)	(quadrupole)	(octapole)	(hexadecapole)			
		at $z=0$	at $z=0$	at $z=0$	at $z=\infty$	at $z=\infty$	at $z=\infty$	at $z=\infty$	at $z=\infty$	at $z=\infty$	

$$\int f dz = \phi(z) = \dots \frac{a_{-3}}{-2} z^{-2} + \frac{a_{-2}}{-1} z^{-1} + a_{-1} \ln z + a_0 z + \frac{a_1}{2} z^2 + \frac{a_2}{3} z^3 + \frac{a_3}{4} z^4 + \frac{a_4}{5} z^5 + \frac{a_5}{6} z^6 + \dots$$

All field terms  $a_{m-1}z^{m-1}$  except *1-pole*  $\frac{a_{-1}}{z}$  have potential term  $a_{m-1}z^m/m$  of a  $2^m$ -pole.

These are located at  $z=0$  for  $m < 0$  and at  $z=\infty$  for  $m > 0$ .

$$\phi(z) = \dots \frac{a_{-4}}{-3} z^{-3} + \frac{a_{-3}}{-2} z^{-2} + \frac{a_{-2}}{-1} z^{-1} + a_{-1} \ln z + a_0 z + \frac{a_1}{2} z^2 + \frac{a_2}{3} z^3 + \dots$$

(octapole)<sub>0</sub>
(quadrupole)<sub>0</sub>
(dipole)<sub>0</sub>
(monopole)
(dipole)<sub>∞</sub>
(quadrupole)<sub>∞</sub>
(octapole)<sub>∞</sub>


$$\phi(w) = \dots \frac{a_{-4}}{-3} w^{-3} + \frac{a_{-3}}{-2} w^{-2} + \frac{a_{-2}}{-1} w^{-1} + a_{-1} \ln w + a_0 w + \frac{a_1}{2} w^2 + \frac{a_2}{3} w^3 + \dots$$

(with  $z \rightarrow w$ )



$$= \dots \frac{a_2}{3} z^{-3} + \frac{a_1}{2} z^{-2} + a_0 z^{-1} - a_{-1} \ln z + \frac{a_{-2}}{-1} z + \frac{a_{-3}}{-2} z^2 + \frac{a_{-4}}{-3} z^3 + \dots$$

(with  $w = z^{-1}$ )





$$\begin{aligned} \phi(z) &= \dots \frac{a_{-4}}{-3} z^{-3} + \frac{a_{-3}}{-2} z^{-2} + \frac{a_{-2}}{-1} z^{-1} + a_{-1} \ln z + a_0 z + \frac{a_1}{2} z^2 + \frac{a_2}{3} z^3 + \dots \\ &\quad \text{(octapole)}_0 \quad \text{(quadrupole)}_0 \quad \text{(dipole)}_0 \quad \text{(monopole)} \quad \text{(dipole)}_\infty \quad \text{(quadrupole)}_\infty \quad \text{(octapole)}_\infty \\ \phi(w) &= \dots \frac{a_{-4}}{-3} w^{-3} + \frac{a_{-3}}{-2} w^{-2} + \frac{a_{-2}}{-1} w^{-1} + a_{-1} \ln w + a_0 w + \frac{a_1}{2} w^2 + \frac{a_2}{3} w^3 + \dots \\ &\quad \text{(with } z \rightarrow w) \\ &= \dots \frac{a_2}{3} z^{-2} + \frac{a_1}{2} z^{-2} + a_0 z^{-1} - a_{-1} \ln z + \frac{a_{-2}}{-1} z + \frac{a_{-3}}{-2} z^2 + \frac{a_{-4}}{-3} z^3 + \dots \\ &\quad \text{(with } w = z^{-1}) \end{aligned}$$



$\phi(z) = \frac{a_{-3}}{-2} z^{-2}$   
 $f(z) = a_{-3} z^{-3}$

quadrupole field centered at North Pole

is quadratic field near South Pole

$\phi(w) = a_0 w^2$   
 $f(w) = a_1 w$

$$f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$



$$f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$

This  $m=1$ -pole constant- $a_{-1}$  formula is just the first in a series of *Laurent coefficient expressions*.

$$\dots a_{-3} = \frac{1}{2\pi i} \oint z^2 f(z)dz, \quad a_{-2} = \frac{1}{2\pi i} \oint z^1 f(z)dz, \quad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz, \quad a_0 = \frac{1}{2\pi i} \oint \frac{f(z)}{z} dz, \quad a_1 = \frac{1}{2\pi i} \oint \frac{f(z)}{z^2} dz, \dots$$

$$f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$

This  $m=1$ -pole constant- $a_{-1}$  formula is just the first in a series of *Laurent coefficient expressions*.

$$\dots a_{-3} = \frac{1}{2\pi i} \oint z^2 f(z)dz, \quad a_{-2} = \frac{1}{2\pi i} \oint z^1 f(z)dz, \quad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz, \quad a_0 = \frac{1}{2\pi i} \oint \frac{f(z)}{z} dz, \quad a_1 = \frac{1}{2\pi i} \oint \frac{f(z)}{z^2} dz, \dots$$

Source analysis starts with 1-pole loop integrals  $\oint z^{-1}dz = 2\pi i$  or, with origin shifted  $\oint (z-a)^{-1}dz = 2\pi i$ .

$$f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$

This  $m=1$ -pole constant- $a_{-1}$  formula is just the first in a series of *Laurent coefficient expressions*.

$$\dots a_{-3} = \frac{1}{2\pi i} \oint z^2 f(z)dz, \quad a_{-2} = \frac{1}{2\pi i} \oint z^1 f(z)dz, \quad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz, \quad a_0 = \frac{1}{2\pi i} \oint \frac{f(z)}{z} dz, \quad a_1 = \frac{1}{2\pi i} \oint \frac{f(z)}{z^2} dz, \dots$$

Source analysis starts with 1-pole loop integrals  $\oint z^{-1}dz = 2\pi i$  or, with origin shifted  $\oint (z-a)^{-1}dz = 2\pi i$ .

They hold for any loop about point- $a$ . Function  $f(z)$  is just  $f(a)$  on a *tiny* circle around point- $a$ .

(assume *tiny* circle around  $z=a$ )

$$\oint \frac{f(z)}{z-a} dz = \oint \frac{f(a)}{z-a} dz = f(a) \oint \frac{1}{z-a} dz = 2\pi i f(a)$$

(but any contour that doesn't "touch  $a$  gives same answer)

$$f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$

This  $m=1$ -pole constant- $a_{-1}$  formula is just the first in a series of *Laurent coefficient expressions*.

$$\dots a_{-3} = \frac{1}{2\pi i} \oint z^2 f(z)dz, \quad a_{-2} = \frac{1}{2\pi i} \oint z^1 f(z)dz, \quad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz, \quad a_0 = \frac{1}{2\pi i} \oint \frac{f(z)}{z} dz, \quad a_1 = \frac{1}{2\pi i} \oint \frac{f(z)}{z^2} dz, \dots$$

Source analysis starts with 1-pole loop integrals  $\oint z^{-1}dz = 2\pi i$  or, with origin shifted  $\oint (z-a)^{-1}dz = 2\pi i$ .

They hold for any loop about point- $a$ . Function  $f(z)$  is just  $f(a)$  on a *tiny* circle around point- $a$ .

(assume *tiny* circle around  $z=a$ )

$$\oint \frac{f(z)}{z-a} dz = \oint \frac{f(a)}{z-a} dz = f(a) \oint \frac{1}{z-a} dz = 2\pi i f(a)$$

(but any contour that doesn't "touch  $a$  gives same answer)

$$f(a) = \frac{1}{2\pi i} \oint \frac{f(z)}{z-a} dz$$

The  $f(a)$  result is called a *Cauchy integral*.

$$f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$

This  $m=1$ -pole constant- $a_{-1}$  formula is just the first in a series of *Laurent coefficient expressions*.

$$\dots a_{-3} = \frac{1}{2\pi i} \oint z^2 f(z)dz, \quad a_{-2} = \frac{1}{2\pi i} \oint z^1 f(z)dz, \quad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz, \quad a_0 = \frac{1}{2\pi i} \oint \frac{f(z)}{z} dz, \quad a_1 = \frac{1}{2\pi i} \oint \frac{f(z)}{z^2} dz, \dots$$

Source analysis starts with 1-pole loop integrals  $\oint z^{-1}dz = 2\pi i$  or, with origin shifted  $\oint (z-a)^{-1}dz = 2\pi i$ .

They hold for any loop about point- $a$ . Function  $f(z)$  is just  $f(a)$  on a *tiny* circle around point- $a$ .

(assume *tiny* circle around  $z=a$ )

$$\oint \frac{f(z)}{z-a} dz = \oint \frac{f(a)}{z-a} dz = f(a) \oint \frac{1}{z-a} dz = 2\pi i f(a)$$

(but any contour that doesn't "touch  $a$  gives same answer)

$$f(a) = \frac{1}{2\pi i} \oint \frac{f(z)}{z-a} dz$$

The  $f(a)$  result is called a *Cauchy integral*. Then repeated  $a$ -derivatives gives a sequence of them.

$$\frac{df(a)}{da} = \frac{1}{2\pi i} \oint \frac{f(z)}{(z-a)^2} dz,$$

$$f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$

This  $m=1$ -pole constant- $a_{-1}$  formula is just the first in a series of *Laurent coefficient expressions*.

$$\dots a_{-3} = \frac{1}{2\pi i} \oint z^2 f(z)dz, \quad a_{-2} = \frac{1}{2\pi i} \oint z^1 f(z)dz, \quad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz, \quad a_0 = \frac{1}{2\pi i} \oint \frac{f(z)}{z} dz, \quad a_1 = \frac{1}{2\pi i} \oint \frac{f(z)}{z^2} dz, \dots$$

Source analysis starts with 1-pole loop integrals  $\oint z^{-1}dz = 2\pi i$  or, with origin shifted  $\oint (z-a)^{-1}dz = 2\pi i$ .

They hold for any loop about point- $a$ . Function  $f(z)$  is just  $f(a)$  on a *tiny* circle around point- $a$ .

(assume *tiny* circle around  $z=a$ )

$$\oint \frac{f(z)}{z-a} dz = \oint \frac{f(a)}{z-a} dz = f(a) \oint \frac{1}{z-a} dz = 2\pi i f(a)$$

(but any contour that doesn't "touch  $a$  gives same answer)

$$f(a) = \frac{1}{2\pi i} \oint \frac{f(z)}{z-a} dz$$

The  $f(a)$  result is called a *Cauchy integral*. Then repeated  $a$ -derivatives gives a sequence of them.

$$\frac{df(a)}{da} = \frac{1}{2\pi i} \oint \frac{f(z)}{(z-a)^2} dz, \quad \frac{d^2 f(a)}{da^2} = \frac{2}{2\pi i} \oint \frac{f(z)}{(z-a)^3} dz,$$

$$f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$

This  $m=1$ -pole constant- $a_{-1}$  formula is just the first in a series of *Laurent coefficient expressions*.

$$\dots a_{-3} = \frac{1}{2\pi i} \oint z^2 f(z)dz, \quad a_{-2} = \frac{1}{2\pi i} \oint z^1 f(z)dz, \quad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz, \quad a_0 = \frac{1}{2\pi i} \oint \frac{f(z)}{z} dz, \quad a_1 = \frac{1}{2\pi i} \oint \frac{f(z)}{z^2} dz, \dots$$

Source analysis starts with 1-pole loop integrals  $\oint z^{-1}dz = 2\pi i$  or, with origin shifted  $\oint (z-a)^{-1}dz = 2\pi i$ .

They hold for any loop about point- $a$ . Function  $f(z)$  is just  $f(a)$  on a *tiny* circle around point- $a$ .

(assume *tiny* circle around  $z=a$ )

$$\oint \frac{f(z)}{z-a} dz = \oint \frac{f(a)}{z-a} dz = f(a) \oint \frac{1}{z-a} dz = 2\pi i f(a)$$

(but any contour that doesn't "touch  $a$  gives same answer)

$$f(a) = \frac{1}{2\pi i} \oint \frac{f(z)}{z-a} dz$$

The  $f(a)$  result is called a *Cauchy integral*. Then repeated  $a$ -derivatives gives a sequence of them.

$$\frac{df(a)}{da} = \frac{1}{2\pi i} \oint \frac{f(z)}{(z-a)^2} dz, \quad \frac{d^2 f(a)}{da^2} = \frac{2}{2\pi i} \oint \frac{f(z)}{(z-a)^3} dz, \quad \frac{d^3 f(a)}{da^3} = \frac{3!}{2\pi i} \oint \frac{f(z)}{(z-a)^4} dz$$



$$f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$

This  $m=1$ -pole constant- $a_{-1}$  formula is just the first in a series of *Laurent coefficient expressions*.

$$\dots a_{-3} = \frac{1}{2\pi i} \oint z^2 f(z)dz, \quad a_{-2} = \frac{1}{2\pi i} \oint z^1 f(z)dz, \quad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz, \quad a_0 = \frac{1}{2\pi i} \oint \frac{f(z)}{z} dz, \quad a_1 = \frac{1}{2\pi i} \oint \frac{f(z)}{z^2} dz, \dots$$

Source analysis starts with 1-pole loop integrals  $\oint z^{-1}dz = 2\pi i$  or, with origin shifted  $\oint (z-a)^{-1}dz = 2\pi i$ .

They hold for any loop about point- $a$ . Function  $f(z)$  is just  $f(a)$  on a *tiny* circle around point- $a$ .

(assume *tiny* circle around  $z=a$ )

$$\oint \frac{f(z)}{z-a} dz = \oint \frac{f(a)}{z-a} dz = f(a) \oint \frac{1}{z-a} dz = 2\pi i f(a)$$

(but any contour that doesn't "touch  $a$  gives same answer)

$$f(a) = \frac{1}{2\pi i} \oint \frac{f(z)}{z-a} dz$$

The  $f(a)$  result is called a *Cauchy integral*. Then repeated  $a$ -derivatives gives a sequence of them.

$$\frac{df(a)}{da} = \frac{1}{2\pi i} \oint \frac{f(z)}{(z-a)^2} dz, \quad \frac{d^2 f(a)}{da^2} = \frac{2}{2\pi i} \oint \frac{f(z)}{(z-a)^3} dz, \quad \frac{d^3 f(a)}{da^3} = \frac{3!}{2\pi i} \oint \frac{f(z)}{(z-a)^4} dz, \quad \dots, \quad \frac{d^n f(a)}{da^n} = \frac{n!}{2\pi i} \oint \frac{f(z)}{(z-a)^{n+1}} dz$$

$$f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$

This  $m=1$ -pole constant- $a_{-1}$  formula is just the first in a series of *Laurent coefficient expressions*.

$$\dots a_{-3} = \frac{1}{2\pi i} \oint z^2 f(z)dz, \quad a_{-2} = \frac{1}{2\pi i} \oint z^1 f(z)dz, \quad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz, \quad a_0 = \frac{1}{2\pi i} \oint \frac{f(z)}{z} dz, \quad a_1 = \frac{1}{2\pi i} \oint \frac{f(z)}{z^2} dz, \dots$$

Source analysis starts with 1-pole loop integrals  $\oint z^{-1}dz = 2\pi i$  or, with origin shifted  $\oint (z-a)^{-1}dz = 2\pi i$ .

They hold for any loop about point- $a$ . Function  $f(z)$  is just  $f(a)$  on a *tiny* circle around point- $a$ .

(assume *tiny* circle around  $z=a$ )

$$\oint \frac{f(z)}{z-a} dz = \oint \frac{f(a)}{z-a} dz = f(a) \oint \frac{1}{z-a} dz = 2\pi i f(a)$$

(but any contour that doesn't "touch  $a$  gives same answer)

$$f(a) = \frac{1}{2\pi i} \oint \frac{f(z)}{z-a} dz$$

The  $f(a)$  result is called a *Cauchy integral*. Then repeated  $a$ -derivatives gives a sequence of them.

$$\frac{df(a)}{da} = \frac{1}{2\pi i} \oint \frac{f(z)}{(z-a)^2} dz, \quad \frac{d^2 f(a)}{da^2} = \frac{2}{2\pi i} \oint \frac{f(z)}{(z-a)^3} dz, \quad \frac{d^3 f(a)}{da^3} = \frac{3!}{2\pi i} \oint \frac{f(z)}{(z-a)^4} dz, \quad \dots, \quad \frac{d^n f(a)}{da^n} = \frac{n!}{2\pi i} \oint \frac{f(z)}{(z-a)^{n+1}} dz$$

This leads to a general *Taylor-Laurent* power series expansion of function  $f(z)$  around point- $a$ .

$$f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$

This  $m=1$ -pole constant- $a_{-1}$  formula is just the first in a series of *Laurent coefficient expressions*.

$$\dots a_{-3} = \frac{1}{2\pi i} \oint z^2 f(z)dz, \quad a_{-2} = \frac{1}{2\pi i} \oint z^1 f(z)dz, \quad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz, \quad a_0 = \frac{1}{2\pi i} \oint \frac{f(z)}{z} dz, \quad a_1 = \frac{1}{2\pi i} \oint \frac{f(z)}{z^2} dz, \dots$$

Source analysis starts with 1-pole loop integrals  $\oint z^{-1}dz = 2\pi i$  or, with origin shifted  $\oint (z-a)^{-1}dz = 2\pi i$ .

They hold for any loop about point- $a$ . Function  $f(z)$  is just  $f(a)$  on a *tiny* circle around point- $a$ .

(assume *tiny* circle around  $z=a$ )

$$\oint \frac{f(z)}{z-a} dz = \oint \frac{f(a)}{z-a} dz = f(a) \oint \frac{1}{z-a} dz = 2\pi i f(a)$$

(but any contour that doesn't "touch  $a$  gives same answer)

$$f(a) = \frac{1}{2\pi i} \oint \frac{f(z)}{z-a} dz$$

The  $f(a)$  result is called a *Cauchy integral*. Then repeated  $a$ -derivatives gives a sequence of them.

$$\frac{df(a)}{da} = \frac{1}{2\pi i} \oint \frac{f(z)}{(z-a)^2} dz, \quad \frac{d^2 f(a)}{da^2} = \frac{2}{2\pi i} \oint \frac{f(z)}{(z-a)^3} dz, \quad \frac{d^3 f(a)}{da^3} = \frac{3!}{2\pi i} \oint \frac{f(z)}{(z-a)^4} dz, \quad \dots, \quad \frac{d^n f(a)}{da^n} = \frac{n!}{2\pi i} \oint \frac{f(z)}{(z-a)^{n+1}} dz$$

This leads to a general *Taylor-Laurent* power series expansion of function  $f(z)$  around point- $a$ .

$$f(z) = \sum_{n=-\infty}^{\infty} a_n (z-a)^n \qquad \text{where : } a_n = \frac{1}{2\pi i} \oint \frac{f(z)}{(z-a)^{n+1}} dz \left( = \frac{1}{n!} \frac{d^n f(a)}{da^n} \quad \text{for : } n \geq 0 \right)$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$

This  $m=1$ -pole constant- $a_{-1}$  formula is just the first in a series of *Laurent coefficient expressions*.

$$\dots a_{-3} = \frac{1}{2\pi i} \oint z^2 f(z)dz, \quad a_{-2} = \frac{1}{2\pi i} \oint z^1 f(z)dz, \quad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz, \quad a_0 = \frac{1}{2\pi i} \oint \frac{f(z)}{z} dz, \quad a_1 = \frac{1}{2\pi i} \oint \frac{f(z)}{z^2} dz, \dots$$

Source analysis starts with 1-pole loop integrals  $\oint z^{-1}dz = 2\pi i$  or, with origin shifted  $\oint (z-a)^{-1}dz = 2\pi i$ .

They hold for any loop about point- $a$ . Function  $f(z)$  is just  $f(a)$  on a *tiny* circle around point- $a$ .

(assume *tiny* circle around  $z=a$ )

$$\oint \frac{f(z)}{z-a} dz = \oint \frac{f(a)}{z-a} dz = f(a) \oint \frac{1}{z-a} dz = 2\pi i f(a)$$

(but any contour that doesn't "touch  $a$  gives same answer)

$$f(a) = \frac{1}{2\pi i} \oint \frac{f(z)}{z-a} dz$$

The  $f(a)$  result is called a *Cauchy integral*. Then repeated  $a$ -derivatives gives a sequence of them.

$$\frac{df(a)}{da} = \frac{1}{2\pi i} \oint \frac{f(z)}{(z-a)^2} dz, \quad \frac{d^2 f(a)}{da^2} = \frac{2}{2\pi i} \oint \frac{f(z)}{(z-a)^3} dz, \quad \frac{d^3 f(a)}{da^3} = \frac{3!}{2\pi i} \oint \frac{f(z)}{(z-a)^4} dz, \dots, \frac{d^n f(a)}{da^n} = \frac{n!}{2\pi i} \oint \frac{f(z)}{(z-a)^{n+1}} dz$$



This leads to a general *Taylor-Laurent* power series expansion of function  $f(z)$  around point- $a$ .

$$f(z) = \sum_{n=-\infty}^{\infty} a_n (z-a)^n \qquad \text{where : } a_n = \frac{1}{2\pi i} \oint \frac{f(z)}{(z-a)^{n+1}} dz \left( = \frac{1}{n!} \frac{d^n f(a)}{da^n} \quad \text{for : } n \geq 0 \right)$$


(*quadrupole*)<sub>0</sub> (dipole)<sub>0</sub> (monopole) (dipole)<sub>∞</sub> (*quadrupole*)<sub>∞</sub> (*octapole*)<sub>∞</sub> (*hexadecapole*)<sub>∞</sub> ...

$$f(z) = \dots a_{-3}z^{-3} + \underbrace{a_{-2}z^{-2}}_{\text{dipole moment}} + \underbrace{a_{-1}z^{-1}}_{\text{monopole moment}} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

$w(z) = z^2$  gives parabolic OCC



Inverse:  $z(w) = w^{1/2}$  gives hyperbolic OCC



$w = (u + iv) = z^2 = (x + iy)^2$  is analytic function of  $z$  and  $w$

Expansion:  $u = x^2 - y^2$  and  $v = 2xy$  may be solved using  $|w| = |z^2| = |z|^2$

Expansion:  $|w| = \sqrt{u^2 + v^2} = x^2 + y^2 = |z|^2$

Solution:  $x^2 = \frac{u + \sqrt{u^2 + v^2}}{2}$   $y^2 = \frac{-u + \sqrt{u^2 + v^2}}{2}$

$$\begin{pmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{pmatrix} = \begin{pmatrix} \bar{\mathbf{E}}^u \\ \bar{\mathbf{E}}^v \end{pmatrix} = \begin{pmatrix} 2x & -2y \\ +2y & 2x \end{pmatrix}$$

$$\begin{pmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{pmatrix} = \begin{pmatrix} \bar{\mathbf{E}}_u & \bar{\mathbf{E}}_v \end{pmatrix} = \frac{\begin{pmatrix} 2x & +2y \\ -2y & 2x \end{pmatrix}}{4(x^2 + y^2)}$$