

Lecture 13  
Wed. 10.09.2019

# Complex Variables, Series, and Field Coordinates II.

(Ch. 10 of Unit 1)

## 1. The Story of $e$ (A Tale of Great \$Interest\$)

*How good are those power series?*

*Taylor-Maclaurin series, imaginary interest, and complex exponentials*

## 2. What good are complex exponentials?

*Easy trig*

*Easy 2D vector analysis*

*Easy oscillator phase analysis*

*Easy rotation and “dot” or “cross” products*

## 3. Easy 2D vector calculus

*Easy 2D vector derivatives*

*Easy 2D source-free field theory*

*Easy 2D vector field-potential theory*

## 4. Riemann-Cauchy relations (What's analytic? What's not?)

*Easy 2D curvilinear coordinate discovery*

*Easy 2D circulation and flux integrals*

*Easy 2D monopole, dipole, and  $2^n$ -pole analysis*

*Easy  $2^n$ -multipole field and potential expansion*

*Easy stereo-projection visualization*

*Cauchy integrals, Laurent-Maclaurin series*

## 5. Mapping and Non-analytic 2D source field analysis

1. Complex numbers provide "automatic trigonometry"

2. Complex numbers add like vectors.

3. Complex exponentials  $Ae^{-i\omega t}$  track position and velocity using Phasor Clock.

4. Complex products provide 2D rotation operations.

5. Complex products provide 2D “dot”( $\cdot$ ) and “cross”(x) products.

*(Review of topics in Lect. 12)*

6. Complex derivative contains “divergence”( $\nabla \cdot \mathbf{F}$ ) and “curl”( $\nabla \times \mathbf{F}$ ) of 2D vector field

7. Invent source-free 2D vector fields [ $\nabla \cdot \mathbf{F}=0$  and  $\nabla \times \mathbf{F}=0$ ]

8. Complex potential  $\phi$  contains “scalar”( $\mathbf{F}=\nabla \phi$ ) and “vector”( $\mathbf{F}=\nabla \times \mathbf{A}$ ) potentials

*The half-n'-half results: (Riemann-Cauchy Derivative Relations)*

9. Complex potentials define 2D Orthogonal Curvilinear Coordinates (OCC) of field

10. Complex integrals  $\int f(z)dz$  count 2D “circulation”( $\int \mathbf{F} \cdot d\mathbf{r}$ ) and “flux”( $\int \mathbf{F} \times d\mathbf{r}$ )

11. Complex integrals define 2D **monopole** fields and potentials

12. Complex derivatives give 2D dipole fields

13. More derivatives give 2D  $2^N$ -pole fields...

14. ...and  $2^N$ -pole multipole expansions of fields and potentials...

15. ...and Laurent Series...

16. ...and non-analytic source analysis.

17. ...and mapping...

Lecture 13 Wed. 10.09.19

Starts review here

Lect. 12

ended here

# This Lecture's Reference Link Listing

[Web Resources - front page](#)

[UAF Physics UTube channel](#)

[Quantum Theory for the Computer Age](#)

[Principles of Symmetry, Dynamics, and Spectroscopy](#)

[Classical Mechanics with a Bang!](#)

[Modern Physics and its Classical Foundations](#)

[2017 Group Theory for QM](#)

[2018 Adv CM](#)

[2018 AMOP](#)

[2019 Advanced Mechanics](#)

## Lectures #12 through #13

*In reverse order*

**Pirelli Relativity Challenge (Introduction level) - Visualizing Waves:**

[Using Earth as a clock,](#)

[Tesla's AC Phasors ,](#)

[Phasors using complex numbers.](#)

[CM wBang Unit 1 - Chapter 10: Calculus of exponentials, logarithms, and complex fields, pdf\\_page=135](#)

[RelaWavity Web Simulation - Unit Circle and Hyperbola \(Mixed labeling\)](#)

## Select, exciting, and related Research & Articles of Interest

*(Many of these may be just beyond this course, but are included to lend added insight):*

[Clifford Algebra And The Projective Model Of Homogeneous Metric Spaces - Foundations - Sokolov-x-2013](#)

[Geometric Algebra 3 - Complex Numbers - MacDonald-yt-2015](#)

[Biquaternion -Complexified Quaternion- Roots of -1 - Sangwine-x-2015](#)

[An Introduction to Clifford Algebras and Spinors - Vaz-Rocha-op-2016](#)

[Unified View on Complex Numbers and Quaternions- Bongardt-wcmms-2015](#)

[Complex Functions and the Cauchy-Riemann Equations - complex2 - Friedman-columbia-2019](#)

Excerpts from the [Geometric Algebra- A Guided Tour through Space and Time - Reimer-www-2019- Page 44-47](#)

(Preliminary Draft)

## Past Articles of Interest:

[An sp-hybridized Molecular Carbon Allotrope- cyclo-18-carbon - Kaiser-s-2019](#)

[An Atomic-Scale View of Cyclocarbon Synthesis - Maier-s-2019](#)

[Discovery Of Topological Weyl Fermion Lines And Drumhead Surface States in a Room Temperature Magnet - Belopolski-s-2019](#)

["Weyl"ing away Time-reversal Symmetry - Neto-s-2019](#)

[Non-Abelian Band Topology in Noninteracting Metals - Wu-s-2019](#)

[What Industry Can Teach Academia - Mao-s-2019](#)

[Rovibrational quantum state resolution of the C60 fullerene - Changala-Ye-s-2019 \(Alt\)](#)

[A Degenerate Fermi Gas of Polar molecules - DeMarco-s-2019](#)

An assist from [Physics Girl](#) (YouTube Channel):

Posted this year:

[How to Make VORTEX RINGS in a Pool](#)

Crazy pool vortex (new inclusion with more background)

[Crazy pool vortex - pg-yt-2014](#)

Posting with the best visuals:

[Fun with Vortex Rings in the Pool - pg-yt-2014](#)

*She covers it beautifully!*

# Running Reference Link Listing

## Lectures #11 through #7

*In reverse order*

### Eric J Heller Gallery:

[Main portal](#), [Consonance and Dissonance II](#), [Bessel 21](#), [Chladni](#)

[The Semiclassical Way to Molecular Spectroscopy - Heller-acs-1981](#)  
[Quantum dynamical tunneling in bound states - Davis-Heller-jcp-1981](#)

[Pendulum Web Simulation](#)

[Cycloidulum Web Simulation](#)

**Links to previous lecture:** [Page=74](#), [Page=75](#), [Page=79](#)

[Pendulum Web Sim](#)

[Cycloidulum Web Sim](#)

**JerkIt Web Simulations:** [Basic/Generic](#); [Inverted](#), [FVPlot](#)

[CMwithBang Lecture 8, page=20](#)

[WWW.sciencenewsforstudents.org: Cassini - Saturnian polar vortex](#)

“RelaWavity” Web Simulations:

[2-CW laser wave](#), [Lagrangian vs Hamiltonian](#),

[Physical Terms Lagrangian L\(u\) vs Hamiltonian H\(p\)](#)

[CoulIt Web Simulation of the Volcanoes of Io](#)

[BohrIt Multi-Panel Plot:](#)

[Relativistically shifted Time-Space plots of 2 CW light waves](#)

### BoxIt Web Simulations:

[Generic/Default](#)

[Most Basic A-Type](#)

[Basic A-Type w/reference lines](#)

[Basic A-Type A-Type with Potential energy](#)

[A-Type with Potential energy and Stokes Plot](#)

[A-Type w/3 time rates of change](#)

[A-Type w/3 time rates of change with Stokes Plot](#)

[B-Type \(A=1.0, B=-0.05, C=0.0, D=1.0\)](#)

### RelaWavity Web Elliptical Motion Simulations:

[Orbits with b/a=0.125](#)

[Orbits with b/a=0.5](#)

[Orbits with b/a=0.7](#)

[Exegesis with b/a=0.125](#)

[Exegesis with b/a=0.5](#)

[Exegesis with b/a=0.7](#)

[Contact Ellipsometry](#)

### CoulIt Web Simulations:

[Basic/Generic](#)

[Exploding Starlet](#)

[Volcanoes of Io \(Color Quantized\)](#)

### JerkIt Web Simulations:

[Basic/Generic](#)

[Catcher in the Eye - IHO with Linear Hooke perturbation - Force-potential-Velocity Plot](#)

### OscillatorPE Web Simulation:

[Coulomb-Newton-Inverse Square](#),

[Hooke-Isotropic Harmonic](#),

[Pendulum-Circular Constraint](#)

[AMOP Ch 0 Space-Time Symmetry - 2019](#)

[Seminar at Rochester Institute of Optics, Aux. slides-2018](#)

[NASA Astronomy Picture of the Day -](#)

[Io: The Prometheus Plume \(Just Image\)](#)

[NASA Galileo - Io's Alien Volcanoes](#)

[New Horizons - Volcanic Eruption Plume on Jupiter's moon IO](#)

[NASA Galileo - A Hawaiian-Style Volcano on Io](#)

[Pirelli Site: Phasors animation](#)

[CMwithBang Lecture #6, page=70 \(9.10.18\)](#)

### Select, exciting, and related Research & Articles of Interest:

[Burning a hole in reality—design for a new laser may be powerful enough to pierce space-time - Sumner-KOS-2019](#)

[Trampoline mirror may push laser pulse through fabric of the Universe - Lee-ArsTechnica-2019](#)

[Achieving Extreme Light Intensities using Optically Curved Relativistic Plasma Mirrors - Vincenti-prl-2019](#)

[A Soft Matter Computer for Soft Robots - Garrad-sr-2019](#)

[Correlated Insulator Behaviour at Half-Filling in Magic-Angle Graphene Superlattices - cao-n-2018](#)

[Sorting ultracold atoms in a three-dimensional optical lattice in a realization of Maxwell's Demon - Kumar-n-2018](#)

[Synthetic three-dimensional atomic structures assembled atom by atom - Barredo-n-2018](#)

Older ones:

[Wave-particle duality of C60 molecules - Arndt-ltn-1999](#)

[Optical Vortex Knots - One Photon At A Time - Tempone-Wiltshire-Sr-2018](#)

[Baryon Deceleration by Strong Chromofields in Ultrarelativistic](#)

[Nuclear Collisions - Mishustin-PhysRevC-2007, APS Link & Abstract](#)

[Hadronic Molecules - Guo-x-2017](#)

[Hidden-charm pentaquark and tetraquark states - Chen-pr-2016](#)



# Running Reference Link Listing

## Lectures #6 through #1

In reverse order

[RelaWavity Web Simulation: Contact Ellipsometry](#)

[BoxIt Web Simulation: Elliptical Motion \(A-Type\)](#)

[CMwBang Course: Site Title Page](#)

[Pirelli Relativity Challenge: Describing Wave Motion With Complex Phasors](#)

[UAF Physics UTube channel](#)

[Velocity Amplification in Collision Experiments Involving Superballs - Harter, 1971](#)

[MIT OpenCourseWare: High School/Physics/Impulse and Momentum](#)

[Hubble Site: Supernova - SN 1987A](#)

### **BounceIt Web Animation - Scenarios:**

[49:1 y vs t, 49:1 V2 vs V1, 1:500:1 - 1D Gas Model w/ faux restorative force \(Cool\),](#)

[1:500:1 - 1D Gas \(Warm\), 1:500:1 - 1D Gas Model \(Cool, Zoomed in\),](#)

[Farey Sequence - Wolfram](#)

[Fractions - Ford-AMM-1938](#)

### **Monstermash BounceIt Animations:**

[1000:1 - V2 vs V1, 1000:1 with t vs x - Minkowski Plot](#)

[Quantum Revivals of Morse Oscillators and Farey-Ford Geometry - Li-Harter-2013](#)

[Quantum Revivals of Morse Oscillators and Farey-Ford Geometry - Li-Harter-cpl-2015](#)

[Quant. Revivals of Morse Oscillators and Farey-Ford Geom. - Harter-Li-CPL-2015 \(Publ.\)](#)

[Velocity Amplification in Collision Experiments Involving Superballs-Harter-1971](#)

### **WaveIt Web Animation - Scenarios:**

[Quantum Carpet, Quantum Carpet wMBars,](#)

[Quantum Carpet BCar, Quantum Carpet BCar wMBars](#)

[Wave Node Dynamics and Revival Symmetry in Quantum Rotors - Harter-JMS-2001](#)

[Wave Node Dynamics and Revival Symmetry in Quantum Rotors - Harter-jms-2001 \(Publ.\)](#)

[AJP article on superball dynamics](#)

[AAPT Summer Reading List](#)

[Scitation.org - AIP publications](#)

[HarterSoft Youtube Channel](#)

### **BounceIt Web Animation - Scenarios:**

[Generic Scenario: 2-Balls dropped no Gravity \(7:1\) - V vs V Plot \(Power=4\)](#)

[1-Ball dropped w/Gravity=0.5 w/Potential Plot: Power=1, Power=4](#)

[7:1 - V vs V Plot: Power=1](#)

[3-Ball Stack \(10:3:1\) w/Newton plot \(y vs t\) - Power=4](#)

[3-Ball Stack \(10:3:1\) w/Newton plot \(y vs t\) - Power=1](#)

[3-Ball Stack \(10:3:1\) w/Newton plot \(y vs t\) - Power=1 w/Gaps](#)

[4-Ball Stack \(27:9:3:1\) w/Newton plot \(y vs t\) - Power=4](#)

[4-Newton's Balls \(1:1:1:1\) w/Newtonian plot \(y vs t\) - Power=4 w/Gaps](#)

[6-Ball Totally Inelastic \(1:1:1:1:1:1\) w/Gaps: Newtonian plot \(t vs x\), V6 vs V5 plot](#)

[5-Ball Totally Inelastic Pile-up w/ 5-Stationary-Balls - Minkowski plot \(t vs x1\) w/Gaps](#)

[1-Ball Totally Inelastic Pile-up w/ 5-Stationary-Balls - Vx2 vs Vx1 plot w/Gaps](#)

### **BounceIt Dual plots**

**$m_1:m_2 = 3:1$**

[v2 vs v1 and V2 vs V1, \(v1, v2\)=\(1, 0.1\), \(v1, v2\)=\(1, 0\)](#)

[y2 vs y1 plots: \(v1, v2\)=\(1, 0.1\), \(v1, v2\)=\(1, 0\), \(v1, v2\)=\(1, -1\)](#)

[Estrangian plot V2 vs V1: \(v1, v2\)=\(0, 1\), \(v1, v2\)=\(1, -1\)](#)

**$m_1:m_2 = 4:1$**

[v2 vs v1, y2 vs y1](#)

**$m_1:m_2 = 100:1$ , (v1, v2)=(1, 0): V2 vs V1 Estrangian plot, y2 vs y1 plot**

[With g=0 and 70:10 mass ratio](#)

[With non zero g, velocity dependent damping and mass ratio of 70:35](#)

[M1=49, M2=1 with Newtonian time plot](#)

[M1=49, M2=1 with V2 vs V1 plot](#)

[Example with friction](#)

[Low force constant with drag displaying a Pass-thru, Fall-Thru, Bounce-Off](#)

[m1:m2= 3:1 and \(v1, v2\) = \(1, 0\) Comparison with Estrangian](#)

X2 paper: [Velocity Amplification in Collision Experiments Involving Superballs - Harter, et. al. 1971 \(pdf\)](#)

Car Collision Web Simulator: <https://modphys.hosted.uark.edu/markup/CMMotionWeb.html>

Superball Collision Web Simulator: <https://modphys.hosted.uark.edu/markup/BounceItWeb.html>; with Scenarios: [1007](#)

[BounceIt web simulation with g=0 and 70:10 mass ratio](#)

[With non zero g, velocity dependent damping and mass ratio of 70:35](#)

Elastic Collision Dual Panel Space vs Space: [Space vs Time \(Newton\)](#), [Time vs. Space\(Minkowski\)](#)

Inelastic Collision Dual Panel Space vs Space: [Space vs Time \(Newton\)](#), [Time vs. Space\(Minkowski\)](#)

Matrix Collision Simulator: [M1=49, M2=1 V2 vs V1 plot](#) <<Under Construction>>

More Advanced QM and classical references will soon be available through our: [Mechanics References Page](#)

(Now in Development)



6. Complex derivative contains “divergence” ( $\nabla \cdot \mathbf{F}$ ) and “curl” ( $\nabla \times \mathbf{F}$ ) of 2D vector field

Relation of  $(z, z^*)$  to  $(x = \text{Re}z, y = \text{Im}z)$  defines a  $z$ -derivative  $\frac{df}{dz}$  and “star”  $z^*$ -derivative.  $\frac{df}{dz^*}$

$$z = x + iy$$

$$x = \frac{1}{2} (z + z^*)$$

$$z^* = x - iy$$

$$y = \frac{1}{2i} (z - z^*)$$

$$\frac{df}{dz} = \frac{\partial x}{\partial z} \frac{\partial f}{\partial x} + \frac{\partial y}{\partial z} \frac{\partial f}{\partial y} = \frac{1}{2} \frac{\partial f}{\partial x} - \frac{i}{2} \frac{\partial f}{\partial y}$$

$$\frac{d}{dz} = \frac{1}{2} \frac{\partial}{\partial x} - \frac{i}{2} \frac{\partial}{\partial y}$$

$$\frac{df}{dz^*} = \frac{\partial x}{\partial z^*} \frac{\partial f}{\partial x} + \frac{\partial y}{\partial z^*} \frac{\partial f}{\partial y} = \frac{1}{2} \frac{\partial f}{\partial x} + \frac{i}{2} \frac{\partial f}{\partial y}$$

$$\frac{d}{dz^*} = \frac{1}{2} \frac{\partial}{\partial x} + \frac{i}{2} \frac{\partial}{\partial y}$$

Derivative chain-rule shows real part of  $\frac{df}{dz}$  has 2D divergence  $\nabla \cdot \mathbf{f}$  and imaginary part has curl  $\nabla \times \mathbf{f}$ .

$$\frac{df}{dz} = \frac{d}{dz} (f_x + i f_y) = \frac{1}{2} \left( \frac{\partial f}{\partial x} - i \frac{\partial f}{\partial y} \right) (f_x + i f_y) = \frac{1}{2} \left( \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial f_y}{\partial x} - \frac{\partial f_x}{\partial y} \right) = \frac{1}{2} \nabla \cdot \mathbf{f} + \frac{i}{2} |\nabla \times \mathbf{f}|_{Z \perp (x,y)}$$

7. Invent source-free 2D vector fields [ $\nabla \cdot \mathbf{F} = 0$  and  $\nabla \times \mathbf{F} = 0$ ]

We can invent *source-free 2D vector fields* that are both *zero-divergence* and *zero-curl*.

Take any function  $f(z)$ , conjugate it (change all  $i$ 's to  $-i$ ) to give  $f^*(z^*)$  for which  $\frac{df^*}{dz^*} = 0$

For example: if  $f(z) = a \cdot z$  then  $f^*(z^*) = a \cdot z^* = a(x - iy)$  is not function of  $z$  so it has zero  $z$ -derivative.

$\mathbf{F} = (F_x, F_y) = (f_x^*, f_y^*) = (a \cdot x, -a \cdot y)$  has *zero divergence*:  $\nabla \cdot \mathbf{F} = 0$  and has *zero curl*:  $|\nabla \times \mathbf{F}| = 0$ .

$$\nabla \cdot \mathbf{F} = \frac{\partial F_x}{\partial x} + \frac{\partial F_y}{\partial y} = \frac{\partial(ax)}{\partial x} + \frac{\partial(-ay)}{\partial y} = 0 \quad |\nabla \times \mathbf{F}|_{Z \perp (x,y)} = \frac{\partial F_y}{\partial x} - \frac{\partial F_x}{\partial y} = \frac{\partial(-ay)}{\partial x} - \frac{\partial(ax)}{\partial y} = 0$$

A *DFL* field  $\mathbf{F}$  (*Divergence-Free-Laminar*)

7. Invent source-free 2D vector fields [ $\nabla \cdot \mathbf{F} = 0$  and  $\nabla \times \mathbf{F} = 0$ ]


We can invent *source-free 2D vector fields* that are both *zero-divergence* and *zero-curl*.

Take any function  $f(z)$ , conjugate it (change all  $i$ 's to  $-i$ ) to give  $f^*(z^*)$  for which  $\frac{df^*}{dz} = 0$ .

For example: if  $f(z) = a \cdot z$  then  $f^*(z^*) = a \cdot z^* = a(x - iy)$  is not function of  $z$  so it has zero  $z$ -derivative.

$\mathbf{F} = (F_x, F_y) = (f^*_x, f^*_y) = (a \cdot x, -a \cdot y)$  has *zero divergence*:  $\nabla \cdot \mathbf{F} = 0$  and has *zero curl*:  $|\nabla \times \mathbf{F}| = 0$ .

$$\nabla \cdot \mathbf{F} = \frac{\partial F_x}{\partial x} + \frac{\partial F_y}{\partial y} = \frac{\partial(ax)}{\partial x} + \frac{\partial(-ay)}{\partial y} = 0 \qquad |\nabla \times \mathbf{F}|_{z \perp (x,y)} = \frac{\partial F_y}{\partial x} - \frac{\partial F_x}{\partial y} = \frac{\partial(-ay)}{\partial x} - \frac{\partial(ax)}{\partial y} = 0$$



precursor to  
Unit 1  
Fig. 10.7

$\mathbf{F} = (f^*_x, f^*_y) = (a \cdot x, -a \cdot y)$  is a *divergence-free laminar (DFL)* field.

8. Complex potential  $\phi$  contains “scalar”( $\mathbf{F}=\nabla\Phi$ ) and “vector”( $\mathbf{F}=\nabla\times\mathbf{A}$ ) potentials

Any *DFL* field  $\mathbf{F}$  is a gradient of a *scalar potential field*  $\Phi$  or a curl of a *vector potential field*  $\mathbf{A}$ .

$$\mathbf{F} = \nabla\Phi \qquad \mathbf{F} = \nabla\times\mathbf{A}$$


A *complex potential*  $\phi(z)=\Phi(x,y)+i\mathbf{A}(x,y)$  exists whose  $z$ -derivative is  $f(z)=d\phi/dz$ .

Its complex conjugate  $\phi^*(z^*)=\Phi(x,y)-i\mathbf{A}(x,y)$  has  $z^*$ -derivative  $f^*(z^*)=d\phi^*/dz^*$  giving *DFL* field  $\mathbf{F}$ .

To find  $\phi=\Phi+i\mathbf{A}$  integrate  $f(z)=a\cdot z$  to get  $\phi$  and isolate real ( $\text{Re } \phi = \Phi$ ) and imaginary ( $\text{Im } \phi = \mathbf{A}$ ) parts.

$$f(z) = \frac{d\phi}{dz} \Rightarrow \phi = \underbrace{\Phi}_{\frac{1}{2}a(x^2 - y^2)} + i \underbrace{\mathbf{A}}_{axy} = \int f \cdot dz = \int az \cdot dz = \frac{1}{2} az^2 = \frac{1}{2} a(x + iy)^2$$

*BONUS!*  
Get a free coordinate system!



Unit 1  
Fig. 10.7

The  $(\Phi, \mathbf{A})$  grid is a GCC coordinate system\*:

$$q^1 = \Phi = (x^2 - y^2)/2 = \text{const.}$$

$$q^2 = \mathbf{A} = (xy) = \text{const.}$$

\*Actually it's OCC.



8. (contd.) Complex potential  $\phi$  contains “scalar”( $\mathbf{F}=\nabla\Phi$ ) and “vector”( $\mathbf{F}=\nabla\times\mathbf{A}$ ) potentials

...and either one (or *half-n'-half!*) works just as well.

Derivative  $\frac{d\phi^*}{dz^*}$  has 2D gradient  $\nabla\Phi = \begin{pmatrix} \frac{\partial\Phi}{\partial x} \\ \frac{\partial\Phi}{\partial y} \end{pmatrix}$  of scalar  $\Phi$  and curl  $\nabla\times\mathbf{A} = \begin{pmatrix} \frac{\partial\mathbf{A}}{\partial y} \\ -\frac{\partial\mathbf{A}}{\partial x} \end{pmatrix}$  of vector  $\mathbf{A}$  (and they're equal!)

The *half-n'-half* result

$$\frac{d}{dz^*} \phi^* = \frac{d}{dz^*} (\Phi - i\mathbf{A}) = \frac{1}{2} \left( \frac{\partial}{\partial x} + i\frac{\partial}{\partial y} \right) (\Phi - i\mathbf{A}) = \frac{1}{2} \left( \frac{\partial\Phi}{\partial x} + i\frac{\partial\Phi}{\partial y} \right) + \frac{1}{2} \left( \frac{\partial\mathbf{A}}{\partial y} - i\frac{\partial\mathbf{A}}{\partial x} \right) = \frac{1}{2} \nabla\Phi + \frac{1}{2} \nabla\times\mathbf{A}$$

Note, *mathematician definition* of force field  $\mathbf{F} = +\nabla\Phi$  replaces usual physicist's definition  $\mathbf{F} = -\nabla\Phi$

Given  $\phi$ :

$$\phi = \Phi + i\mathbf{A} = \frac{1}{2} a(x^2 - y^2) + i axy$$

The *half-n'-half* result


find:

$$\nabla\Phi = \begin{pmatrix} \frac{\partial\Phi}{\partial x} \\ \frac{\partial\Phi}{\partial y} \end{pmatrix} = \begin{pmatrix} \frac{\partial}{\partial x} \frac{a}{2}(x^2 - y^2) \\ \frac{\partial}{\partial y} \frac{a}{2}(x^2 - y^2) \end{pmatrix} = \begin{pmatrix} ax \\ -ay \end{pmatrix} = \mathbf{F}$$

or find:

$$\nabla\times\mathbf{A} = \begin{pmatrix} \frac{\partial\mathbf{A}}{\partial y} \\ -\frac{\partial\mathbf{A}}{\partial x} \end{pmatrix} = \begin{pmatrix} \frac{\partial}{\partial y} axy \\ -\frac{\partial}{\partial x} axy \end{pmatrix} = \begin{pmatrix} ax \\ -ay \end{pmatrix} = \mathbf{F}$$

Scalar *static potential lines*  $\Phi = \text{const.}$  and vector *flux potential lines*  $\mathbf{A} = \text{const.}$  define *DFL field-net*.



The *half-n'-half* results

are called

*Riemann-Cauchy*

*Derivative Relations*

$$\frac{\partial\Phi}{\partial x} = \frac{\partial\mathbf{A}}{\partial y} \quad \text{is:} \quad \frac{\partial\text{Re}f(z)}{\partial x} = \frac{\partial\text{Im}f(z)}{\partial y}$$

$$\frac{\partial\Phi}{\partial y} = -\frac{\partial\mathbf{A}}{\partial x} \quad \text{is:} \quad \frac{\partial\text{Re}f(z)}{\partial y} = -\frac{\partial\text{Im}f(z)}{\partial x}$$

Review  $(z, z^*)$  to  $(x, y)$  transformation relations

(Review of topics in Lect.12)

$$z = x + iy \quad x = \frac{1}{2}(z + z^*)$$

$$z^* = x - iy \quad y = \frac{1}{2i}(z - z^*)$$

$$\frac{df}{dz} = \frac{\partial x}{\partial z} \frac{\partial f}{\partial x} + \frac{\partial y}{\partial z} \frac{\partial f}{\partial y} = \frac{1}{2} \frac{\partial f}{\partial x} + \frac{1}{2i} \frac{\partial f}{\partial y} = \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) f$$

$$\frac{df}{dz^*} = \frac{\partial x}{\partial z^*} \frac{\partial f}{\partial x} + \frac{\partial y}{\partial z^*} \frac{\partial f}{\partial y} = \frac{1}{2} \frac{\partial f}{\partial x} - \frac{1}{2i} \frac{\partial f}{\partial y} = \frac{1}{2} \left( \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right) f$$

Criteria for a field function  $f = f_x(x, y) + i f_y(x, y)$  to be an **analytic function  $f(z)$**  of  $z = x + iy$ :

First,  $f(z)$  must not be a function of  $z^* = x - iy$ , that is:  $\frac{df}{dz^*} = 0$

This implies  $f(z)$  satisfies differential equations known as the **Riemann-Cauchy conditions**

$$\frac{df}{dz^*} = 0 = \frac{1}{2} \left( \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right) (f_x + i f_y) = \frac{1}{2} \left( \frac{\partial f_x}{\partial x} - \frac{\partial f_y}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial f_y}{\partial x} + \frac{\partial f_x}{\partial y} \right) \text{ implies: } \frac{\partial f_x}{\partial x} = \frac{\partial f_y}{\partial y} \quad \text{and:} \quad \frac{\partial f_y}{\partial x} = -\frac{\partial f_x}{\partial y}$$

$$\frac{df}{dz} = \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) (f_x + i f_y) = \frac{1}{2} \left( \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial f_y}{\partial x} - \frac{\partial f_x}{\partial y} \right) = \frac{\partial f_x}{\partial x} + i \frac{\partial f_y}{\partial x} = \frac{\partial f_y}{\partial y} - i \frac{\partial f_x}{\partial y} = \frac{\partial}{\partial x} (f_x + i f_y) = \frac{\partial}{\partial iy} (f_x + i f_y)$$

Criteria for a field function  $f = f_x(x, y) + i f_y(x, y)$  to be an **analytic function  $f(z^*)$**  of  $z^* = x - iy$ :

First,  $f(z^*)$  must not be a function of  $z = x + iy$ , that is:  $\frac{df}{dz} = 0$

This implies  $f(z^*)$  satisfies differential equations we call **Anti-Riemann-Cauchy conditions**

$$\frac{df}{dz} = 0 = \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) (f_x + i f_y) = \frac{1}{2} \left( \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial f_y}{\partial x} - \frac{\partial f_x}{\partial y} \right) = \text{implies: } \frac{\partial f_x}{\partial x} = -\frac{\partial f_y}{\partial y} \quad \text{and:} \quad \frac{\partial f_y}{\partial x} = \frac{\partial f_x}{\partial y}$$

$$\frac{df}{dz^*} = \frac{1}{2} \left( \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right) (f_x + i f_y) = \frac{1}{2} \left( \frac{\partial f_x}{\partial x} - \frac{\partial f_y}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial f_y}{\partial x} + \frac{\partial f_x}{\partial y} \right) = \frac{\partial f_x}{\partial x} + i \frac{\partial f_y}{\partial x} = -\frac{\partial f_y}{\partial y} + i \frac{\partial f_x}{\partial y} = \frac{\partial}{\partial x} (f_x + i f_y) = -\frac{\partial}{\partial iy} (f_x + i f_y)$$





## 4. Riemann-Cauchy conditions *What's analytic? (...and what's not?)*

→ *Easy 2D circulation and flux integrals*

*Easy 2D curvilinear coordinate discovery*

*Easy 2D monopole, dipole, and  $2^n$ -pole analysis*

*Easy  $2^n$ -multipole field and potential expansion*

*Easy stereo-projection visualization*

9. Complex integrals  $\int f(z)dz$  count 2D “circulation” ( $\int \mathbf{F} \cdot d\mathbf{r}$ ) and “flux” ( $\int \mathbf{F} \times d\mathbf{r}$ )

Integral of  $f(z)$  between point  $z_1$  and point  $z_2$  is potential difference  $\Delta\phi = \phi(z_2) - \phi(z_1)$


$$\Delta\phi = \phi(z_2) - \phi(z_1) = \int_{z_1}^{z_2} f(z)dz = \underbrace{\Phi(x_2, y_2) - \Phi(x_1, y_1)}_{\Delta\Phi} + i \underbrace{[\mathbf{A}(x_2, y_2) - \mathbf{A}(x_1, y_1)]}_{\Delta\mathbf{A}}$$

$$\Delta\phi = \Delta\Phi + i \Delta\mathbf{A}$$


In *DFL*-field  $\mathbf{F}$ ,  $\Delta\phi$  is independent of the integration path  $z(t)$  connecting  $z_1$  and  $z_2$ .

$$\begin{aligned} \int f(z) dz &= \int \left( f^*(z^*) \right)^* dz = \int \left( f^*(z^*) \right)^* (dx + i dy) = \int \left( f_x^* + i f_y^* \right)^* (dx + i dy) = \int \left( f_x^* - i f_y^* \right) (dx + i dy) \\ &= \int (f_x^* dx + f_y^* dy) + i \int (f_x^* dy - f_y^* dx) \\ &= \int \mathbf{F} \cdot d\mathbf{r} + i \int \mathbf{F} \times d\mathbf{r} \cdot \hat{\mathbf{e}}_z \\ &= \int \mathbf{F} \cdot d\mathbf{r} + i \int \mathbf{F} \cdot d\mathbf{r} \times \hat{\mathbf{e}}_z \\ &= \boxed{\int \mathbf{F} \cdot d\mathbf{r}} + i \boxed{\int \mathbf{F} \cdot d\mathbf{S}} \quad \text{where: } d\mathbf{S} = d\mathbf{r} \times \hat{\mathbf{e}}_z \end{aligned}$$

**Real part**  $\int_1^2 \mathbf{F} \cdot d\mathbf{r} = \Delta\Phi$   
 sums  $\mathbf{F}$  projections *along* path  $d\mathbf{r}$  that is, *circulation* on path to get  $\Delta\Phi = \Phi(x_2, y_2) - \Phi(x_1, y_1)$




**Imaginary part**  $\int_1^2 \mathbf{F} \cdot d\mathbf{S} = \Delta\mathbf{A}$   
 sums  $\mathbf{F}$  projection *across* path  $d\mathbf{r}$  that is, *flux* thru surface elements  $d\mathbf{S} = d\mathbf{r} \times \mathbf{e}_z$  normal to  $d\mathbf{r}$  to get  $\Delta\mathbf{A} = \mathbf{A}(x_2, y_2) - \mathbf{A}(x_1, y_1)$



Here the scalar potential  $\Phi=(x^2-y^2)/2$  is stereo-plotted vs.  $(x,y)$

The  $\Phi=(x^2-y^2)/2=const.$  curves are topography lines

The  $A=(xy)=const.$  curves are streamlines normal to topography lines





## 4. Riemann-Cauchy conditions *What's analytic? (...and what's not?)*

*Easy 2D circulation and flux integrals*

 *Easy 2D curvilinear coordinate discovery*

*Easy 2D monopole, dipole, and  $2^n$ -pole analysis*

*Easy  $2^n$ -multipole field and potential expansion*

*Easy stereo-projection visualization*


10. Complex potentials define 2D Orthogonal Curvilinear Coordinates (OCC) of field

The  $(\Phi, A)$  grid is a GCC coordinate system\*:

$$q^1 = \Phi = (x^2 - y^2)/2 = \text{const.}$$

$$q^2 = A = (xy) = \text{const.}$$

\*Actually it's OCC.



$$Kajobian = \begin{pmatrix} \frac{\partial q^1}{\partial x} & \frac{\partial q^1}{\partial y} \\ \frac{\partial q^2}{\partial x} & \frac{\partial q^2}{\partial y} \end{pmatrix} = \begin{pmatrix} \frac{\partial \Phi}{\partial x} & \frac{\partial \Phi}{\partial y} \\ \frac{\partial A}{\partial x} & \frac{\partial A}{\partial y} \end{pmatrix} = \begin{pmatrix} x & -y \\ y & x \end{pmatrix} \leftarrow \begin{matrix} \mathbf{E}^\Phi \\ \mathbf{E}^A \end{matrix}$$

$$Jacobian = \begin{pmatrix} \frac{\partial x}{\partial q^1} & \frac{\partial x}{\partial q^2} \\ \frac{\partial y}{\partial q^1} & \frac{\partial y}{\partial q^2} \end{pmatrix} = \begin{pmatrix} \frac{\partial \Phi}{\partial \Phi} & \frac{\partial \Phi}{\partial A} \\ \frac{\partial A}{\partial \Phi} & \frac{\partial A}{\partial A} \end{pmatrix} = \frac{1}{r^2} \begin{pmatrix} x & y \\ -y & x \end{pmatrix}$$

$$Metric\ tensor = \begin{pmatrix} g_{\Phi\Phi} & g_{\Phi A} \\ g_{A\Phi} & g_{AA} \end{pmatrix} = \begin{pmatrix} \mathbf{E}_\Phi \cdot \mathbf{E}_\Phi & \mathbf{E}_\Phi \cdot \mathbf{E}_A \\ \mathbf{E}_A \cdot \mathbf{E}_\Phi & \mathbf{E}_A \cdot \mathbf{E}_A \end{pmatrix} = \begin{pmatrix} r^2 & 0 \\ 0 & r^2 \end{pmatrix} \text{ where: } r^2 = x^2 + y^2$$


### 10. Complex potentials define 2D Orthogonal Curvilinear Coordinates (OCC) of field

The  $(\Phi, A)$  grid is a GCC coordinate system\*:

$$q^1 = \Phi = (x^2 - y^2)/2 = \text{const.}$$

$$q^2 = A = (xy) = \text{const.}$$

\*Actually it's OCC.



$$Kajobian = \begin{pmatrix} \frac{\partial q^1}{\partial x} & \frac{\partial q^1}{\partial y} \\ \frac{\partial q^2}{\partial x} & \frac{\partial q^2}{\partial y} \end{pmatrix} = \begin{pmatrix} \frac{\partial \Phi}{\partial x} & \frac{\partial \Phi}{\partial y} \\ \frac{\partial A}{\partial x} & \frac{\partial A}{\partial y} \end{pmatrix} = \begin{pmatrix} x & -y \\ y & x \end{pmatrix} \leftarrow \begin{matrix} \mathbf{E}^\Phi \\ \mathbf{E}^A \end{matrix}$$

$$Jacobian = \begin{pmatrix} \frac{\partial x}{\partial q^1} & \frac{\partial x}{\partial q^2} \\ \frac{\partial y}{\partial q^1} & \frac{\partial y}{\partial q^2} \end{pmatrix} = \begin{pmatrix} \frac{\partial \Phi}{\partial \Phi} & \frac{\partial \Phi}{\partial A} \\ \frac{\partial A}{\partial \Phi} & \frac{\partial A}{\partial A} \end{pmatrix} = \frac{1}{r^2} \begin{pmatrix} x & y \\ -y & x \end{pmatrix}$$

$$Metric\ tensor = \begin{pmatrix} g_{\Phi\Phi} & g_{\Phi A} \\ g_{A\Phi} & g_{AA} \end{pmatrix} = \begin{pmatrix} \mathbf{E}_\Phi \cdot \mathbf{E}_\Phi & \mathbf{E}_\Phi \cdot \mathbf{E}_A \\ \mathbf{E}_A \cdot \mathbf{E}_\Phi & \mathbf{E}_A \cdot \mathbf{E}_A \end{pmatrix} = \begin{pmatrix} r^2 & 0 \\ 0 & r^2 \end{pmatrix} \text{ where: } r^2 = x^2 + y^2$$

Riemann-Cauchy Derivative Relations make coordinates orthogonal

$$\nabla \Phi = \begin{pmatrix} \frac{\partial \Phi}{\partial x} \\ \frac{\partial \Phi}{\partial y} \end{pmatrix} = \begin{pmatrix} \frac{\partial}{\partial x} \frac{a}{2} (x^2 - y^2) \\ \frac{\partial}{\partial y} \frac{a}{2} (x^2 - y^2) \end{pmatrix} = \begin{pmatrix} ax \\ -ay \end{pmatrix} = \mathbf{F}$$

The half-n'-half results assure

$$\mathbf{E}_\Phi \cdot \mathbf{E}_A = \frac{\partial \Phi}{\partial x} \frac{\partial A}{\partial x} + \frac{\partial \Phi}{\partial y} \frac{\partial A}{\partial y}$$

$$= -\frac{\partial \Phi}{\partial x} \frac{\partial \Phi}{\partial y} + \frac{\partial \Phi}{\partial y} \frac{\partial \Phi}{\partial x} = 0$$

$$\nabla \times \mathbf{A} = \begin{pmatrix} \frac{\partial A}{\partial y} \\ -\frac{\partial A}{\partial x} \end{pmatrix} = \begin{pmatrix} \frac{\partial}{\partial y} axy \\ -\frac{\partial}{\partial x} axy \end{pmatrix} = \begin{pmatrix} ax \\ -ay \end{pmatrix} = \mathbf{F}$$


10. Complex potentials define 2D Orthogonal Curvilinear Coordinates (OCC) of field

The  $(\Phi, A)$  grid is a GCC coordinate system\*:

$$q^1 = \Phi = (x^2 - y^2)/2 = \text{const.}$$

$$q^2 = A = (xy) = \text{const.}$$

\*Actually it's OCC.



$$Kajobian = \begin{pmatrix} \frac{\partial q^1}{\partial x} & \frac{\partial q^1}{\partial y} \\ \frac{\partial q^2}{\partial x} & \frac{\partial q^2}{\partial y} \end{pmatrix} = \begin{pmatrix} \frac{\partial \Phi}{\partial x} & \frac{\partial \Phi}{\partial y} \\ \frac{\partial A}{\partial x} & \frac{\partial A}{\partial y} \end{pmatrix} = \begin{pmatrix} x & -y \\ y & x \end{pmatrix} \leftarrow \begin{matrix} \mathbf{E}^\Phi \\ \mathbf{E}^A \end{matrix}$$

$$Jacobian = \begin{pmatrix} \frac{\partial x}{\partial q^1} & \frac{\partial x}{\partial q^2} \\ \frac{\partial y}{\partial q^1} & \frac{\partial y}{\partial q^2} \end{pmatrix} = \begin{pmatrix} \frac{\partial \Phi}{\partial \Phi} & \frac{\partial \Phi}{\partial A} \\ \frac{\partial A}{\partial \Phi} & \frac{\partial A}{\partial A} \end{pmatrix} = \frac{1}{r^2} \begin{pmatrix} x & y \\ -y & x \end{pmatrix}$$

$$Metric\ tensor = \begin{pmatrix} g_{\Phi\Phi} & g_{\Phi A} \\ g_{A\Phi} & g_{AA} \end{pmatrix} = \begin{pmatrix} \mathbf{E}_\Phi \cdot \mathbf{E}_\Phi & \mathbf{E}_\Phi \cdot \mathbf{E}_A \\ \mathbf{E}_A \cdot \mathbf{E}_\Phi & \mathbf{E}_A \cdot \mathbf{E}_A \end{pmatrix} = \begin{pmatrix} r^2 & 0 \\ 0 & r^2 \end{pmatrix} \text{ where: } r^2 = x^2 + y^2$$

Riemann-Cauchy Derivative Relations make coordinates orthogonal

$$\nabla \Phi = \begin{pmatrix} \frac{\partial \Phi}{\partial x} \\ \frac{\partial \Phi}{\partial y} \end{pmatrix} = \begin{pmatrix} \frac{\partial}{\partial x} \frac{a}{2} (x^2 - y^2) \\ \frac{\partial}{\partial y} \frac{a}{2} (x^2 - y^2) \end{pmatrix} = \begin{pmatrix} ax \\ -ay \end{pmatrix} = \mathbf{F}$$

The half-n'-half results assure

$$\mathbf{E}_\Phi \cdot \mathbf{E}_A = \frac{\partial \Phi}{\partial x} \frac{\partial A}{\partial x} + \frac{\partial \Phi}{\partial y} \frac{\partial A}{\partial y} = -\frac{\partial \Phi}{\partial x} \frac{\partial \Phi}{\partial y} + \frac{\partial \Phi}{\partial y} \frac{\partial \Phi}{\partial x} = 0$$

$$\nabla \times \mathbf{A} = \begin{pmatrix} \frac{\partial A}{\partial y} \\ -\frac{\partial A}{\partial x} \end{pmatrix} = \begin{pmatrix} \frac{\partial}{\partial y} axy \\ -\frac{\partial}{\partial x} axy \end{pmatrix} = \begin{pmatrix} ax \\ -ay \end{pmatrix} = \mathbf{F}$$

or Riemann-Cauchy

Zero divergence requirement:  $0 = \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} = \frac{\partial}{\partial x} \frac{\partial \Phi}{\partial x} + \frac{\partial}{\partial y} \frac{\partial \Phi}{\partial y} = \frac{\partial^2 \Phi}{\partial x^2} + \frac{\partial^2 \Phi}{\partial y^2} = 0$

and so does A

potential  $\Phi$  obeys Laplace equation



## *4. Riemann-Cauchy conditions What's analytic? (...and what's not?)*

*Easy 2D circulation and flux integrals*

*Easy 2D curvilinear coordinate discovery*

 *Easy 2D monopole, dipole, and  $2^n$ -pole analysis*

*Easy  $2^n$ -multipole field and potential expansion*

*Easy stereo-projection visualization*

## What Good Are Complex Exponentials? (contd.)

### 11. Complex integrals define 2D *monopole* fields and potentials

Of all power-law fields  $f(z)=az^n$  one lacks a power-law potential  $\phi(z)=\frac{a}{n+1}z^{n+1}$ . It is the  $n = -1$  case.

Unit *monopole* field:  $f(z)=\frac{1}{z}=z^{-1}$

$f(z)=\frac{a}{z}=az^{-1}$  Source- $a$  *monopole*

It has a *logarithmic potential*  $\phi(z)=a\cdot\ln(z)=a\cdot\ln(x+iy)$ .

## What Good Are Complex Exponentials? (contd.)

### 11. Complex integrals define 2D *monopole* fields and potentials

Of all power-law fields  $f(z)=az^n$  one lacks a power-law potential  $\phi(z)=\frac{a}{n+1}z^{n+1}$ . It is the  $n = -1$  case.

Unit *monopole* field:  $f(z)=\frac{1}{z}=z^{-1}$        $f(z)=\frac{a}{z}=az^{-1}$  Source- $a$  *monopole*

It has a *logarithmic potential*  $\phi(z)=a\cdot\ln(z)=a\cdot\ln(x+iy)$ .

$$\phi(z) = \Phi + i\mathbf{A} = \int f(z)dz = \int \frac{a}{z} dz = a \ln(z)$$

## What Good Are Complex Exponentials? (contd.)

### 11. Complex integrals define 2D *monopole* fields and potentials

Of all power-law fields  $f(z)=az^n$  one lacks a power-law potential  $\phi(z)=\frac{a}{n+1}z^{n+1}$ . It is the  $n = -1$  case.

$$\text{Unit monopole field: } f(z)=\frac{1}{z}=z^{-1} \qquad f(z)=\frac{a}{z}=az^{-1} \text{ Source-}a \text{ monopole}$$

It has a *logarithmic potential*  $\phi(z)=a \cdot \ln(z)=a \cdot \ln(x+iy)$ . Note:  $\ln(a \cdot b)=\ln(a)+\ln(b)$ ,  $\ln(e^{i\theta})=i\theta$ , and  $z=re^{i\theta}$ .

$$\begin{aligned} \phi(z) &= \underbrace{\Phi}_{a \ln(r)} + \underbrace{i\mathbf{A}}_{i a \theta} = \int f(z) dz = \int \frac{a}{z} dz = a \ln(z) = a \ln(re^{i\theta}) \\ &= a \ln(r) + i a \theta \end{aligned}$$



## What Good Are Complex Exponentials? (contd.)

### 11. Complex integrals define 2D *monopole* fields and potentials


Of all power-law fields  $f(z)=az^n$  one lacks a power-law potential  $\phi(z)=\frac{a}{n+1}z^{n+1}$ . It is the  $n = -1$  case.

Unit *monopole* field:  $f(z)=\frac{1}{z}=z^{-1}$        $f(z)=\frac{a}{z}=az^{-1}$  Source- $a$  *monopole*

It has a *logarithmic potential*  $\phi(z)=a\cdot\ln(z)=a\cdot\ln(x+iy)$ . Note:  $\ln(a\cdot b)=\ln(a)+\ln(b)$ ,  $\ln(e^{i\theta})=i\theta$ , and  $z=re^{i\theta}$ .

$$\begin{aligned}\phi(z) &= \Phi + i\mathbf{A} = \int f(z)dz = \int \frac{a}{z} dz = a \ln(z) = a \ln(re^{i\theta}) \\ &= \underbrace{a \ln(r)} + i \underbrace{a\theta}\end{aligned}$$

(a) Unit Z-line-flux field  $f(z)=1/z$



## What Good Are Complex Exponentials? (contd.)

### 11. Complex integrals define 2D *monopole* fields and potentials

Of all power-law fields  $f(z)=az^n$  one lacks a power-law potential  $\phi(z)=\frac{a}{n+1}z^{n+1}$ . It is the  $n = -1$  case.


Unit *monopole* field:  $f(z)=\frac{1}{z}=z^{-1}$        $f(z)=\frac{a}{z}=az^{-1}$  Source- $a$  *monopole*

It has a *logarithmic potential*  $\phi(z)=a\cdot\ln(z)=a\cdot\ln(x+iy)$ . Note:  $\ln(a\cdot b)=\ln(a)+\ln(b)$ ,  $\ln(e^{i\theta})=i\theta$ , and  $z=re^{i\theta}$ .


$$\begin{aligned} \phi(z) &= \Phi + i\mathbf{A} = \int f(z)dz = \int \frac{a}{z} dz = a \ln(z) = a \ln(re^{i\theta}) \\ &= \underbrace{a \ln(r)} + i \underbrace{a\theta} \end{aligned}$$

(a) Unit Z-line-flux field  $f(z)=1/z$

(b) Unit Z-line-vortex field  $f(z)=i/z$



Field:  
 $f^*(z^*)=1/z^*=e^{i\theta}/r$   
 $\mathbf{F}_{(x,y)}=(x,y)/r^2$   
Potential:  
 $\phi(z)=\ln z$   
 $=\ln r+i\theta$   
 $=\Phi+i\mathbf{A}$



Field:  
 $f^*(z^*)=-i/z^*=-ie^{i\theta}/r$   
 $\mathbf{F}_{(x,y)}=(y,-x)/r^2$   
Potential:  
 $\phi(z)=i \ln z$   
 $=\theta+i \ln r$   
 $=\Phi+i\mathbf{A}$

## What Good Are Complex Exponentials? (contd.)

### 11. Complex integrals define 2D *monopole* fields and potentials



Of all power-law fields  $f(z)=az^n$  one lacks a power-law potential  $\phi(z)=\frac{a}{n+1}z^{n+1}$ . It is the  $n = -1$  case.

Unit *monopole* field:  $f(z)=\frac{1}{z}=z^{-1}$        $f(z)=\frac{a}{z}=az^{-1}$  Source- $a$  *monopole*

It has a *logarithmic potential*  $\phi(z)=a\cdot\ln(z)=a\cdot\ln(x+iy)$ . Note:  $\ln(a\cdot b)=\ln(a)+\ln(b)$ ,  $\ln(e^{i\theta})=i\theta$ , and  $z=re^{i\theta}$ .

$$\begin{aligned} \phi(z) &= \underbrace{\Phi}_{= a \ln(r)} + \underbrace{i\mathbf{A}}_{+ i a \theta} = \int f(z)dz = \int \frac{a}{z} dz = a \ln(z) = a \ln(re^{i\theta}) \\ &= a \ln(r) + i a \theta \end{aligned}$$


A *monopole* field is the only power-law field whose integral (potential) depends on *path of integration*.



*path that goes N times around origin ( $r=0$ ) at constant  $r = R$ .*


$$\Delta\phi = \oint f(z)dz = a \oint \frac{dz}{z} = a \int_{\theta=0}^{\theta=2\pi N} \frac{d(Re^{i\theta})}{Re^{i\theta}} = a \int_{\theta=0}^{\theta=2\pi N} id\theta = ai \theta \Big|_0^{2\pi N} = 2a\pi iN$$

(a) Unit Z-line-flux field  $f(z)=1/z$




$$\phi(z) = \underbrace{\Phi}_{\ln(r)} + \underbrace{iA}_{i\theta} = \int f(z)dz = \int \frac{a}{z} dz = a \ln(re^{i\theta})$$

*(For a=1)*




*Each turn around origin adds  $2\pi i$  to vector potential  $iA$*








*(For a=1)*



(a) Unit Z-line-flux field  $f(z)=1/z$




(b) Unit Z-line-vortex field  $f(z)=i/z$



# What Good Are Complex Exponentials? (contd.)


$$f(z) = (0.5 + i0.5)/z = e^{i\pi/4}/z\sqrt{2}$$

“Vortex”



$$f(z) = (0.75 + i0.25)/z = e^{i18^\circ}/z\sqrt{n}$$


“Hurricane”





# What Good Are Complex Exponentials? (contd.)

An assist from *Physics Girl* (YouTube Channel):



Posted this year:

[How to Make VORTEX RINGS in a Pool](#)




Crazy pool vortex (new inclusion with more background)

[Crazy pool vortex - pg-yt-2014](#)

Posting with the best visuals:

[Fun with Vortex Rings in the Pool - pg-yt-2014](#)

*She covers it beautifully!*



## *4. Riemann-Cauchy conditions What's analytic? (...and what's not?)*

*Easy 2D circulation and flux integrals*

*Easy 2D curvilinear coordinate discovery*

 *Easy 2D monopole, dipole, and  $2^n$ -pole analysis*

*Easy  $2^n$ -multipole field and potential expansion*

*Easy stereo-projection visualization*



# What Good Are Complex Exponentials? (2D monopole, dipole, and $2^n$ -pole analysis)

## 12. Complex derivatives give 2D dipole fields

Start with  $f(z)=az^{-1}$ : 2D line *monopole field* and is its *monopole potential*  $\phi(z)=a \ln z$  of source strength  $a$ .

$$f^{1-pole}(z) = \frac{a}{z} = \frac{d\phi^{1-pole}}{dz} \quad \phi^{1-pole}(z) = a \ln z$$

Now let these two line-sources of equal but opposite source constants  $+a$  and  $-a$  be located at  $z=\pm\Delta/2$  separated by a small interval  $\Delta$ . This sum (actually difference) of  $f^{1-pole}$ -fields is called a *dipole field*.

$$f^{dipole}(z) = \frac{a}{z+\frac{\Delta}{2}} - \frac{a}{z-\frac{\Delta}{2}} = \frac{-a \cdot \Delta}{z^2 - \frac{\Delta^2}{4}} \quad \phi^{dipole}(z) = a \ln(z - \frac{\Delta}{2}) - a \ln(z + \frac{\Delta}{2}) = a \ln \frac{z - \frac{\Delta}{2}}{z + \frac{\Delta}{2}}$$

*This is like the derivative definition:*


$$\frac{df}{dz} = \frac{f(z+\Delta) - f(z)}{\Delta}$$

*or:*

$$\frac{df}{dz} = \frac{f(z+\frac{\Delta}{2}) - f(z-\frac{\Delta}{2})}{\Delta}$$

*if  $\Delta$  is infinitesimal*

$$(\Delta \rightarrow 0)$$



*So-called “physical dipole” has finite  $\Delta$  (+)(-) separation*

## What Good Are Complex Exponentials? (2D monopole, dipole, and 2<sup>n</sup>-pole analysis)

### 12. Complex derivatives give 2D dipole fields

Start with  $f(z)=az^{-1}$ : 2D line *monopole field* and is its *monopole potential*  $\phi(z)=a \ln z$  of source strength  $a$ .

$$f^{1-pole}(z) = \frac{a}{z} = \frac{d\phi^{1-pole}}{dz} \quad \phi^{1-pole}(z) = a \ln z$$

Now let these two line-sources of equal but opposite source constants  $+a$  and  $-a$  be located at  $z=\pm\Delta/2$  separated by a small interval  $\Delta$ . This sum (actually difference) of  $f^{1-pole}$ -fields is called a *dipole field*.

$$f^{dipole}(z) = \frac{a}{z+\frac{\Delta}{2}} - \frac{a}{z-\frac{\Delta}{2}} = \frac{-a \cdot \Delta}{z^2 - \frac{\Delta^2}{4}} \quad \phi^{dipole}(z) = a \ln\left(z - \frac{\Delta}{2}\right) - a \ln\left(z + \frac{\Delta}{2}\right) = a \ln \frac{z - \frac{\Delta}{2}}{z + \frac{\Delta}{2}}$$

If interval  $\Delta$  is *tiny* and is divided out we get a *point-dipole field*  $f^{2-pole}$  that is the  $z$ -derivative of  $f^{1-pole}$ .

$$f^{2-pole} = \frac{-a}{z^2} = \frac{df^{1-pole}}{dz} = \frac{d\phi^{2-pole}}{dz} \quad \phi^{2-pole} = \frac{a}{z} = \frac{d\phi^{1-pole}}{dz}$$

## What Good Are Complex Exponentials? (2D monopole, dipole, and $2^n$ -pole analysis)

### 12. Complex derivatives give 2D dipole fields

Start with  $f(z)=az^{-1}$ : 2D line *monopole field* and is its *monopole potential*  $\phi(z)=a \ln z$  of source strength  $a$ .


$$f^{1-pole}(z) = \frac{a}{z} = \frac{d\phi^{1-pole}}{dz} \quad \phi^{1-pole}(z) = a \ln z$$

Now let these two line-sources of equal but opposite source constants  $+a$  and  $-a$  be located at  $z=\pm\Delta/2$  separated by a small interval  $\Delta$ . This sum (actually difference) of  $f^{1-pole}$ -fields is called a *dipole field*.

$$f^{dipole}(z) = \frac{a}{z + \frac{\Delta}{2}} - \frac{a}{z - \frac{\Delta}{2}} = \frac{-a \cdot \Delta}{z^2 - \frac{\Delta^2}{4}} \quad \phi^{dipole}(z) = a \ln\left(z - \frac{\Delta}{2}\right) - a \ln\left(z + \frac{\Delta}{2}\right) = a \ln \frac{z - \frac{\Delta}{2}}{z + \frac{\Delta}{2}}$$

If interval  $\Delta$  is *tiny* and is divided out we get a *point-dipole field*  $f^{2-pole}$  that is the  $z$ -derivative of  $f^{1-pole}$ .

$$f^{2-pole} = \frac{-a}{z^2} = \frac{df^{1-pole}}{dz} = \frac{d\phi^{2-pole}}{dz} \quad \phi^{2-pole} = \frac{a}{z} = \frac{d\phi^{1-pole}}{dz}$$



## What Good Are Complex Exponentials? (2D monopole, dipole, and 2<sup>n</sup>-pole analysis)

### 12. Complex derivatives give 2D dipole fields

Start with  $f(z)=az^{-1}$ : 2D line *monopole field* and is its *monopole potential*  $\phi(z)=a \ln z$  of source strength  $a$ .

$$f^{1-pole}(z) = \frac{a}{z} = \frac{d\phi^{1-pole}}{dz} \quad \phi^{1-pole}(z) = a \ln z$$

Now let these two line-sources of equal but opposite source constants  $+a$  and  $-a$  be located at  $z=\pm\Delta/2$  separated by a small interval  $\Delta$ . This sum (actually difference) of  $f^{1-pole}$ -fields is called a *dipole field*.

$$f^{dipole}(z) = \frac{a}{z+\frac{\Delta}{2}} - \frac{a}{z-\frac{\Delta}{2}} = \frac{-a \cdot \Delta}{z^2 - \frac{\Delta^2}{4}} \quad \phi^{dipole}(z) = a \ln\left(z - \frac{\Delta}{2}\right) - a \ln\left(z + \frac{\Delta}{2}\right) = a \ln \frac{z - \frac{\Delta}{2}}{z + \frac{\Delta}{2}}$$

If interval  $\Delta$  is *tiny* and is divided out we get a *point-dipole field*  $f^{2-pole}$  that is the  $z$ -derivative of  $f^{1-pole}$ .


$$f^{2-pole} = \frac{-a}{z^2} = \frac{df^{1-pole}}{dz} = \frac{d\phi^{2-pole}}{dz} \quad \phi^{2-pole} = \frac{a}{z} = \frac{d\phi^{1-pole}}{dz}$$

A *point-dipole potential*  $\phi^{2-pole}$  (whose  $z$ -derivative is  $f^{2-pole}$ ) is a  $z$ -derivative of  $\phi^{1-pole}$ .

$$\begin{aligned} \phi^{2-pole} &= \frac{a}{z} = \frac{a}{x+iy} = \frac{a}{x+iy} \frac{x-iy}{x-iy} = \frac{ax}{x^2+y^2} + i \frac{-ay}{x^2+y^2} = \frac{a}{r} \cos \theta - i \frac{a}{r} \sin \theta \\ &= \Phi^{2-pole} + i \mathbf{A}^{2-pole} \end{aligned}$$

A *point-dipole potential*  $\phi^{2-pole}$  (whose  $z$ -derivative is  $f^{2-pole}$ ) is a  $z$ -derivative of  $\phi^{1-pole}$ .


$$\begin{aligned} \phi^{2-pole} &= \frac{a}{z} = \frac{a}{x+iy} = \frac{a}{x+iy} \frac{x-iy}{x-iy} = \frac{ax}{x^2+y^2} + i \frac{-ay}{x^2+y^2} = \frac{a}{r} \cos \theta - i \frac{a}{r} \sin \theta \\ &= \Phi^{2-pole} + i A^{2-pole} \end{aligned}$$



Field:  
 $f^*(z^*) = 1/z^{2*} = e^{i2\theta}/r^2$   
 $\mathbf{F}(x,y) = (\cos 2\theta, \sin 2\theta)/r^2$   
Potential:  
 $\phi(z) = 1/z$   
 $= (\cos \theta)/r + i(\sin \theta)/r$   
 $= \Phi + i A$

Scalar potentials

$$\Phi = (a/r) \cos \theta = const.$$



Vector potentials

$$A = (a/r) \sin \theta = const.$$



## $2^n$ -pole analysis (quadrupole: $2^2=4$ -pole, octapole: $2^3=8$ -pole, ..., pole dancer,

What if we put a (-)copy of a 2-pole near its original?

Well, the result is 4-pole or *quadrupole* field  $f^{4-pole}$  and potential  $\phi^{4-pole}$ .

Each a z-derivative of  $f^{2-pole}$  and  $\phi^{2-pole}$ .

$$f^{4-pole} = \frac{a}{z^3} = \frac{1}{2} \frac{df^{2-pole}}{dz} = \frac{d\phi^{4-pole}}{dz}$$

$$\phi^{4-pole} = -\frac{a}{2z^2} = \frac{1}{2} \frac{d\phi^{2-pole}}{dz}$$

# $2^n$ -pole analysis (quadrupole: $2^2=4$ -pole, octapole: $2^3=8$ -pole, ..., pole dancer,



What if we put a (-)copy of a 2-pole near its original?

Well, the result is 4-pole or *quadrupole* field  $f^{4-pole}$  and potential  $\phi^{4-pole}$ .

Each a z-derivative of  $f^{2-pole}$  and  $\phi^{2-pole}$ , respectively.

$$f^{4-pole} = \frac{a}{z^3} = \frac{1}{2} \frac{df^{2-pole}}{dz} = \frac{d\phi^{4-pole}}{dz}$$

$$\phi^{4-pole} = -\frac{a}{2z^2} = \frac{1}{2} \frac{d\phi^{2-pole}}{dz}$$



Field:  
 $f^*(z^*) = 1/z^3 = e^{i3\theta}/r^3$   
 $\mathbf{F}(x,y) = (\cos 3\theta, \sin 3\theta)/r^3$   
Potential:  
 $-2\phi(z) = 1/z^2$   
 $= (\cos 2\theta)/r^2 + i(\sin 2\theta)/r^2$   
 $= \Phi + iA$

## *4. Riemann-Cauchy conditions What's analytic? (...and what's not?)*

*Easy 2D circulation and flux integrals*

*Easy 2D curvilinear coordinate discovery*

*Easy 2D monopole, dipole, and  $2^n$ -pole analysis*

*Easy  $2^n$ -multipole field and potential expansion*

*Easy stereo-projection visualization*



## $2^n$ -pole analysis: Laurent series (Generalization of Maclaurin-Taylor series)

*Laurent series* or *multipole expansion* of a given complex field function  $f(z)$  around  $z=0$ .

$$\begin{aligned} \frac{d\phi}{dz} = f(z) &= \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots \\ &\quad \dots \begin{array}{l} 2^2\text{-pole} \\ \text{(quadrupole)} \\ \text{at } z=0 \end{array} + \begin{array}{l} 2^1\text{-pole} \\ \text{(dipole)} \\ \text{at } z=0 \end{array} + \begin{array}{l} 2^0\text{-pole} \\ \text{(monopole)} \\ \text{at } z=0 \end{array} + \begin{array}{l} 2^1\text{-pole} \\ \text{(dipole)} \\ \text{at } z=\infty \end{array} + \begin{array}{l} 2^2\text{-pole} \\ \text{(quadrupole)} \\ \text{at } z=\infty \end{array} + \begin{array}{l} 2^3\text{-pole} \\ \text{(octapole)} \\ \text{at } z=\infty \end{array} + \begin{array}{l} 2^4\text{-pole} \\ \text{(hexadecapole)} \\ \text{at } z=\infty \end{array} + \begin{array}{l} 2^5\text{-pole} \\ \text{at } z=\infty \end{array} + \begin{array}{l} 2^6\text{-pole} \\ \text{at } z=\infty \end{array} \dots \\ \int f dz = \phi(z) &= \dots \frac{a_{-3}}{-2}z^{-2} + \frac{a_{-2}}{-1}z^{-1} + a_{-1} \ln z + a_0z + \frac{a_1}{2}z^2 + \frac{a_2}{3}z^3 + \frac{a_3}{4}z^4 + \frac{a_4}{5}z^5 + \frac{a_5}{6}z^6 + \dots \end{aligned}$$

All field terms  $a_{m-1}z^{m-1}$  except  $1\text{-pole } \frac{a_{-1}}{z}$  have potential term  $a_{m-1}z^m/m$  of a  $2^m$ -pole.

These are located at  $z=0$  for  $m < 0$  and at  $z=\infty$  for  $m > 0$ .

$$\phi(z) = \dots \begin{array}{l} \text{(octapole)}_0 \\ \frac{a_{-4}}{-3}z^{-3} \end{array} + \begin{array}{l} \text{(quadrupole)}_0 \\ \frac{a_{-3}}{-2}z^{-2} \end{array} + \begin{array}{l} \text{(dipole)}_0 \\ \frac{a_{-2}}{-1}z^{-1} \end{array} + \begin{array}{l} \text{(monopole)} \\ a_{-1} \ln z \end{array} + \begin{array}{l} \text{(dipole)}_\infty \\ a_0z \end{array} + \begin{array}{l} \text{(quadrupole)}_\infty \\ \frac{a_1}{2}z^2 \end{array} + \begin{array}{l} \text{(octapole)}_\infty \\ \frac{a_2}{3}z^3 \end{array} + \dots$$

## $2^n$ -pole analysis: Laurent series (Generalization of Maclaurin-Taylor series)

*Laurent series* or *multipole expansion* of a given complex field function  $f(z)$  around  $z=0$ .

$$\begin{aligned} \frac{d\phi}{dz} = f(z) &= \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots \\ &\quad \dots \begin{array}{l} 2^2\text{-pole} \\ \text{(quadrupole)} \\ \text{at } z=0 \end{array} + \begin{array}{l} 2^1\text{-pole} \\ \text{(dipole)} \\ \text{at } z=0 \end{array} + \begin{array}{l} 2^0\text{-pole} \\ \text{(monopole)} \\ \text{at } z=0 \end{array} + \begin{array}{l} 2^1\text{-pole} \\ \text{(dipole)} \\ \text{at } z=\infty \end{array} + \begin{array}{l} 2^2\text{-pole} \\ \text{(quadrupole)} \\ \text{at } z=\infty \end{array} + \begin{array}{l} 2^3\text{-pole} \\ \text{(octapole)} \\ \text{at } z=\infty \end{array} + \begin{array}{l} 2^4\text{-pole} \\ \text{(hexadecapole)} \\ \text{at } z=\infty \end{array} + \begin{array}{l} 2^5\text{-pole} \\ \text{at } z=\infty \end{array} + \begin{array}{l} 2^6\text{-pole} \\ \text{at } z=\infty \end{array} \dots \\ \int f dz = \phi(z) &= \dots \frac{a_{-3}}{-2}z^{-2} + \frac{a_{-2}}{-1}z^{-1} + a_{-1} \ln z + a_0z + \frac{a_1}{2}z^2 + \frac{a_2}{3}z^3 + \frac{a_3}{4}z^4 + \frac{a_4}{5}z^5 + \frac{a_5}{6}z^6 + \dots \end{aligned}$$

All field terms  $a_{m-1}z^{m-1}$  except  $1\text{-pole } \frac{a_{-1}}{z}$  have potential term  $a_{m-1}z^m/m$  of a  $2^m$ -pole.

These are located at  $z=0$  for  $m < 0$  and at  $z=\infty$  for  $m > 0$ .

$$\phi(z) = \dots \begin{array}{l} \text{(octapole)}_0 \\ \frac{a_{-4}}{-3}z^{-3} \end{array} + \begin{array}{l} \text{(quadrupole)}_0 \\ \frac{a_{-3}}{-2}z^{-2} \end{array} + \begin{array}{l} \text{(dipole)}_0 \\ \frac{a_{-2}}{-1}z^{-1} \end{array} + \begin{array}{l} \text{(monopole)} \\ a_{-1} \ln z \end{array} + \begin{array}{l} \text{(dipole)}_\infty \\ a_0z \end{array} + \begin{array}{l} \text{(quadrupole)}_\infty \\ \frac{a_1}{2}z^2 \end{array} + \begin{array}{l} \text{(octapole)}_\infty \\ \frac{a_2}{3}z^3 \end{array} + \dots$$

$$\phi(w) = \dots \frac{a_{-4}}{-3}w^{-3} + \frac{a_{-3}}{-2}w^{-2} + \frac{a_{-2}}{-1}w^{-1} + a_{-1} \ln w + a_0w + \frac{a_1}{2}w^2 + \frac{a_2}{3}w^3 + \dots$$

(with  $z=w^{-1}$ )



# $2^n$ -pole analysis: Laurent series (Generalization of Maclaurin-Taylor series)

*Laurent series* or *multipole expansion* of a given complex field function  $f(z)$  around  $z=0$ .

$$\frac{d\phi}{dz} = f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

	...	2 <sup>2</sup> -pole	2 <sup>1</sup> -pole	2 <sup>0</sup> -pole	2 <sup>1</sup> -pole	2 <sup>2</sup> -pole	2 <sup>3</sup> -pole	2 <sup>4</sup> -pole	2 <sup>5</sup> -pole	2 <sup>6</sup> -pole	...
		(quadrupole)	(dipole)	(monopole)	(dipole)	(quadrupole)	(octapole)	(hexadecapole)			
		at $z=0$	at $z=0$	at $z=0$	at $z=\infty$	at $z=\infty$	at $z=\infty$	at $z=\infty$	at $z=\infty$	at $z=\infty$	

$$\int f dz = \phi(z) = \dots \frac{a_{-3}}{-2} z^{-2} + \frac{a_{-2}}{-1} z^{-1} + a_{-1} \ln z + a_0 z + \frac{a_1}{2} z^2 + \frac{a_2}{3} z^3 + \frac{a_3}{4} z^4 + \frac{a_4}{5} z^5 + \frac{a_5}{6} z^6 + \dots$$

All field terms  $a_{m-1}z^{m-1}$  except *1-pole*  $\frac{a_{-1}}{z}$  have potential term  $a_{m-1}z^m/m$  of a  $2^m$ -pole.

These are located at  $z=0$  for  $m < 0$  and at  $z=\infty$  for  $m > 0$ .

$$\phi(z) = \dots \frac{a_{-4}}{-3} z^{-3} + \frac{a_{-3}}{-2} z^{-2} + \frac{a_{-2}}{-1} z^{-1} + a_{-1} \ln z + a_0 z + \frac{a_1}{2} z^2 + \frac{a_2}{3} z^3 + \dots$$


(octapole)<sub>0</sub>
(quadrupole)<sub>0</sub>
(dipole)<sub>0</sub>
(monopole)
(dipole)<sub>∞</sub>
(quadrupole)<sub>∞</sub>
(octapole)<sub>∞</sub>

$$\phi(w) = \dots \frac{a_{-4}}{-3} w^{-3} + \frac{a_{-3}}{-2} w^{-2} + \frac{a_{-2}}{-1} w^{-1} + a_{-1} \ln w + a_0 w + \frac{a_1}{2} w^2 + \frac{a_2}{3} w^3 + \dots$$

(with  $z \rightarrow w$ )

$$= \dots \frac{a_2}{3} z^{-3} + \frac{a_1}{2} z^{-2} + a_0 z^{-1} - a_{-1} \ln z + \frac{a_{-2}}{-1} z + \frac{a_{-3}}{-2} z^2 + \frac{a_{-4}}{-3} z^3 + \dots$$



(with  $w = z^{-1}$ )



$$\begin{aligned} \phi(z) &= \dots \frac{a_{-4}}{-3} z^{-3} + \frac{a_{-3}}{-2} z^{-2} + \frac{a_{-2}}{-1} z^{-1} + a_{-1} \ln z + a_0 z + \frac{a_1}{2} z^2 + \frac{a_2}{3} z^3 + \dots \\ &\quad \text{(octapole)}_0 \quad \text{(quadrupole)}_0 \quad \text{(dipole)}_0 \quad \text{(monopole)} \quad \text{(dipole)}_\infty \quad \text{(quadrupole)}_\infty \quad \text{(octapole)}_\infty \\ \phi(w) &= \dots \frac{a_{-4}}{-3} w^{-3} + \frac{a_{-3}}{-2} w^{-2} + \frac{a_{-2}}{-1} w^{-1} + a_{-1} \ln w + a_0 w + \frac{a_1}{2} w^2 + \frac{a_2}{3} w^3 + \dots \\ &= \dots \frac{a_2}{3} z^{-2} + \frac{a_1}{2} z^{-2} + a_0 z^{-1} - a_{-1} \ln z + \frac{a_{-2}}{-1} z + \frac{a_{-3}}{-2} z^2 + \frac{a_{-4}}{-3} z^3 + \dots \end{aligned}$$

(with  $z \rightarrow w$ )

(with  $w = z^{-1}$ )



quadrupole field centered at North Pole

is quadratic field near South Pole

$\phi(z) = \frac{a_{-3}}{-2} z^{-2}$   
 $f(z) = a_{-3} z^{-3}$

$\phi(w) = a_0 w^2$   
 $f(w) = a_1 w$

$$f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$

$$f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$

This  $m=1$ -pole constant- $a_{-1}$  formula is just the first in a series of *Laurent coefficient expressions*.

$$\dots a_{-3} = \frac{1}{2\pi i} \oint z^2 f(z)dz, \quad a_{-2} = \frac{1}{2\pi i} \oint z^1 f(z)dz, \quad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz, \quad a_0 = \frac{1}{2\pi i} \oint \frac{f(z)}{z} dz, \quad a_1 = \frac{1}{2\pi i} \oint \frac{f(z)}{z^2} dz, \dots$$



$$f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$

This  $m=1$ -pole constant- $a_{-1}$  formula is just the first in a series of *Laurent coefficient expressions*.

$$\dots a_{-3} = \frac{1}{2\pi i} \oint z^2 f(z)dz, \quad a_{-2} = \frac{1}{2\pi i} \oint z^1 f(z)dz, \quad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz, \quad a_0 = \frac{1}{2\pi i} \oint \frac{f(z)}{z} dz, \quad a_1 = \frac{1}{2\pi i} \oint \frac{f(z)}{z^2} dz, \dots$$

Source analysis starts with 1-pole loop integrals  $\oint z^{-1}dz = 2\pi i$  or, with origin shifted  $\oint (z-a)^{-1}dz = 2\pi i$ .

$$f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$

This  $m=1$ -pole constant- $a_{-1}$  formula is just the first in a series of *Laurent coefficient expressions*.

$$\dots a_{-3} = \frac{1}{2\pi i} \oint z^2 f(z)dz, \quad a_{-2} = \frac{1}{2\pi i} \oint z^1 f(z)dz, \quad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz, \quad a_0 = \frac{1}{2\pi i} \oint \frac{f(z)}{z} dz, \quad a_1 = \frac{1}{2\pi i} \oint \frac{f(z)}{z^2} dz, \dots$$

Source analysis starts with 1-pole loop integrals  $\oint z^{-1}dz = 2\pi i$  or, with origin shifted  $\oint (z-a)^{-1}dz = 2\pi i$ .

They hold for any loop about point- $a$ . Function  $f(z)$  is just  $f(a)$  on a *tiny* circle around point- $a$ .

(assume *tiny* circle around  $z=a$ )

$$\oint \frac{f(z)}{z-a} dz = \oint \frac{f(a)}{z-a} dz = f(a) \oint \frac{1}{z-a} dz = 2\pi i f(a)$$

(but any contour that doesn't "touch  $a$  gives same answer)

$$f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$

This  $m=1$ -pole constant- $a_{-1}$  formula is just the first in a series of *Laurent coefficient expressions*.

$$\dots a_{-3} = \frac{1}{2\pi i} \oint z^2 f(z)dz, \quad a_{-2} = \frac{1}{2\pi i} \oint z^1 f(z)dz, \quad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz, \quad a_0 = \frac{1}{2\pi i} \oint \frac{f(z)}{z} dz, \quad a_1 = \frac{1}{2\pi i} \oint \frac{f(z)}{z^2} dz, \dots$$

Source analysis starts with 1-pole loop integrals  $\oint z^{-1}dz = 2\pi i$  or, with origin shifted  $\oint (z-a)^{-1}dz = 2\pi i$ .

They hold for any loop about point- $a$ . Function  $f(z)$  is just  $f(a)$  on a *tiny* circle around point- $a$ .

(assume *tiny* circle around  $z=a$ )

$$\oint \frac{f(z)}{z-a} dz = \oint \frac{f(a)}{z-a} dz = f(a) \oint \frac{1}{z-a} dz = 2\pi i f(a)$$

(but any contour that doesn't "touch  $a$  gives same answer)

$$f(a) = \frac{1}{2\pi i} \oint \frac{f(z)}{z-a} dz$$

The  $f(a)$  result is called a *Cauchy integral*.

$$f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$

This  $m=1$ -pole constant- $a_{-1}$  formula is just the first in a series of *Laurent coefficient expressions*.

$$\dots a_{-3} = \frac{1}{2\pi i} \oint z^2 f(z)dz, \quad a_{-2} = \frac{1}{2\pi i} \oint z^1 f(z)dz, \quad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz, \quad a_0 = \frac{1}{2\pi i} \oint \frac{f(z)}{z} dz, \quad a_1 = \frac{1}{2\pi i} \oint \frac{f(z)}{z^2} dz, \dots$$

Source analysis starts with 1-pole loop integrals  $\oint z^{-1}dz = 2\pi i$  or, with origin shifted  $\oint (z-a)^{-1}dz = 2\pi i$ .

They hold for any loop about point- $a$ . Function  $f(z)$  is just  $f(a)$  on a *tiny* circle around point- $a$ .

(assume *tiny* circle around  $z=a$ )

$$\oint \frac{f(z)}{z-a} dz = \oint \frac{f(a)}{z-a} dz = f(a) \oint \frac{1}{z-a} dz = 2\pi i f(a)$$

(but any contour that doesn't "touch  $a$  gives same answer)

$$f(a) = \frac{1}{2\pi i} \oint \frac{f(z)}{z-a} dz$$

The  $f(a)$  result is called a *Cauchy integral*. Then repeated  $a$ -derivatives gives a sequence of them.

$$\frac{df(a)}{da} = \frac{1}{2\pi i} \oint \frac{f(z)}{(z-a)^2} dz,$$

$$f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$

This  $m=1$ -pole constant- $a_{-1}$  formula is just the first in a series of *Laurent coefficient expressions*.

$$\dots a_{-3} = \frac{1}{2\pi i} \oint z^2 f(z)dz, \quad a_{-2} = \frac{1}{2\pi i} \oint z^1 f(z)dz, \quad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz, \quad a_0 = \frac{1}{2\pi i} \oint \frac{f(z)}{z} dz, \quad a_1 = \frac{1}{2\pi i} \oint \frac{f(z)}{z^2} dz, \dots$$

Source analysis starts with 1-pole loop integrals  $\oint z^{-1}dz = 2\pi i$  or, with origin shifted  $\oint (z-a)^{-1}dz = 2\pi i$ .

They hold for any loop about point- $a$ . Function  $f(z)$  is just  $f(a)$  on a *tiny* circle around point- $a$ .

(assume *tiny* circle around  $z=a$ )

$$\oint \frac{f(z)}{z-a} dz = \oint \frac{f(a)}{z-a} dz = f(a) \oint \frac{1}{z-a} dz = 2\pi i f(a)$$

(but any contour that doesn't "touch  $a$  gives same answer)

$$f(a) = \frac{1}{2\pi i} \oint \frac{f(z)}{z-a} dz$$

The  $f(a)$  result is called a *Cauchy integral*. Then repeated  $a$ -derivatives gives a sequence of them.

$$\frac{df(a)}{da} = \frac{1}{2\pi i} \oint \frac{f(z)}{(z-a)^2} dz, \quad \frac{d^2 f(a)}{da^2} = \frac{2}{2\pi i} \oint \frac{f(z)}{(z-a)^3} dz,$$



$$f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$

This  $m=1$ -pole constant- $a_{-1}$  formula is just the first in a series of *Laurent coefficient expressions*.

$$\dots a_{-3} = \frac{1}{2\pi i} \oint z^2 f(z)dz, \quad a_{-2} = \frac{1}{2\pi i} \oint z^1 f(z)dz, \quad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz, \quad a_0 = \frac{1}{2\pi i} \oint \frac{f(z)}{z} dz, \quad a_1 = \frac{1}{2\pi i} \oint \frac{f(z)}{z^2} dz, \dots$$

Source analysis starts with 1-pole loop integrals  $\oint z^{-1}dz = 2\pi i$  or, with origin shifted  $\oint (z-a)^{-1}dz = 2\pi i$ .

They hold for any loop about point- $a$ . Function  $f(z)$  is just  $f(a)$  on a *tiny* circle around point- $a$ .

(assume *tiny* circle around  $z=a$ )

$$\oint \frac{f(z)}{z-a} dz = \oint \frac{f(a)}{z-a} dz = f(a) \oint \frac{1}{z-a} dz = 2\pi i f(a)$$

(but any contour that doesn't "touch  $a$  gives same answer)

$$f(a) = \frac{1}{2\pi i} \oint \frac{f(z)}{z-a} dz$$

The  $f(a)$  result is called a *Cauchy integral*. Then repeated  $a$ -derivatives gives a sequence of them.

$$\frac{df(a)}{da} = \frac{1}{2\pi i} \oint \frac{f(z)}{(z-a)^2} dz, \quad \frac{d^2 f(a)}{da^2} = \frac{2}{2\pi i} \oint \frac{f(z)}{(z-a)^3} dz, \quad \frac{d^3 f(a)}{da^3} = \frac{3!}{2\pi i} \oint \frac{f(z)}{(z-a)^4} dz$$

$$f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$

This  $m=1$ -pole constant- $a_{-1}$  formula is just the first in a series of *Laurent coefficient expressions*.

$$\dots a_{-3} = \frac{1}{2\pi i} \oint z^2 f(z)dz, \quad a_{-2} = \frac{1}{2\pi i} \oint z^1 f(z)dz, \quad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz, \quad a_0 = \frac{1}{2\pi i} \oint \frac{f(z)}{z} dz, \quad a_1 = \frac{1}{2\pi i} \oint \frac{f(z)}{z^2} dz, \dots$$

Source analysis starts with 1-pole loop integrals  $\oint z^{-1}dz = 2\pi i$  or, with origin shifted  $\oint (z-a)^{-1}dz = 2\pi i$ .

They hold for any loop about point- $a$ . Function  $f(z)$  is just  $f(a)$  on a *tiny* circle around point- $a$ .

(assume *tiny* circle around  $z=a$ )

$$\oint \frac{f(z)}{z-a} dz = \oint \frac{f(a)}{z-a} dz = f(a) \oint \frac{1}{z-a} dz = 2\pi i f(a)$$

(but any contour that doesn't "touch  $a$  gives same answer)

$$f(a) = \frac{1}{2\pi i} \oint \frac{f(z)}{z-a} dz$$

The  $f(a)$  result is called a *Cauchy integral*. Then repeated  $a$ -derivatives gives a sequence of them.

$$\frac{df(a)}{da} = \frac{1}{2\pi i} \oint \frac{f(z)}{(z-a)^2} dz, \quad \frac{d^2 f(a)}{da^2} = \frac{2}{2\pi i} \oint \frac{f(z)}{(z-a)^3} dz, \quad \frac{d^3 f(a)}{da^3} = \frac{3!}{2\pi i} \oint \frac{f(z)}{(z-a)^4} dz, \quad \dots, \quad \frac{d^n f(a)}{da^n} = \frac{n!}{2\pi i} \oint \frac{f(z)}{(z-a)^{n+1}} dz$$

$$f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$

This  $m=1$ -pole constant- $a_{-1}$  formula is just the first in a series of *Laurent coefficient expressions*.

$$\dots a_{-3} = \frac{1}{2\pi i} \oint z^2 f(z)dz, \quad a_{-2} = \frac{1}{2\pi i} \oint z^1 f(z)dz, \quad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz, \quad a_0 = \frac{1}{2\pi i} \oint \frac{f(z)}{z} dz, \quad a_1 = \frac{1}{2\pi i} \oint \frac{f(z)}{z^2} dz, \dots$$

Source analysis starts with 1-pole loop integrals  $\oint z^{-1}dz = 2\pi i$  or, with origin shifted  $\oint (z-a)^{-1}dz = 2\pi i$ .

They hold for any loop about point- $a$ . Function  $f(z)$  is just  $f(a)$  on a *tiny* circle around point- $a$ .

(assume *tiny* circle around  $z=a$ )

$$\oint \frac{f(z)}{z-a} dz = \oint \frac{f(a)}{z-a} dz = f(a) \oint \frac{1}{z-a} dz = 2\pi i f(a)$$

(but any contour that doesn't "touch  $a$  gives same answer)

$$f(a) = \frac{1}{2\pi i} \oint \frac{f(z)}{z-a} dz$$

The  $f(a)$  result is called a *Cauchy integral*. Then repeated  $a$ -derivatives gives a sequence of them.

$$\frac{df(a)}{da} = \frac{1}{2\pi i} \oint \frac{f(z)}{(z-a)^2} dz, \quad \frac{d^2 f(a)}{da^2} = \frac{2}{2\pi i} \oint \frac{f(z)}{(z-a)^3} dz, \quad \frac{d^3 f(a)}{da^3} = \frac{3!}{2\pi i} \oint \frac{f(z)}{(z-a)^4} dz, \quad \dots, \quad \frac{d^n f(a)}{da^n} = \frac{n!}{2\pi i} \oint \frac{f(z)}{(z-a)^{n+1}} dz$$

This leads to a general *Taylor-Laurent* power series expansion of function  $f(z)$  around point- $a$ .

$$f(z) = \dots a_{-3}z^{-3} + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$

This  $m=1$ -pole constant- $a_{-1}$  formula is just the first in a series of *Laurent coefficient expressions*.

$$\dots a_{-3} = \frac{1}{2\pi i} \oint z^2 f(z)dz, \quad a_{-2} = \frac{1}{2\pi i} \oint z^1 f(z)dz, \quad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz, \quad a_0 = \frac{1}{2\pi i} \oint \frac{f(z)}{z} dz, \quad a_1 = \frac{1}{2\pi i} \oint \frac{f(z)}{z^2} dz, \dots$$

Source analysis starts with 1-pole loop integrals  $\oint z^{-1}dz = 2\pi i$  or, with origin shifted  $\oint (z-a)^{-1}dz = 2\pi i$ .

They hold for any loop about point- $a$ . Function  $f(z)$  is just  $f(a)$  on a *tiny* circle around point- $a$ .

(assume *tiny* circle around  $z=a$ )

$$\oint \frac{f(z)}{z-a} dz = \oint \frac{f(a)}{z-a} dz = f(a) \oint \frac{1}{z-a} dz = 2\pi i f(a)$$

(but any contour that doesn't "touch  $a$  gives same answer)

$$f(a) = \frac{1}{2\pi i} \oint \frac{f(z)}{z-a} dz$$

The  $f(a)$  result is called a *Cauchy integral*. Then repeated  $a$ -derivatives gives a sequence of them.

$$\frac{df(a)}{da} = \frac{1}{2\pi i} \oint \frac{f(z)}{(z-a)^2} dz, \quad \frac{d^2 f(a)}{da^2} = \frac{2}{2\pi i} \oint \frac{f(z)}{(z-a)^3} dz, \quad \frac{d^3 f(a)}{da^3} = \frac{3!}{2\pi i} \oint \frac{f(z)}{(z-a)^4} dz, \dots, \frac{d^n f(a)}{da^n} = \frac{n!}{2\pi i} \oint \frac{f(z)}{(z-a)^{n+1}} dz$$

This leads to a general *Taylor-Laurent* power series expansion of function  $f(z)$  around point- $a$ .

$$f(z) = \sum_{n=-\infty}^{\infty} a_n (z-a)^n \qquad \text{where : } a_n = \frac{1}{2\pi i} \oint \frac{f(z)}{(z-a)^{n+1}} dz \left( = \frac{1}{n!} \frac{d^n f(a)}{da^n} \quad \text{for : } n \geq 0 \right)$$

Of all  $2^m$ -pole field terms  $a_{m-1}z^{m-1}$ , only the  $m=0$  monopole  $a_{-1}z^{-1}$  has a non-zero loop integral (10.39).

$$\oint f(z)dz = \oint a_{-1}z^{-1}dz = 2\pi i a_{-1} \qquad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz$$

This  $m=1$ -pole constant- $a_{-1}$  formula is just the first in a series of *Laurent coefficient expressions*.

$$\dots a_{-3} = \frac{1}{2\pi i} \oint z^2 f(z)dz, \quad a_{-2} = \frac{1}{2\pi i} \oint z^1 f(z)dz, \quad a_{-1} = \frac{1}{2\pi i} \oint f(z)dz, \quad a_0 = \frac{1}{2\pi i} \oint \frac{f(z)}{z} dz, \quad a_1 = \frac{1}{2\pi i} \oint \frac{f(z)}{z^2} dz, \dots$$

Source analysis starts with 1-pole loop integrals  $\oint z^{-1}dz = 2\pi i$  or, with origin shifted  $\oint (z-a)^{-1}dz = 2\pi i$ .

They hold for any loop about point- $a$ . Function  $f(z)$  is just  $f(a)$  on a *tiny* circle around point- $a$ .

(assume *tiny* circle around  $z=a$ )

$$\oint \frac{f(z)}{z-a} dz = \oint \frac{f(a)}{z-a} dz = f(a) \oint \frac{1}{z-a} dz = 2\pi i f(a)$$

(but any contour that doesn't "touch  $a$  gives same answer)

$$f(a) = \frac{1}{2\pi i} \oint \frac{f(z)}{z-a} dz$$

The  $f(a)$  result is called a *Cauchy integral*. Then repeated  $a$ -derivatives gives a sequence of them.

$$\frac{df(a)}{da} = \frac{1}{2\pi i} \oint \frac{f(z)}{(z-a)^2} dz, \quad \frac{d^2 f(a)}{da^2} = \frac{2}{2\pi i} \oint \frac{f(z)}{(z-a)^3} dz, \quad \frac{d^3 f(a)}{da^3} = \frac{3!}{2\pi i} \oint \frac{f(z)}{(z-a)^4} dz, \dots, \frac{d^n f(a)}{da^n} = \frac{n!}{2\pi i} \oint \frac{f(z)}{(z-a)^{n+1}} dz$$

This leads to a general *Taylor-Laurent* power series expansion of function  $f(z)$  around point- $a$ .

$$f(z) = \sum_{n=-\infty}^{\infty} a_n (z-a)^n \qquad \text{where : } a_n = \frac{1}{2\pi i} \oint \frac{f(z)}{(z-a)^{n+1}} dz \left( = \frac{1}{n!} \frac{d^n f(a)}{da^n} \quad \text{for : } n \geq 0 \right)$$

(*quadrupole*)<sub>0</sub> (*dipole*)<sub>0</sub> (*monopole*) (*dipole*)<sub>∞</sub> (*quadrupole*)<sub>∞</sub> (*octapole*)<sub>∞</sub> (*hexadecapole*)<sub>∞</sub> ...

$$f(z) = \dots a_{-3}z^{-3} + \underbrace{a_{-2}z^{-2}}_{\text{dipole moment}} + \underbrace{a_{-1}z^{-1}}_{\text{monopole moment}} + a_0 + a_1z + a_2z^2 + a_3z^3 + a_4z^4 + a_5z^5 + \dots$$



## *5. Mapping and Non-analytic 2D source field analysis*

The *half-n'-half* results

are called

*Riemann-Cauchy*

*Derivative Relations*

$\frac{\partial \Phi}{\partial x} = \frac{\partial A}{\partial y}$	is:	$\frac{\partial \operatorname{Re}\phi(z)}{\partial x} = \frac{\partial \operatorname{Im}\phi(z)}{\partial y}$	or:	$\frac{\partial \operatorname{Re}f(z)}{\partial x} = \frac{\partial \operatorname{Im}f(z)}{\partial y}$	is:	$\frac{\partial f_x(z)}{\partial x} = \frac{\partial f_y(z)}{\partial y}$
$\frac{\partial \Phi}{\partial y} = -\frac{\partial A}{\partial x}$	is:	$\frac{\partial \operatorname{Re}\phi(z)}{\partial y} = -\frac{\partial \operatorname{Im}\phi(z)}{\partial x}$	or:	$\frac{\partial \operatorname{Re}f(z)}{\partial y} = -\frac{\partial \operatorname{Im}f(z)}{\partial x}$	is:	$\frac{\partial f_x(z)}{\partial y} = -\frac{\partial f_y(z)}{\partial x}$


*RC applies to analytic potential  $\phi(z) = \Phi + iA$  and analytic field  $f(z) = f_x + if_y$  and any analytic function*

The *half-n'-half* results  
are called  
*Riemann-Cauchy*  
*Derivative Relations*

$\frac{\partial \Phi}{\partial x} = \frac{\partial A}{\partial y}$	is:	$\frac{\partial \operatorname{Re} \phi(z)}{\partial x} = \frac{\partial \operatorname{Im} \phi(z)}{\partial y}$	or:	$\frac{\partial \operatorname{Re} f(z)}{\partial x} = \frac{\partial \operatorname{Im} f(z)}{\partial y}$	is:	$\frac{\partial f_x(z)}{\partial x} = \frac{\partial f_y(z)}{\partial y}$
$\frac{\partial \Phi}{\partial y} = -\frac{\partial A}{\partial x}$	is:	$\frac{\partial \operatorname{Re} \phi(z)}{\partial y} = -\frac{\partial \operatorname{Im} \phi(z)}{\partial x}$	or:	$\frac{\partial \operatorname{Re} f(z)}{\partial y} = -\frac{\partial \operatorname{Im} f(z)}{\partial x}$	is:	$\frac{\partial f_x(z)}{\partial y} = -\frac{\partial f_y(z)}{\partial x}$

RC applies to analytic potential  $\phi(z) = \Phi + iA$  and analytic field  $f(z) = f_x + if_y$  and any analytic function

Common notation for mapping:  $w(z) = u + iv$




The *half-n'-half* results  
are called  
*Riemann-Cauchy  
Derivative Relations*

$\frac{\partial \Phi}{\partial x} = \frac{\partial A}{\partial y}$	is:	$\frac{\partial \operatorname{Re}\phi(z)}{\partial x} = \frac{\partial \operatorname{Im}\phi(z)}{\partial y}$	or:	$\frac{\partial \operatorname{Re}f(z)}{\partial x} = \frac{\partial \operatorname{Im}f(z)}{\partial y}$	is:	$\frac{\partial f_x(z)}{\partial x} = \frac{\partial f_y(z)}{\partial y}$
$\frac{\partial \Phi}{\partial y} = -\frac{\partial A}{\partial x}$	is:	$\frac{\partial \operatorname{Re}\phi(z)}{\partial y} = -\frac{\partial \operatorname{Im}\phi(z)}{\partial x}$	or:	$\frac{\partial \operatorname{Re}f(z)}{\partial y} = -\frac{\partial \operatorname{Im}f(z)}{\partial x}$	is:	$\frac{\partial f_x(z)}{\partial y} = -\frac{\partial f_y(z)}{\partial x}$

RC applies to analytic potential  $\phi(z) = \Phi + iA$  and analytic field  $f(z) = f_x + if_y$  and any analytic function

Common notation for mapping:  $w(z) = u + iv$



Jacobian for mapping:

$$du = \frac{\partial u}{\partial x} dx + \frac{\partial u}{\partial y} dy$$

$$dv = \frac{\partial v}{\partial x} dx + \frac{\partial v}{\partial y} dy$$

$$\begin{pmatrix} du \\ dv \end{pmatrix} = \begin{pmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{pmatrix} \begin{pmatrix} dx \\ dy \end{pmatrix}$$

Complex derivative for mapping:


$$\frac{dw}{dz} = \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) (u + iv) = \frac{1}{2} \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right)$$

The *half-n'-half* results  
are called  
*Riemann-Cauchy  
Derivative Relations*

$\frac{\partial \Phi}{\partial x} = \frac{\partial A}{\partial y}$	is:	$\frac{\partial \operatorname{Re}\phi(z)}{\partial x} = \frac{\partial \operatorname{Im}\phi(z)}{\partial y}$	or:	$\frac{\partial \operatorname{Re}f(z)}{\partial x} = \frac{\partial \operatorname{Im}f(z)}{\partial y}$	is:	$\frac{\partial f_x(z)}{\partial x} = \frac{\partial f_y(z)}{\partial y}$
$\frac{\partial \Phi}{\partial y} = -\frac{\partial A}{\partial x}$	is:	$\frac{\partial \operatorname{Re}\phi(z)}{\partial y} = -\frac{\partial \operatorname{Im}\phi(z)}{\partial x}$	or:	$\frac{\partial \operatorname{Re}f(z)}{\partial y} = -\frac{\partial \operatorname{Im}f(z)}{\partial x}$	is:	$\frac{\partial f_x(z)}{\partial y} = -\frac{\partial f_y(z)}{\partial x}$

RC applies to analytic potential  $\phi(z) = \Phi + iA$  and analytic field  $f(z) = f_x + if_y$  and any analytic function

Common notation for mapping:  $w(z) = u + iv$



Jacobian for mapping:

$$du = \frac{\partial u}{\partial x} dx + \frac{\partial u}{\partial y} dy$$

$$dv = \frac{\partial v}{\partial x} dx + \frac{\partial v}{\partial y} dy$$

$$\begin{pmatrix} du \\ dv \end{pmatrix} = \begin{pmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{pmatrix} \begin{pmatrix} dx \\ dy \end{pmatrix}$$

$$= \begin{pmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ -\frac{\partial u}{\partial y} & \frac{\partial u}{\partial x} \end{pmatrix} \begin{pmatrix} dx \\ dy \end{pmatrix} = \begin{pmatrix} \frac{\partial v}{\partial y} & -\frac{\partial v}{\partial x} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{pmatrix} \begin{pmatrix} dx \\ dy \end{pmatrix}$$

Complex derivative for mapping:

$$\frac{dw}{dz} = \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) (u + iv) = \frac{1}{2} \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right)$$

$$= \frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y} = \frac{\partial v}{\partial y} + i \frac{\partial v}{\partial x}$$

Complex derivative abs-square:

$$\left| \frac{dw}{dz} \right|^2 = \left( \frac{\partial u}{\partial x} \right)^2 + \left( \frac{\partial u}{\partial y} \right)^2 = \left( \frac{\partial v}{\partial y} \right)^2 + \left( \frac{\partial v}{\partial x} \right)^2$$




The *half-n'-half* results  
are called  
*Riemann-Cauchy  
Derivative Relations*

$\frac{\partial \Phi}{\partial x} = \frac{\partial A}{\partial y}$	is:	$\frac{\partial \operatorname{Re}\phi(z)}{\partial x} = \frac{\partial \operatorname{Im}\phi(z)}{\partial y}$	or:	$\frac{\partial \operatorname{Re}f(z)}{\partial x} = \frac{\partial \operatorname{Im}f(z)}{\partial y}$	is:	$\frac{\partial f_x(z)}{\partial x} = \frac{\partial f_y(z)}{\partial y}$
$\frac{\partial \Phi}{\partial y} = -\frac{\partial A}{\partial x}$	is:	$\frac{\partial \operatorname{Re}\phi(z)}{\partial y} = -\frac{\partial \operatorname{Im}\phi(z)}{\partial x}$	or:	$\frac{\partial \operatorname{Re}f(z)}{\partial y} = -\frac{\partial \operatorname{Im}f(z)}{\partial x}$	is:	$\frac{\partial f_x(z)}{\partial y} = -\frac{\partial f_y(z)}{\partial x}$

RC applies to analytic potential  $\phi(z) = \Phi + iA$  and analytic field  $f(z) = f_x + if_y$  and any analytic function

Common notation for mapping:  $w(z) = u + iv$



Jacobian for mapping:

$$du = \frac{\partial u}{\partial x} dx + \frac{\partial u}{\partial y} dy$$

$$dv = \frac{\partial v}{\partial x} dx + \frac{\partial v}{\partial y} dy$$

$$\begin{pmatrix} du \\ dv \end{pmatrix} = \begin{pmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{pmatrix} \begin{pmatrix} dx \\ dy \end{pmatrix}$$

$$= \begin{pmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ -\frac{\partial u}{\partial y} & \frac{\partial u}{\partial x} \end{pmatrix} \begin{pmatrix} dx \\ dy \end{pmatrix} = \begin{pmatrix} \frac{\partial v}{\partial y} & -\frac{\partial v}{\partial x} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{pmatrix} \begin{pmatrix} dx \\ dy \end{pmatrix}$$

Complex derivative for mapping:

$$\frac{dw}{dz} = \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) (u + iv) = \frac{1}{2} \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right)$$

$$= \frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y} = \frac{\partial v}{\partial y} + i \frac{\partial v}{\partial x}$$

Complex derivative abs-square:

$$\left| \frac{dw}{dz} \right|^2 = \left( \frac{\partial u}{\partial x} \right)^2 + \left( \frac{\partial u}{\partial y} \right)^2 = \left( \frac{\partial v}{\partial y} \right)^2 + \left( \frac{\partial v}{\partial x} \right)^2 = \det|J|$$


...equals Jacobian Determinant

The *half-n'-half* results  
are called  
*Riemann-Cauchy  
Derivative Relations*

$\frac{\partial \Phi}{\partial x} = \frac{\partial A}{\partial y}$	is:	$\frac{\partial \text{Re}\phi(z)}{\partial x} = \frac{\partial \text{Im}\phi(z)}{\partial y}$	or:	$\frac{\partial \text{Re}f(z)}{\partial x} = \frac{\partial \text{Im}f(z)}{\partial y}$	is:	$\frac{\partial f_x(z)}{\partial x} = \frac{\partial f_y(z)}{\partial y}$
$\frac{\partial \Phi}{\partial y} = -\frac{\partial A}{\partial x}$	is:	$\frac{\partial \text{Re}\phi(z)}{\partial y} = -\frac{\partial \text{Im}\phi(z)}{\partial x}$	or:	$\frac{\partial \text{Re}f(z)}{\partial y} = -\frac{\partial \text{Im}f(z)}{\partial x}$	is:	$\frac{\partial f_x(z)}{\partial y} = -\frac{\partial f_y(z)}{\partial x}$

RC applies to analytic potential  $\phi(z) = \Phi + iA$  and analytic field  $f(z) = f_x + if_y$  and any analytic function

Common notation for mapping:  $w(z) = u + iv$



Important result:  
 $dw = \sqrt{J} \cdot e^{i\theta} \cdot dz$   
is scaled rotation of  $dz$ .

$$du = \frac{\partial u}{\partial x} dx + \frac{\partial u}{\partial y} dy$$

$$dv = \frac{\partial v}{\partial x} dx + \frac{\partial v}{\partial y} dy$$

Jacobian for mapping is scaled rotation:

$$\begin{pmatrix} du \\ dv \end{pmatrix} = \begin{pmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{pmatrix} \begin{pmatrix} dx \\ dy \end{pmatrix} = \sqrt{\det J} \begin{pmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{pmatrix} \begin{pmatrix} dx \\ dy \end{pmatrix}$$

$$= \begin{pmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ -\frac{\partial u}{\partial y} & \frac{\partial u}{\partial x} \end{pmatrix} \begin{pmatrix} dx \\ dy \end{pmatrix} = \begin{pmatrix} \frac{\partial v}{\partial y} & -\frac{\partial v}{\partial x} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{pmatrix} \begin{pmatrix} dx \\ dy \end{pmatrix}$$

Complex derivative for mapping:

$$\frac{dw}{dz} = \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) (u + iv) = \frac{1}{2} \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right)$$



$$= \frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y} = \frac{\partial v}{\partial y} + i \frac{\partial v}{\partial x}$$

Complex derivative abs-square:



$$\left| \frac{dw}{dz} \right|^2 = \left( \frac{\partial u}{\partial x} \right)^2 + \left( \frac{\partial u}{\partial y} \right)^2 = \left( \frac{\partial v}{\partial y} \right)^2 + \left( \frac{\partial v}{\partial x} \right)^2 = \det|J|$$

...equals Jacobian Determinant


$w(z) = z^2$  gives parabolic OCC



$w(z) = z^2$  gives parabolic OCC



Inverse:  $z(w) = w^{1/2}$  gives hyperbolic OCC



## *5. Mapping and Non-analytic 2D source field analysis*



*Non-analytic potential, force, and source field functions (Excerpts of Unit 1-Ch.10 and AnalyIt)*

A general 2D complex field may have:

1. non-analytic *potential field function*  $\phi(z, z^*) = \Phi(x, y) + iA(x, y)$ ,
2. non-analytic *force field function*  $f(z, z^*) = f_x(x, y) + if_y(x, y)$  ,
3. non-analytic *source distribution function*  $s(z, z^*) = \rho(x, y) + i I(x, y)$ .

## *Non-analytic potential, force, and source field functions (Excerpts of Unit 1-Ch.10 and AnalyIt)*

A general 2D complex field may have:

1. non-analytic *potential field function*  $\phi(z, z^*) = \Phi(x, y) + iA(x, y)$ ,
2. non-analytic *force field function*  $f(z, z^*) = f_x(x, y) + if_y(x, y)$ ,
3. non-analytic *source distribution function*  $s(z, z^*) = \rho(x, y) + i I(x, y)$ .

Source definitions generalize source-free fields ( $\frac{df(z^*)}{dz} = 0 = \frac{df(z)}{dz^*}$ ) based on relations.  $\frac{df}{dz} = \frac{1}{2} \left( \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial f_y}{\partial x} - \frac{\partial f_x}{\partial y} \right) = \frac{1}{2} \nabla \cdot \mathbf{F} + \frac{i}{2} |\nabla \times \mathbf{F}|$

## Non-analytic potential, force, and source field functions (Excerpts of Unit 1-Ch.10 and AnalyIt)

A general 2D complex field may have:

1. non-analytic *potential field function*  $\phi(z, z^*) = \Phi(x, y) + iA(x, y)$ ,
2. non-analytic *force field function*  $f(z, z^*) = f_x(x, y) + if_y(x, y)$ ,
3. non-analytic *source distribution function*  $s(z, z^*) = \rho(x, y) + i I(x, y)$ .

Source definitions generalize source-free fields ( $\frac{df(z^*)}{dz} = 0 = \frac{df(z)}{dz^*}$ ) based on relations.  $\frac{df}{dz} = \frac{1}{2}(\frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y}) + \frac{i}{2}(\frac{\partial f_y}{\partial x} - \frac{\partial f_x}{\partial y}) = \frac{1}{2}\nabla \cdot \mathbf{F} + \frac{i}{2}|\nabla \times \mathbf{F}|$

$$2 \frac{df^*}{dz} = s^*(z, z^*)$$

$$2 \frac{df}{dz^*} = s(z, z^*)$$

## Non-analytic potential, force, and source field functions (Excerpts of Unit 1-Ch.10 and AnalyIt)

A general 2D complex field may have:

1. non-analytic *potential field function*  $\phi(z, z^*) = \Phi(x, y) + iA(x, y)$ ,
2. non-analytic *force field function*  $f(z, z^*) = f_x(x, y) + if_y(x, y)$ ,
3. non-analytic *source distribution function*  $s(z, z^*) = \rho(x, y) + i I(x, y)$ .

Source definitions generalize source-free fields ( $\frac{df(z^*)}{dz} = 0 = \frac{df(z)}{dz^*}$ ) based on relations.  $\frac{df}{dz} = \frac{1}{2}(\frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y}) + \frac{i}{2}(\frac{\partial f_y}{\partial x} - \frac{\partial f_x}{\partial y}) = \frac{1}{2}\nabla \cdot \mathbf{F} + \frac{i}{2}|\nabla \times \mathbf{F}|$

$$2 \frac{df^*}{dz} = s^*(z, z^*)$$

$$2 \frac{df}{dz^*} = s(z, z^*)$$

Field- $f$ -from-potential- $\phi$  equations are like the older ( $f(z) = \frac{d\phi}{dz}$  or  $f^*(z^*) = \frac{d\phi^*}{dz^*}$ ) but with an extra factor of 2.

$$2 \frac{d\phi}{dz} = f(z, z^*)$$

$$2 \frac{d\phi^*}{dz^*} = f^*(z, z^*)$$

## Non-analytic potential, force, and source field functions (Excerpts of Unit 1-Ch.10 and AnalyIt)

A general 2D complex field may have:

1. non-analytic *potential field function*  $\phi(z, z^*) = \Phi(x, y) + iA(x, y)$ ,
2. non-analytic *force field function*  $f(z, z^*) = f_x(x, y) + if_y(x, y)$ ,
3. non-analytic *source distribution function*  $s(z, z^*) = \rho(x, y) + i I(x, y)$ .

Source definitions generalize source-free fields ( $\frac{df(z^*)}{dz} = 0 = \frac{df(z)}{dz^*}$ ) based on relations.  $\frac{df}{dz} = \frac{1}{2}(\frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y}) + \frac{i}{2}(\frac{\partial f_y}{\partial x} - \frac{\partial f_x}{\partial y}) = \frac{1}{2}\nabla \cdot \mathbf{F} + \frac{i}{2}|\nabla \times \mathbf{F}|$

$$2 \frac{df^*}{dz} = s^*(z, z^*) \qquad 2 \frac{df}{dz^*} = s(z, z^*)$$

Field- $f$ -from-potential- $\phi$  equations are like the older ( $f(z) = \frac{d\phi}{dz}$  or  $f^*(z^*) = \frac{d\phi^*}{dz^*}$ ) but with an extra factor of 2.

$$2 \frac{d\phi}{dz} = f(z, z^*) \qquad 2 \frac{d\phi^*}{dz^*} = f^*(z, z^*)$$

The new source equations expand into a real and imaginary parts that are divergence and curl terms, respectively.

$$s^*(z, z^*) = 2 \frac{df^*}{dz} = \left[ \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right] \left[ f_x^*(x, y) + if_y^*(x, y) \right] = \rho - i I, \quad \text{where: } f_x^* = f_x, \text{ and: } f_y^* = -f_y$$

## Non-analytic potential, force, and source field functions (Excerpts of Unit 1-Ch.10 and AnalyIt)

A general 2D complex field may have:

1. non-analytic *potential field function*  $\phi(z, z^*) = \Phi(x, y) + iA(x, y)$ ,
2. non-analytic *force field function*  $f(z, z^*) = f_x(x, y) + if_y(x, y)$ ,
3. non-analytic *source distribution function*  $s(z, z^*) = \rho(x, y) + i I(x, y)$ .

Source definitions generalize source-free fields ( $\frac{df(z^*)}{dz} = 0 = \frac{df(z)}{dz^*}$ ) based on relations.  $\frac{df}{dz} = \frac{1}{2} \left( \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial f_y}{\partial x} - \frac{\partial f_x}{\partial y} \right) = \frac{1}{2} \nabla \cdot \mathbf{F} + \frac{i}{2} |\nabla \times \mathbf{F}|$

$$2 \frac{df^*}{dz} = s^*(z, z^*) \qquad 2 \frac{df}{dz^*} = s(z, z^*)$$

Field- $f$ -from-potential- $\phi$  equations are like the older ( $f(z) = \frac{d\phi}{dz}$  or  $f^*(z^*) = \frac{d\phi^*}{dz^*}$ ) but with an extra factor of 2.

$$2 \frac{d\phi}{dz} = f(z, z^*) \qquad 2 \frac{d\phi^*}{dz^*} = f^*(z, z^*)$$

The new source equations expand into a real and imaginary parts that are divergence and curl terms, respectively.

$$\begin{aligned} s^*(z, z^*) = 2 \frac{df^*}{dz} &= \left[ \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right] \left[ f_x^*(x, y) + if_y^*(x, y) \right] = \rho - i I, \quad \text{where: } f_x^* = f_x, \text{ and: } f_y^* = -f_y \\ &= \left[ \frac{\partial f_x^*}{\partial x} + \frac{\partial f_y^*}{\partial y} \right] + i \left[ \frac{\partial f_y^*}{\partial x} - \frac{\partial f_x^*}{\partial y} \right] = \left[ \nabla \cdot \mathbf{f}^* \right] + i \left[ \nabla \times \mathbf{f}^* \right]_z \end{aligned}$$



## Non-analytic potential, force, and source field functions (Excerpts of Unit 1-Ch.10 and AnalyIt)

A general 2D complex field may have:

1. non-analytic *potential field function*  $\phi(z, z^*) = \Phi(x, y) + iA(x, y)$ ,
2. non-analytic *force field function*  $f(z, z^*) = f_x(x, y) + if_y(x, y)$ ,
3. non-analytic *source distribution function*  $s(z, z^*) = \rho(x, y) + i I(x, y)$ .

Source definitions generalize source-free fields ( $\frac{df(z^*)}{dz} = 0 = \frac{df(z)}{dz^*}$ ) based on relations.  $\frac{df}{dz} = \frac{1}{2} \left( \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial f_y}{\partial x} - \frac{\partial f_x}{\partial y} \right) = \frac{1}{2} \nabla \cdot \mathbf{F} + \frac{i}{2} |\nabla \times \mathbf{F}|$

$$2 \frac{df^*}{dz} = s^*(z, z^*) \qquad 2 \frac{df}{dz^*} = s(z, z^*)$$

Field- $f$ -from-potential- $\phi$  equations are like the older ( $f(z) = \frac{d\phi}{dz}$  or  $f^*(z^*) = \frac{d\phi^*}{dz^*}$ ) but with an extra factor of 2.

$$2 \frac{d\phi}{dz} = f(z, z^*) \qquad 2 \frac{d\phi^*}{dz^*} = f^*(z, z^*)$$

The new source equations expand into a real and imaginary parts that are divergence and curl terms, respectively.

$$\begin{aligned} s^*(z, z^*) = 2 \frac{df^*}{dz} &= \left[ \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right] \left[ f_x^*(x, y) + if_y^*(x, y) \right] = \rho - i I, \quad \text{where: } f_x^* = f_x, \text{ and: } f_y^* = -f_y \\ &= \left[ \frac{\partial f_x^*}{\partial x} + \frac{\partial f_y^*}{\partial y} \right] + i \left[ \frac{\partial f_y^*}{\partial x} - \frac{\partial f_x^*}{\partial y} \right] = \left[ \nabla \cdot \mathbf{f}^* \right] + i \left[ \nabla \times \mathbf{f}^* \right]_z \end{aligned}$$

Real part: *Poisson scalar source equation* (charge density  $\rho$ ).      Imaginary part: *Biot-Savart vector source equation* (current density  $I$ )

$$\nabla \cdot \mathbf{f}^* = \rho$$

$$\nabla \times \mathbf{f}^* = -I$$

## Non-analytic potential, force, and source field functions (Excerpts of Unit 1-Ch.10 and AnalyIt)

A general 2D complex field may have:

1. non-analytic *potential field function*  $\phi(z, z^*) = \Phi(x, y) + iA(x, y)$ ,
2. non-analytic *force field function*  $f(z, z^*) = f_x(x, y) + if_y(x, y)$ ,
3. non-analytic *source distribution function*  $s(z, z^*) = \rho(x, y) + i I(x, y)$ .

Source definitions generalize source-free fields ( $\frac{df(z^*)}{dz} = 0 = \frac{df(z)}{dz^*}$ ) based on relations.  $\frac{df}{dz} = \frac{1}{2} \left( \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial f_y}{\partial x} - \frac{\partial f_x}{\partial y} \right) = \frac{1}{2} \nabla \cdot \mathbf{F} + \frac{i}{2} |\nabla \times \mathbf{F}|$

$$2 \frac{df^*}{dz} = s^*(z, z^*) \qquad 2 \frac{df}{dz^*} = s(z, z^*)$$

Field- $f$ -from-potential- $\phi$  equations are like the older ( $f(z) = \frac{d\phi}{dz}$  or  $f^*(z^*) = \frac{d\phi^*}{dz^*}$ ) but with an extra factor of 2.

$$2 \frac{d\phi}{dz} = f(z, z^*) \qquad 2 \frac{d\phi^*}{dz^*} = f^*(z, z^*)$$

The new source equations expand into a real and imaginary parts that are divergence and curl terms, respectively.

$$\begin{aligned} s^*(z, z^*) = 2 \frac{df^*}{dz} &= \left[ \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right] \left[ f_x^*(x, y) + if_y^*(x, y) \right] = \rho - i I, \quad \text{where: } f_x^* = f_x, \text{ and: } f_y^* = -f_y \\ &= \left[ \frac{\partial f_x^*}{\partial x} + \frac{\partial f_y^*}{\partial y} \right] + i \left[ \frac{\partial f_y^*}{\partial x} - \frac{\partial f_x^*}{\partial y} \right] = \left[ \nabla \cdot \mathbf{f}^* \right] + i \left[ \nabla \times \mathbf{f}^* \right]_z \end{aligned}$$

Real part: *Poisson scalar source equation* (charge density  $\rho$ ).      Imaginary part: *Biot-Savart vector source equation* (current density  $I$ )

$$\nabla \cdot \mathbf{f}^* = \rho$$

$$\nabla \times \mathbf{f}^* = -I$$

Field-vs-potential has Re and Im parts that are  $x$  and  $y$  components of  $\text{grad } \Phi$  and  $\text{curl } A_z$  from potential  $\phi = \Phi + iA$  or  $\phi^* = \Phi - iA$ .

$$f^*(z, z^*) = 2 \frac{d\phi^*}{dz^*} = \left[ \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right] (\Phi - iA) = f_x^* + if_y^*$$

## Non-analytic potential, force, and source field functions (Excerpts of Unit 1-Ch.10 and AnalyIt)

A general 2D complex field may have:

1. non-analytic *potential field function*  $\phi(z, z^*) = \Phi(x, y) + iA(x, y)$ ,
2. non-analytic *force field function*  $f(z, z^*) = f_x(x, y) + if_y(x, y)$ ,
3. non-analytic *source distribution function*  $s(z, z^*) = \rho(x, y) + i I(x, y)$ .

Source definitions generalize source-free fields ( $\frac{df(z^*)}{dz} = 0 = \frac{df(z)}{dz^*}$ ) based on relations.  $\frac{df}{dz} = \frac{1}{2} \left( \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial f_y}{\partial x} - \frac{\partial f_x}{\partial y} \right) = \frac{1}{2} \nabla \cdot \mathbf{F} + \frac{i}{2} |\nabla \times \mathbf{F}|$

$$2 \frac{df^*}{dz} = s^*(z, z^*) \qquad 2 \frac{df}{dz^*} = s(z, z^*)$$

Field- $f$ -from-potential- $\phi$  equations are like the older ( $f(z) = \frac{d\phi}{dz}$  or  $f^*(z^*) = \frac{d\phi^*}{dz^*}$ ) but with an extra factor of 2.

$$2 \frac{d\phi}{dz} = f(z, z^*) \qquad 2 \frac{d\phi^*}{dz^*} = f^*(z, z^*)$$

The new source equations expand into a real and imaginary parts that are divergence and curl terms, respectively.

$$\begin{aligned} s^*(z, z^*) = 2 \frac{df^*}{dz} &= \left[ \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right] \left[ f_x^*(x, y) + if_y^*(x, y) \right] = \rho - i I, \quad \text{where: } f_x^* = f_x, \text{ and: } f_y^* = -f_y \\ &= \left[ \frac{\partial f_x^*}{\partial x} + \frac{\partial f_y^*}{\partial y} \right] + i \left[ \frac{\partial f_y^*}{\partial x} - \frac{\partial f_x^*}{\partial y} \right] = \left[ \nabla \cdot \mathbf{f}^* \right] + i \left[ \nabla \times \mathbf{f}^* \right]_z \end{aligned}$$

Real part: *Poisson scalar source equation (charge density  $\rho$ )*.      Imaginary part: *Biot-Savart vector source equation (current density  $I$ )*

$$\nabla \cdot \mathbf{f}^* = \rho$$

$$\nabla \times \mathbf{f}^* = -I$$

Field-vs-potential has Re and Im parts that are  $x$  and  $y$  components of  $\text{grad } \Phi$  and  $\text{curl } A_z$  from potential  $\phi = \Phi + iA$  or  $\phi^* = \Phi - iA$ .

$$\begin{aligned} f^*(z, z^*) = 2 \frac{d\phi^*}{dz^*} &= \left[ \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right] (\Phi - iA) = f_x^* + if_y^* \\ &= \left[ \frac{\partial \Phi}{\partial x} + i \frac{\partial \Phi}{\partial y} \right] + \left[ \frac{\partial A}{\partial y} - i \frac{\partial A}{\partial x} \right] = \left[ \nabla \Phi \right] + \left[ \nabla \times \mathbf{A}_z \right] \end{aligned} \qquad \text{(For source-free analytic functions these two fields are identical.)}$$

## Non-analytic potential, force, and source field functions (Excerpts of Unit 1-Ch.10 and AnalyIt)

A general 2D complex field may have:

1. non-analytic *potential field function*  $\phi(z, z^*) = \Phi(x, y) + iA(x, y)$ ,
2. non-analytic *force field function*  $f(z, z^*) = f_x(x, y) + if_y(x, y)$ ,
3. non-analytic *source distribution function*  $s(z, z^*) = \rho(x, y) + i I(x, y)$ .

Source definitions generalize source-free fields ( $\frac{df(z^*)}{dz} = 0 = \frac{df(z)}{dz^*}$ ) based on relations.  $\frac{df}{dz} = \frac{1}{2} \left( \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial f_y}{\partial x} - \frac{\partial f_x}{\partial y} \right) = \frac{1}{2} \nabla \cdot \mathbf{F} + \frac{i}{2} |\nabla \times \mathbf{F}|$

$$2 \frac{df^*}{dz} = s^*(z, z^*) \qquad 2 \frac{df}{dz^*} = s(z, z^*)$$

Field- $f$ -from-potential- $\phi$  equations are like the older ( $f(z) = \frac{d\phi}{dz}$  or  $f^*(z^*) = \frac{d\phi^*}{dz^*}$ ) but with an extra factor of 2.

$$2 \frac{d\phi}{dz} = f(z, z^*) \qquad 2 \frac{d\phi^*}{dz^*} = f^*(z, z^*)$$

The new source equations expand into a real and imaginary parts that are divergence and curl terms, respectively.

$$\begin{aligned} s^*(z, z^*) = 2 \frac{df^*}{dz} &= \left[ \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right] \left[ f_x^*(x, y) + if_y^*(x, y) \right] = \rho - i I, \quad \text{where: } f_x^* = f_x, \text{ and: } f_y^* = -f_y \\ &= \left[ \frac{\partial f_x^*}{\partial x} + \frac{\partial f_y^*}{\partial y} \right] + i \left[ \frac{\partial f_y^*}{\partial x} - \frac{\partial f_x^*}{\partial y} \right] = \left[ \nabla \cdot \mathbf{f}^* \right] + i \left[ \nabla \times \mathbf{f}^* \right]_z \end{aligned}$$

Real part: *Poisson scalar source equation (charge density  $\rho$ )*.      Imaginary part: *Biot-Savart vector source equation (current density  $I$ )*

$$\nabla \cdot \mathbf{f}^* = \rho$$

$$\nabla \times \mathbf{f}^* = -I$$

Field-vs-potential has Re and Im parts that are  $x$  and  $y$  components of grad  $\Phi$  and curl  $A_z$  from potential  $\phi = \Phi + iA$  or  $\phi^* = \Phi - iA$ .

$$\begin{aligned} f^*(z, z^*) = 2 \frac{d\phi^*}{dz^*} &= \left[ \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right] (\Phi - iA) = f_x^* + if_y^* \\ &= \left[ \frac{\partial \Phi}{\partial x} + i \frac{\partial \Phi}{\partial y} \right] + \left[ \frac{\partial A}{\partial y} - i \frac{\partial A}{\partial x} \right] = \left[ \nabla \Phi \right] + \left[ \nabla \times \mathbf{A}_z \right] \quad \text{(For source-free analytic functions these two fields are identical.)} \end{aligned}$$

Gradient of scalar potential is the *longitudinal field*  $\mathbf{f}_L^*$  and curl of a vector potential is the *transverse field*  $\mathbf{f}_T^*$ .


$$\text{Total field is: } \mathbf{f}^* = \mathbf{f}_L^* + \mathbf{f}_T^*$$

$$\mathbf{f}_L^* = \nabla \Phi$$


$$\mathbf{f}_T^* = \nabla \times \mathbf{A}$$

Potential, force, and source field equations vs. position, velocity, and acceleration equations

Field equations



Newton equations



Potential and source field theory reduced to sophomore mechanics of 1D-motion!

*Example 1* Consider a non-analytic field  $f(z) = (z^*)^2$  or  $f^*(z) = z^2$ .

Non-analytic source  $s^*$  is derivative of field  $f^*$

$$s^*(z, z^*) = 2 \frac{df^*}{dz} = 4z = 4x + i4y,$$

$$\text{or: } \rho = 4x, \quad \text{and: } I = -4y.$$

Non-analytic potential  $\phi$  is integral of field  $f^*$

$$\phi(z, z^*) = \frac{1}{2} \int f(z) dz = \frac{1}{2} \int (z^*)^2 dz = \frac{z(z^*)^2}{2} = \frac{(x+iy)(x^2-y^2-i2xy)}{2},$$

$$\text{or: } \Phi = \frac{x^3 + xy^2}{2}, \quad \text{and: } A = \frac{-y^3 - yx^2}{2}.$$



*Example 1* Consider a non-analytic field  $f(z) = (z^*)^2$  or  $f^*(z) = z^2$ .

Non-analytic source  $s^*$  is derivative of field  $f^*$

$$s^*(z, z^*) = 2 \frac{df^*}{dz} = 4z = 4x + i4y,$$

or:  $\rho = 4x$ , and:  $I = -4y$ .

Non-analytic potential  $\phi$  is integral of field  $f^*$

$$\phi(z, z^*) = \frac{1}{2} \int f(z) dz = \frac{1}{2} \int (z^*)^2 dz = \frac{z(z^*)^2}{2} = \frac{(x+iy)(x^2 - y^2 - i2xy)}{2},$$

or:  $\Phi = \frac{x^3 + xy^2}{2}$ , and:  $\mathbf{A} = \frac{-y^3 - yx^2}{2}$ .

The longitudinal field  $\mathbf{f}_L^*$  is quite different from the transverse field  $\mathbf{f}_T^*$

$$\mathbf{f}_L^* = \nabla \Phi = \nabla \left( \frac{x^3 + xy^2}{2} \right) = \begin{pmatrix} \frac{\partial \Phi}{\partial x} \\ \frac{\partial \Phi}{\partial y} \end{pmatrix} = \begin{pmatrix} \frac{3x^2 + y^2}{2} \\ xy \end{pmatrix}, \quad \mathbf{f}_T^* = \nabla \times \mathbf{A} = \nabla \times \left( \frac{-y^3 - yx^2}{2} \mathbf{e}_z \right) = \begin{pmatrix} \frac{\partial A}{\partial y} \\ -\frac{\partial A}{\partial x} \end{pmatrix} = \begin{pmatrix} \frac{-3y^2 - x^2}{2} \\ xy \end{pmatrix}.$$

*Example 1* Consider a non-analytic field  $f(z) = (z^*)^2$  or  $f^*(z) = z^2$ .

Non-analytic source  $s^*$  is derivative of field  $f^*$

$$s^*(z, z^*) = 2 \frac{df^*}{dz} = 4z = 4x + i4y,$$

or:  $\rho = 4x$ , and:  $I = -4y$ .

Non-analytic potential  $\phi$  is integral of field  $f^*$

$$\phi(z, z^*) = \frac{1}{2} \int f(z) dz = \frac{1}{2} \int (z^*)^2 dz = \frac{z(z^*)^2}{2} = \frac{(x+iy)(x^2-y^2-i2xy)}{2},$$

or:  $\Phi = \frac{x^3 + xy^2}{2}$ , and:  $A = \frac{-y^3 - yx^2}{2}$ .

The longitudinal field  $\mathbf{f}_L^*$  is quite different from the transverse field  $\mathbf{f}_T^*$

$$\mathbf{f}_L^* = \nabla \Phi = \nabla \left( \frac{x^3 + xy^2}{2} \right) = \begin{pmatrix} \frac{\partial \Phi}{\partial x} \\ \frac{\partial \Phi}{\partial y} \end{pmatrix} = \begin{pmatrix} \frac{3x^2 + y^2}{2} \\ xy \end{pmatrix}, \quad \mathbf{f}_T^* = \nabla \times \mathbf{A} = \nabla \times \left( \frac{-y^3 - yx^2}{2} \mathbf{e}_z \right) = \begin{pmatrix} \frac{\partial A}{\partial y} \\ -\frac{\partial A}{\partial x} \end{pmatrix} = \begin{pmatrix} \frac{-3y^2 - x^2}{2} \\ xy \end{pmatrix}.$$

Longitudinal field  $\mathbf{f}_L^*$  has no curl and the transverse field  $\mathbf{f}_T^*$  has no divergence. Sum field  $\mathbf{f}$  has both.

$$\mathbf{f}^* = \mathbf{f}_L^* + \mathbf{f}_T^* = \begin{pmatrix} \frac{3x^2 + y^2}{2} \\ xy \end{pmatrix} + \begin{pmatrix} \frac{-3y^2 - x^2}{2} \\ xy \end{pmatrix} = \begin{pmatrix} x^2 - y^2 \\ 2xy \end{pmatrix}, \quad \nabla \cdot \mathbf{f}^* = \nabla \cdot \mathbf{f}_L^* = 4x = \rho, \quad \nabla \times \mathbf{f}^* = \nabla \times \mathbf{f}_T^* = 4y = -I.$$

**Example 1** Consider a non-analytic field  $f(z) = (z^*)^2$  or  $f^*(z) = z^2$ .

Non-analytic source  $s^*$  is derivative of field  $f^*$

$$s^*(z, z^*) = 2 \frac{df^*}{dz} = 4z = 4x + i4y,$$

or:  $\rho = 4x$ , and:  $I = -4y$ .

Non-analytic potential  $\phi$  is integral of field  $f^*$

$$\phi(z, z^*) = \frac{1}{2} \int f(z) dz = \frac{1}{2} \int (z^*)^2 dz = \frac{z(z^*)^2}{2} = \frac{(x+iy)(x^2-y^2-i2xy)}{2},$$


or:  $\Phi = \frac{x^3 + xy^2}{2}$ , and:  $A = \frac{-y^3 - yx^2}{2}$ .

The longitudinal field  $\mathbf{f}_L^*$  is quite different from the transverse field  $\mathbf{f}_T^*$

$$\mathbf{f}_L^* = \nabla \Phi = \nabla \left( \frac{x^3 + xy^2}{2} \right) = \begin{pmatrix} \frac{\partial \Phi}{\partial x} \\ \frac{\partial \Phi}{\partial y} \end{pmatrix} = \begin{pmatrix} \frac{3x^2 + y^2}{2} \\ xy \end{pmatrix}, \quad \mathbf{f}_T^* = \nabla \times \mathbf{A} = \nabla \times \left( \frac{-y^3 - yx^2}{2} \mathbf{e}_z \right) = \begin{pmatrix} \frac{\partial A}{\partial y} \\ -\frac{\partial A}{\partial x} \end{pmatrix} = \begin{pmatrix} \frac{-3y^2 - x^2}{2} \\ xy \end{pmatrix}.$$



Longitudinal field  $\mathbf{f}_L^*$  has no curl and the transverse field  $\mathbf{f}_T^*$  has no divergence. Sum field  $\mathbf{f}$  has both.

$$\mathbf{f}^* = \mathbf{f}_L^* + \mathbf{f}_T^* = \begin{pmatrix} \frac{3x^2 + y^2}{2} \\ xy \end{pmatrix} + \begin{pmatrix} \frac{-3y^2 - x^2}{2} \\ xy \end{pmatrix} = \begin{pmatrix} x^2 - y^2 \\ 2xy \end{pmatrix}, \quad \nabla \cdot \mathbf{f}^* = \nabla \cdot \mathbf{f}_L^* = 4x = \rho, \quad \nabla \times \mathbf{f}^* = \nabla \times \mathbf{f}_T^* = 4y = -I.$$





$\mathbf{f}_L^* + \mathbf{f}_T^*$



Fig.10.17 Force field vectors for non-analytic function  $f(z) = (z^*)^2$

















$$Sum = \begin{pmatrix} x^2 - y^2 \\ 2xy \end{pmatrix}$$

