

# *Group Theory in Quantum Mechanics*

## *Lecture 18* (4.4.13)

### *Hexagonal $D_6 \subset D_{6h}$ and octahedral-tetrahedral $O \sim T_d$ symmetry*

(Int.J.Mol.Sci, 14, 714(2013) p.755-774 , QTCA Unit 5 Ch. 15 )  
(PSDS - Ch. 4 )

*Review: Symmetry reduction and splitting: Subduced irep  $D^\alpha(D_3) \downarrow C_2 = d^{0_2} \oplus d^{1_2} \oplus \dots$  correlation*  
*Symmetry induction and clustering: Induced rep  $d^a(C_2) \uparrow D_3 = D^\alpha \oplus D^\beta \oplus \dots$  correlation*

*$D_3$ - $C_2$  Coset structure of  $d^{m_2}(C_2) \uparrow D_3$  induced representation basis*

*$D_3$ -Projection of  $d^{m_2}(C_2) \uparrow D_3$  induced representation basis*

*Derivation of Frobenius reciprocity*

*$D_6 \supset D_2 \supset C_2 = D_3 \times C_2$  symmetry and outer product geometry*

*Irreducible characters*

*Irreducible representations*

*Correlations with  $D_6$  characters:*

*...and  $C_2(\mathbf{i}_3)$  characters.....and  $C_6(\mathbf{1}, \mathbf{h}^1, \mathbf{h}^2, \dots)$  characters*

*$D_6$  symmetry and induced representation band structure*

*Introduction to octahedral tetrahedral symmetry  $O_h \supset O \sim T_d \supset T$*

→ Review: *Symmetry reduction and splitting: Subduced irep  $D^\alpha(D_3) \downarrow C_2 = d^{0_2} \oplus d^{1_2} \oplus \dots$  correlation* ←  
*Symmetry induction and clustering: Induced rep  $d^a(C_2) \uparrow D_3 = D^\alpha \oplus D^\beta \oplus \dots$  correlation*

*$D_3$ - $C_2$  Coset structure of  $d^{m_2}(C_2) \uparrow D_3$  induced representation basis*

*$D_3$ -Projection of  $d^{m_2}(C_2) \uparrow D_3$  induced representation basis*

*Derivation of Frobenius reciprocity*

*$D_6 \supset D_2 \supset C_2 = D_3 \times C_2$  symmetry and outer product geometry*

*Irreducible characters*

*Irreducible representations*

*Correlations with  $D_6$  characters:*

*...and  $C_2(\mathbf{i}_3)$  characters.....and  $C_6(\mathbf{1}, \mathbf{h}^1, \mathbf{h}^2, \dots)$  characters*

*$D_6$  symmetry and induced representation band structure*

*Introduction to octahedral tetrahedral symmetry  $O_h \supset O \sim T_d \supset T$*

Applied symmetry reduction and splitting: Subduced irep  $D^\alpha(D_3) \downarrow C_2 = d^{0_2} \oplus d^{1_2} \oplus \dots$  correlation

$D_3 \supset C_2$	$\underline{\mathbf{P}^\alpha \text{ relabel/split}}$	$\underline{D^\alpha \text{ relabel/reduce}}$	$\underline{\omega^\alpha \text{ relabel/split}}$	$\underline{D_3 \supset C_2}$	$0_2$	$1_2$	$D^{A_1}(D_3) \downarrow C_2 \sim d^{0_2}$
$A_1$	$\mathbf{P}^{A_1} = \mathbf{P}^{A_1} \mathbf{P}^{0_2} = \mathbf{P}_{0_2 0_2}^{A_1}$	$\Rightarrow D^{A_1} \downarrow C_2 \sim d^{0_2}$	$\Rightarrow \omega^{A_1} \rightarrow \omega^{0_2}$	$A_1$	1	.	$D^{A_1}(D_3) \downarrow C_2 \sim d^{0_2}$
$A_2$	$\mathbf{P}^{A_2} = \mathbf{P}^{A_2} \mathbf{P}^{1_2} = \mathbf{P}_{1_2 1_2}^{A_2}$	$\Rightarrow D^{A_2} \downarrow C_2 \sim d^{1_2}$	$\Rightarrow \omega^{A_2} \rightarrow \omega^{1_2}$	$A_2$	.	1	$D^{A_2}(D_3) \downarrow C_2 \sim d^{1_2}$
$E_1$	$\mathbf{P}^{E_1} = \mathbf{P}^{E_1} \mathbf{P}^{0_2} + \mathbf{P}^{E_1} \mathbf{P}^{1_2}$ $= \mathbf{P}_{0_2 0_2}^{E_1} + \mathbf{P}_{1_2 1_2}^{E_1}$	$\Rightarrow D^{E_1} \downarrow C_2 \sim$ $d^{0_2} \oplus d^{1_2}$	$\Rightarrow \omega^{E_1} \rightarrow \omega^{0_2}$ $\searrow \omega^{1_2}$	$E_1$	1	1	$D^{E_1}(D_3) \downarrow C_2 \sim d^{0_2} \oplus d^{1_2}$

Applied symmetry reduction and splitting: Subduced irep  $D^\alpha(D_3) \downarrow C_2 = d^{0_2} \oplus d^{1_2} \oplus \dots$  correlation

$D_3 \supset C_2$	$\underline{\mathbf{P}^\alpha \text{ relabel/split}}$	$\underline{D^\alpha \text{ relabel/reduce}}$	$\underline{\omega^\alpha \text{ relabel/split}}$	$\underline{D_3 \supset C_2}$	$0_2$	$1_2$	
$A_1$	$\mathbf{P}^{A_1} = \mathbf{P}^{A_1} \mathbf{P}^{0_2} = \mathbf{P}_{0_2 0_2}^{A_1}$	$\Rightarrow D^{A_1} \downarrow C_2 \sim d^{0_2}$	$\Rightarrow \omega^{A_1} \rightarrow \omega^{0_2}$	$A_1$	1	.	$D^{A_1}(D_3) \downarrow C_2 \sim d^{0_2}$
$A_2$	$\mathbf{P}^{A_2} = \mathbf{P}^{A_2} \mathbf{P}^{1_2} = \mathbf{P}_{1_2 1_2}^{A_2}$	$\Rightarrow D^{A_2} \downarrow C_2 \sim d^{1_2}$	$\Rightarrow \omega^{A_2} \rightarrow \omega^{1_2}$	$A_2$	.	1	$D^{A_2}(D_3) \downarrow C_2 \sim d^{1_2}$
$E_1$	$\mathbf{P}^{E_1} = \mathbf{P}^{E_1} \mathbf{P}^{0_2} + \mathbf{P}^{E_1} \mathbf{P}^{1_2}$ $= \mathbf{P}_{0_2 0_2}^{E_1} + \mathbf{P}_{1_2 1_2}^{E_1}$	$\Rightarrow D^{E_1} \downarrow C_2 \sim d^{0_2} \oplus d^{1_2}$	$\Rightarrow \omega^{E_1} \rightarrow \omega^{0_2}$ $\searrow \omega^{1_2}$	$E_1$	1	1	$D^{E_1}(D_3) \downarrow C_2 \sim d^{0_2} \oplus d^{1_2}$
					$d^{0_2}(C_2) \uparrow D_3$		

Spontaneous symmetry breaking

and clustering: Induced rep  $d^a(C_2) \uparrow D_3 = D^\alpha \oplus D^\beta \oplus \dots$  correlation

$$\sim D^{A_1} \oplus D^{E_1}$$

$$d^{1_2}(C_2) \uparrow D_3$$

$$\sim D^{A_2} \oplus D^{E_1}$$

Applied symmetry reduction and splitting: Subduced irep  $D^\alpha(D_3) \downarrow C_2 = d^{0_2} \oplus d^{l_2} \oplus \dots$  correlation

$D_3 \supset C_2$	$\frac{\mathbf{P}^\alpha \text{ relabel/split}}{\mathbf{P}^{A_1} = \mathbf{P}^{A_1} \mathbf{P}^{0_2} = \mathbf{P}_{0_2 0_2}^{A_1}}$	$\frac{D^\alpha \text{ relabel/reduce}}{\Rightarrow D^{A_1} \downarrow C_2 \sim d^{0_2}}$	$\frac{\omega^\alpha \text{ relabel/split}}{\Rightarrow \omega^{A_1} \rightarrow \omega^{0_2}}$	$\frac{D_3 \supset C_2}{\begin{array}{c cc} & \mathbf{0}_2 & \mathbf{1}_2 \\ \hline A_1 & 1 & \cdot \\ A_2 & \cdot & 1 \\ E_1 & 1 & 1 \end{array}}$	$D^{A_1}(D_3) \downarrow C_2 \sim d^{0_2}$
$A_1$	$\mathbf{P}^{A_2} = \mathbf{P}^{A_2} \mathbf{P}^{1_2} = \mathbf{P}_{1_2 1_2}^{A_2}$	$\Rightarrow D^{A_2} \downarrow C_2 \sim d^{1_2}$	$\Rightarrow \omega^{A_2} \rightarrow \omega^{1_2}$		$D^{A_2}(D_3) \downarrow C_2 \sim d^{1_2}$
$E_1$	$\mathbf{P}^{E_1} = \mathbf{P}^{E_1} \mathbf{P}^{0_2} + \mathbf{P}^{E_1} \mathbf{P}^{1_2}$ $= \mathbf{P}_{0_2 0_2}^{E_1} + \mathbf{P}_{1_2 1_2}^{E_1}$	$\Rightarrow D^{E_1} \downarrow C_2 \sim d^{0_2} \oplus d^{1_2}$	$\Rightarrow \omega^{E_1} \rightarrow \omega^{0_2}$ $\searrow \omega^{1_2}$		$D^{E_1}(D_3) \downarrow C_2 \sim d^{0_2} \oplus d^{1_2}$
				$d^{0_2}(C_2) \uparrow D_3$ $\sim D^{A_1} \oplus D^{E_1}$	
				$d^{1_2}(C_2) \uparrow D_3$ $\sim D^{A_2} \oplus D^{E_1}$	

Spontaneous symmetry breaking

and clustering: Induced rep  $d^a(C_2) \uparrow D_3 = D^\alpha \oplus D^\beta \oplus \dots$  correlation

Applied symmetry reduction and splitting: Subduced irep  $D^\alpha(D_3) \downarrow C_3 = d^{0_3} \oplus d^{l_3} \oplus \dots$  correlation

$D_3 \supset C_3$	$\frac{\mathbf{P}^\alpha \text{ relabel/split}}{\mathbf{P}^{A_1} = \mathbf{P}^{A_1} \mathbf{P}^{0_3} = \mathbf{P}_{0_3 0_3}^{A_1}}$	$\frac{D^\alpha \text{ relabel/reduce}}{\Rightarrow D^{A_1} \downarrow C_3 \sim d^{0_3}}$	$\frac{\omega^\alpha \text{ relabel/split}}{\Rightarrow \omega^{A_1} \rightarrow \omega^{0_3}}$	$\frac{D_3 \supset C_3}{\begin{array}{c ccc} & \mathbf{0}_3 & \mathbf{1}_3 & \mathbf{2}_3 \\ \hline A_1 & 1 & \cdot & \cdot \\ A_2 & 1 & \cdot & \cdot \\ E_1 & \cdot & 1 & 1 \end{array}}$	$D^{A_1}(D_3) \downarrow C_3 \sim d^{0_3}$
$A_1$	$\mathbf{P}^{A_2} = \mathbf{P}^{A_2} \mathbf{P}^{0_3} = \mathbf{P}_{0_3 0_3}^{A_2}$	$\Rightarrow D^{A_2} \downarrow C_3 \sim d^{0_3}$	$\Rightarrow \omega^{A_2} \rightarrow \omega^{0_3}$		$D^{A_2}(D_3) \downarrow C_3 \sim d^{0_3}$
$E_1$	$\mathbf{P}^{E_1} = \mathbf{P}^{E_1} \mathbf{P}^{1_3} + \mathbf{P}^{E_1} \mathbf{P}^{2_3}$ $= \mathbf{P}_{1_3 1_3}^{E_1} + \mathbf{P}_{2_3 2_3}^{E_1}$	$\Rightarrow D^{E_1} \downarrow C_3 \sim d^{1_3} \oplus d^{2_3}$	$\Rightarrow \omega^{E_1} \rightarrow \omega^{1_3}$ $\searrow \omega^{2_3}$		$D^{E_1}(D_3) \downarrow C_3 \sim d^{1_3} \oplus d^{2_3}$

Applied symmetry reduction and splitting: Subduced irep  $D^\alpha(D_3) \downarrow C_2 = d^{0_2} \oplus d^{l_2} \oplus \dots$  correlation

$D_3 \supset C_2$	$\mathbf{P}^\alpha$ relabel/split	$D^\alpha$ relabel/reduce	$\omega^\alpha$ relabel/split	$D_3 \supset C_2$	$0_2$	$1_2$	$D^{\mathbf{A}_1}(D_3) \downarrow C_2 \sim d^{0_2}$
$A_1$	$\mathbf{P}^{\mathbf{A}_1} = \mathbf{P}^{\mathbf{A}_1} \mathbf{P}^{0_2} = \mathbf{P}_{0_2 0_2}^{\mathbf{A}_1}$	$\Rightarrow D^{\mathbf{A}_1} \downarrow C_2 \sim d^{0_2}$	$\Rightarrow \omega^{\mathbf{A}_1} \rightarrow \omega^{0_2}$	$A_1$	1	.	$D^{\mathbf{A}_1}(D_3) \downarrow C_2 \sim d^{0_2}$
$A_2$	$\mathbf{P}^{\mathbf{A}_2} = \mathbf{P}^{\mathbf{A}_2} \mathbf{P}^{1_2} = \mathbf{P}_{1_2 1_2}^{\mathbf{A}_2}$	$\Rightarrow D^{\mathbf{A}_2} \downarrow C_2 \sim d^{1_2}$	$\Rightarrow \omega^{\mathbf{A}_2} \rightarrow \omega^{1_2}$	$A_2$	.	1	$D^{\mathbf{A}_2}(D_3) \downarrow C_2 \sim d^{1_2}$
$E_1$	$\mathbf{P}^{\mathbf{E}_1} = \mathbf{P}^{\mathbf{E}_1} \mathbf{P}^{0_2} + \mathbf{P}^{\mathbf{E}_1} \mathbf{P}^{1_2}$ $= \mathbf{P}_{0_2 0_2}^{\mathbf{E}_1} + \mathbf{P}_{1_2 1_2}^{\mathbf{E}_1}$	$\Rightarrow D^{\mathbf{E}_1} \downarrow C_2 \sim d^{0_2} \oplus d^{1_2}$	$\Rightarrow \omega^{\mathbf{E}_1} \rightarrow \omega^{0_2}$ $\searrow \omega^{1_2}$	$E_1$	1	1	$D^{\mathbf{E}_1}(D_3) \downarrow C_2 \sim d^{0_2} \oplus d^{1_2}$
					$d^{0_2}(C_2) \uparrow D_3$		
					$\sim D^{\mathbf{A}_1} \oplus D^{\mathbf{E}_1}$		
					$d^{1_2}(C_2) \uparrow D_3$		
					$\sim D^{\mathbf{A}_2} \oplus D^{\mathbf{E}_1}$		

Spontaneous symmetry breaking

and clustering: Induced rep  $d^a(C_2) \uparrow D_3 = D^\alpha \oplus D^\beta \oplus \dots$  correlation

Applied symmetry reduction and splitting: Subduced irep  $D^\alpha(D_3) \downarrow C_3 = d^{0_3} \oplus d^{l_3} \oplus \dots$  correlation

$D_3 \supset C_3$	$\mathbf{P}^\alpha$ relabel/split	$D^\alpha$ relabel/reduce	$\omega^\alpha$ relabel/split	$D_3 \supset C_3$	$0_3$	$1_3$	$2_3$	$D^{\mathbf{A}_1}(D_3) \downarrow C_3 \sim d^{0_3}$
$A_1$	$\mathbf{P}^{\mathbf{A}_1} = \mathbf{P}^{\mathbf{A}_1} \mathbf{P}^{0_3} = \mathbf{P}_{0_3 0_3}^{\mathbf{A}_1}$	$\Rightarrow D^{\mathbf{A}_1} \downarrow C_3 \sim d^{0_3}$	$\Rightarrow \omega^{\mathbf{A}_1} \rightarrow \omega^{0_3}$	$A_1$	1	.	.	$D^{\mathbf{A}_1}(D_3) \downarrow C_3 \sim d^{0_3}$
$A_2$	$\mathbf{P}^{\mathbf{A}_2} = \mathbf{P}^{\mathbf{A}_2} \mathbf{P}^{0_3} = \mathbf{P}_{0_3 0_3}^{\mathbf{A}_2}$	$\Rightarrow D^{\mathbf{A}_2} \downarrow C_3 \sim d^{0_3}$	$\Rightarrow \omega^{\mathbf{A}_2} \rightarrow \omega^{0_3}$	$A_2$	1	.	.	$D^{\mathbf{A}_2}(D_3) \downarrow C_3 \sim d^{0_3}$
$E_1$	$\mathbf{P}^{\mathbf{E}_1} = \mathbf{P}^{\mathbf{E}_1} \mathbf{P}^{1_3} + \mathbf{P}^{\mathbf{E}_1} \mathbf{P}^{2_3}$ $= \mathbf{P}_{1_3 1_3}^{\mathbf{E}_1} + \mathbf{P}_{2_3 2_3}^{\mathbf{E}_1}$	$\Rightarrow D^{\mathbf{E}_1} \downarrow C_3 \sim d^{1_3} \oplus d^{2_3}$	$\Rightarrow \omega^{\mathbf{E}_1} \rightarrow \omega^{1_3}$ $\searrow \omega^{2_3}$	$E_1$	.	1	1	$D^{\mathbf{E}_1}(D_3) \downarrow C_3 \sim d^{1_3} \oplus d^{2_3}$
					$d^{0_3}(C_3) \uparrow D_3$			
					$\sim D^{\mathbf{A}_1} \oplus D^{\mathbf{A}_2}$			
					$d^{1_3}(C_3) \uparrow D_3$			
					$\sim D^{\mathbf{E}_1}$			
					$d^{2_3}(C_3) \uparrow D_3$			
					$\sim D^{\mathbf{E}_1}$			

Spontaneous symmetry breaking

and clustering: Induced rep  $d^a(C_3) \uparrow D_3 = D^\alpha \oplus D^\beta \oplus \dots$  correlation

Review: Symmetry reduction and splitting: Subduced irep  $D^\alpha(D_3) \downarrow C_2 = d^{0_2} \oplus d^{1_2} \oplus \dots$  correlation

Symmetry induction and clustering: Induced rep  $d^a(C_2) \uparrow D_3 = D^\alpha \oplus D^\beta \oplus \dots$  correlation

→  **$D_3$ -C<sub>2</sub> Coset structure of  $d^{m_2}(C_2) \uparrow D_3$  induced representation basis** ←  
 **$D_3$ -Projection of  $d^{m_2}(C_2) \uparrow D_3$  induced representation basis**  
**Derivation of Frobenius reciprocity**

$D_6 \supset D_2 \supset C_2 = D_3 \times C_2$  symmetry and outer product geometry

Irreducible characters

Irreducible representations

Correlations with  $D_6$  characters:


...and  $C_2(\mathbf{i}_3)$  characters.....and  $C_6(\mathbf{1}, \mathbf{h}^1, \mathbf{h}^2, \dots)$  characters

$D_6$  symmetry and induced representation band structure

Introduction to octahedral tetrahedral symmetry  $O_h \supset O \sim T_d \supset T$


# $D_3$ - $C_2$ Coset structure of $d^{m_2}(C_2) \uparrow D^3$ induced representation basis

Left cosets [ $\mathbf{1}C_2 = (\mathbf{1}, \mathbf{i}_3)$ ,  $\mathbf{r}^1 C_2 = (\mathbf{r}^1, \mathbf{i}_2)$ ,  $\mathbf{r}^2 C_2 = (\mathbf{r}^2, \mathbf{i}_1)$ ] relate to sets of  $\mathbf{r}^{\textcolor{teal}{P}}$ -transformed kets



# $D_3$ - $C_2$ Coset structure of $d^{m_2}(C_2) \uparrow D^3$ induced representation basis


Left cosets  $[1C_2 = (1, \mathbf{i}_3), \quad \mathbf{r}^1 C_2 = (\mathbf{r}^1, \mathbf{i}_2), \quad \mathbf{r}^2 C_2 = (\mathbf{r}^2, \mathbf{i}_1)]$  relate to sets of  $\mathbf{r}^{\textcolor{teal}{P}}$ -transformed kets  
 $[1(|1\rangle, |\mathbf{i}_3\rangle) = (|1\rangle, |\mathbf{i}_3\rangle), \quad \mathbf{r}^1(|1\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{r}^1\rangle, |\mathbf{i}_2\rangle), \quad \mathbf{r}^2(|1\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{r}^2\rangle, |\mathbf{i}_1\rangle)]$



# $D_3$ - $C_2$ Coset structure of $d^{m_2}(C_2) \uparrow D^3$ induced representation basis

Left cosets  $[1C_2 = (1, \mathbf{i}_3), \mathbf{r}^1 C_2 = (\mathbf{r}^1, \mathbf{i}_2), \mathbf{r}^2 C_2 = (\mathbf{r}^2, \mathbf{i}_1)]$  relate to sets of  $\mathbf{r}^{\textcolor{teal}{P}}$ -transformed kets  
 $[1(|1\rangle, |\mathbf{i}_3\rangle) = (|1\rangle, |\mathbf{i}_3\rangle), \mathbf{r}^1(|1\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{r}^1\rangle, |\mathbf{i}_2\rangle), \mathbf{r}^2(|1\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{r}^2\rangle, |\mathbf{i}_1\rangle)]$

Right cosets  $[C_2 = (1, \mathbf{i}_3), C_2 \mathbf{r}^2 = (\mathbf{r}^2, \mathbf{i}_2), C_2 \mathbf{r} = (\mathbf{r}, \mathbf{i}_1)]$  relate to sets of bras




# $D_3$ - $C_2$ Coset structure of $d^{m_2}(C_2) \uparrow D^3$ induced representation basis

Left cosets  $[1C_2 = (1, \mathbf{i}_3), \mathbf{r}^1 C_2 = (\mathbf{r}^1, \mathbf{i}_2), \mathbf{r}^2 C_2 = (\mathbf{r}^2, \mathbf{i}_1)]$  relate to sets of  $\mathbf{r}^{\textcolor{teal}{P}}$ -transformed kets  
 $[(1|1\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{1}\rangle, |\mathbf{i}_3\rangle), \mathbf{r}^1(1|1\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{r}^1\rangle, |\mathbf{i}_2\rangle), \mathbf{r}^2(1|1\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{r}^2\rangle, |\mathbf{i}_1\rangle)]$

Right cosets  $[C_2 = (1, \mathbf{i}_3), C_2 \mathbf{r}^2 = (\mathbf{r}^2, \mathbf{i}_2), C_2 \mathbf{r} = (\mathbf{r}, \mathbf{i}_1)]$  relate to sets of bras

$[(\langle 1|, \langle \mathbf{i}_3|)1 = (\langle 1|, \langle \mathbf{i}_3|), (\langle 1|, \langle \mathbf{i}_3|)\mathbf{r}^2 = (\langle \mathbf{r}^1|, \langle \mathbf{i}_2|), (\langle 1|, \langle \mathbf{i}_3|)\mathbf{r}^1 = (\langle \mathbf{r}^2|, \langle \mathbf{i}_1|)]$




## D<sub>3</sub>-C<sub>2</sub> Coset structure of $d^{m_2}(C_2) \uparrow D^3$ induced representation basis

*Left cosets* [ $\mathbf{1}C_2 = (\mathbf{1}, \mathbf{i}_3)$ ,  $\mathbf{r}^1 C_2 = (\mathbf{r}^1, \mathbf{i}_2)$ ,  $\mathbf{r}^2 C_2 = (\mathbf{r}^2, \mathbf{i}_1)$ ] relate to sets of  $\mathbf{r}^{\text{teal}}$ -transformed kets  
 $[\mathbf{1}(|\mathbf{1}\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{1}\rangle, |\mathbf{i}_3\rangle), \mathbf{r}^1(|\mathbf{1}\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{r}^1\rangle, |\mathbf{i}_2\rangle), \mathbf{r}^2(|\mathbf{1}\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{r}^2\rangle, |\mathbf{i}_1\rangle)]$

*Right cosets* [ $C_2 = (\mathbf{1}, \mathbf{i}_3)$ ,  $C_2 \mathbf{r}^2 = (\mathbf{r}^2, \mathbf{i}_2)$ ,  $C_2 \mathbf{r} = (\mathbf{r}, \mathbf{i}_1)$ ] relate to sets of bras

$[(\langle \mathbf{1}|, \langle \mathbf{i}_3|) \mathbf{1} = (\langle \mathbf{1}|, \langle \mathbf{i}_3|), (\langle \mathbf{1}|, \langle \mathbf{i}_3|) \mathbf{r}^2 = (\langle \mathbf{r}^1|, \langle \mathbf{i}_2|), (\langle \mathbf{1}|, \langle \mathbf{i}_3|) \mathbf{r}^1 = (\langle \mathbf{r}^2|, \langle \mathbf{i}_1|)]$

C<sub>2</sub> projectors  $\mathbf{P}^{0_2} = \frac{1}{2}(\mathbf{1} + \mathbf{i}_3) = \mathbf{P}^x$  and  $\mathbf{P}^{1_2} = \frac{1}{2}(\mathbf{1} - \mathbf{i}_3) = \mathbf{P}^y$  split ket  $|\mathbf{r}\rangle = \mathbf{r}|\mathbf{1}\rangle$  or bra  $\langle \mathbf{r}| = \langle \mathbf{1}|\mathbf{r}^\dagger$  into  $\pm$  coset sums



# $D_3$ - $C_2$ Coset structure of $d^{m_2}(C_2) \uparrow D^3$ induced representation basis


Left cosets  $[1C_2 = (1, \mathbf{i}_3), \mathbf{r}^1 C_2 = (\mathbf{r}^1, \mathbf{i}_2), \mathbf{r}^2 C_2 = (\mathbf{r}^2, \mathbf{i}_1)]$  relate to sets of  $\mathbf{r}^{\text{teal}}$ -transformed kets  
 $[1(|1\rangle, |\mathbf{i}_3\rangle) = (|1\rangle, |\mathbf{i}_3\rangle), \mathbf{r}^1(|1\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{r}^1\rangle, |\mathbf{i}_2\rangle), \mathbf{r}^2(|1\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{r}^2\rangle, |\mathbf{i}_1\rangle)]$

Right cosets  $[C_2 = (1, \mathbf{i}_3), C_2 \mathbf{r}^2 = (\mathbf{r}^2, \mathbf{i}_2), C_2 \mathbf{r} = (\mathbf{r}, \mathbf{i}_1)]$  relate to sets of bras

$[(\langle 1 |, \langle \mathbf{i}_3 |) \mathbf{1} = (\langle 1 |, \langle \mathbf{i}_3 |), (\langle 1 |, \langle \mathbf{i}_3 |) \mathbf{r}^2 = (\langle \mathbf{r}^1 |, \langle \mathbf{i}_2 |), (\langle 1 |, \langle \mathbf{i}_3 |) \mathbf{r}^1 = (\langle \mathbf{r}^2 |, \langle \mathbf{i}_1 |)]$

$C_2$  projectors  $\mathbf{P}^{0_2} = \frac{1}{2}(1 + \mathbf{i}_3) = \mathbf{P}^x$  and  $\mathbf{P}^{1_2} = \frac{1}{2}(1 - \mathbf{i}_3) = \mathbf{P}^y$  split ket  $|\mathbf{r}\rangle = \mathbf{r}|1\rangle$  or bra  $\langle \mathbf{r}| = \langle 1|\mathbf{r}^\dagger$  into  $\pm$  coset sums

$$\left[ \mathbf{P}^{n_2} |1\rangle = \frac{1}{2} (|1\rangle \pm |\mathbf{i}_3\rangle), \quad \right] = \left[ |\mathbf{r}_n^0\rangle \quad , \quad \right] \text{basis of } d^{n_2} \uparrow D_3$$




# $D_3$ - $C_2$ Coset structure of $d^{m_2}(C_2) \uparrow D^3$ induced representation basis

Left cosets  $[1C_2 = (1, \mathbf{i}_3), \mathbf{r}^1 C_2 = (\mathbf{r}^1, \mathbf{i}_2), \mathbf{r}^2 C_2 = (\mathbf{r}^2, \mathbf{i}_1)]$  relate to sets of  $\mathbf{r}^{\text{teal}}$ -transformed kets  
 $[1(|1\rangle, |\mathbf{i}_3\rangle) = (|1\rangle, |\mathbf{i}_3\rangle), \mathbf{r}^1(|1\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{r}^1\rangle, |\mathbf{i}_2\rangle), \mathbf{r}^2(|1\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{r}^2\rangle, |\mathbf{i}_1\rangle)]$

Right cosets  $[C_2 = (1, \mathbf{i}_3), C_2 \mathbf{r}^2 = (\mathbf{r}^2, \mathbf{i}_2), C_2 \mathbf{r} = (\mathbf{r}, \mathbf{i}_1)]$  relate to sets of bras

$$[(\langle 1 |, \langle \mathbf{i}_3 |) \mathbf{1} = (\langle 1 |, \langle \mathbf{i}_3 |), (\langle 1 |, \langle \mathbf{i}_3 |) \mathbf{r}^2 = (\langle \mathbf{r}^1 |, \langle \mathbf{i}_2 |), (\langle 1 |, \langle \mathbf{i}_3 |) \mathbf{r}^1 = (\langle \mathbf{r}^2 |, \langle \mathbf{i}_1 |)]$$



$C_2$  projectors  $\mathbf{P}^{0_2} = \frac{1}{2}(1 + \mathbf{i}_3) = \mathbf{P}^x$  and  $\mathbf{P}^{1_2} = \frac{1}{2}(1 - \mathbf{i}_3) = \mathbf{P}^y$  split ket  $|\mathbf{r}\rangle = \mathbf{r}|1\rangle$  or bra  $\langle \mathbf{r}| = \langle 1|\mathbf{r}^\dagger$  into  $\pm$  coset sums

$$\left[ \mathbf{P}^{n_2} |1\rangle = \frac{1}{2} (|1\rangle \pm |\mathbf{i}_3\rangle), \quad \right] = \left[ |\mathbf{r}_n^0\rangle \quad , \quad \right] \text{basis of } d^{n_2} \uparrow D_3$$

$$\left[ \langle 1| \mathbf{P}^{n_2} = \frac{1}{2} (\langle 1| \pm \langle \mathbf{i}_3|), \quad \right] = \left[ \langle \mathbf{r}_n^0 | \quad , \quad \right] \text{basis of } d^{n_2} \uparrow D_3$$

Review: Symmetry reduction and splitting: Subduced irep  $D^\alpha(D_3) \downarrow C_2 = d^{0_2} \oplus d^{1_2} \oplus \dots$  correlation

Symmetry induction and clustering: Induced rep  $d^a(C_2) \uparrow D_3 = D^\alpha \oplus D^\beta \oplus \dots$  correlation

*$D_3$ - $C_2$  Coset structure of  $d^{m_2}(C_2) \uparrow D_3$  induced representation basis*

$\rightarrow$   *$D_3$ -Projection of  $d^{m_2}(C_2) \uparrow D_3$  induced representation basis*

*Derivation of Frobenius reciprocity*



$D_6 \supset D_2 \supset C_2 = D_3 \times C_2$  symmetry and outer product geometry

*Irreducible characters*

*Irreducible representations*

*Correlations with  $D_6$  characters:*

*...and  $C_2(\mathbf{i}_3)$  characters.....and  $C_6(\mathbf{1}, \mathbf{h}^1, \mathbf{h}^2, \dots)$  characters*

*$D_6$  symmetry and induced representation band structure*


*Introduction to octahedral tetrahedral symmetry  $O_h \supset O \sim T_d \supset T$*

# $D_3$ - $C_2$ Coset structure of $d^{m_2}(C_2) \uparrow D^3$ induced representation basis

Left cosets [ $1C_2 = (1, \mathbf{i}_3)$ ,  $\mathbf{r}^1 C_2 = (\mathbf{r}^1, \mathbf{i}_2)$ ,  $\mathbf{r}^2 C_2 = (\mathbf{r}^2, \mathbf{i}_1)$ ] relate to sets of  $\mathbf{r}^{\text{p}}$ -transformed kets  
 $[|1\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{1}\rangle, |\mathbf{i}_3\rangle), \quad \mathbf{r}^1(|\mathbf{1}\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{r}^1\rangle, |\mathbf{i}_2\rangle), \quad \mathbf{r}^2(|\mathbf{1}\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{r}^2\rangle, |\mathbf{i}_1\rangle)]$

Right cosets [ $C_2 = (1, \mathbf{i}_3)$ ,  $C_2 \mathbf{r}^2 = (\mathbf{r}^2, \mathbf{i}_2)$ ,  $C_2 \mathbf{r} = (\mathbf{r}, \mathbf{i}_1)$ ] relate to sets of bras


$$[(\langle \mathbf{1}|, \langle \mathbf{i}_3|) \mathbf{1} = (\langle \mathbf{1}|, \langle \mathbf{i}_3|), \quad (\langle \mathbf{1}|, \langle \mathbf{i}_3|) \mathbf{r}^2 = (\langle \mathbf{r}^1|, \langle \mathbf{i}_2|), \quad (\langle \mathbf{1}|, \langle \mathbf{i}_3|) \mathbf{r}^1 = (\langle \mathbf{r}^2|, \langle \mathbf{i}_1|)]$$



$C_2$  projectors  $\mathbf{P}^{0_2} = \frac{1}{2}(1 + \mathbf{i}_3) = \mathbf{P}^x$  and  $\mathbf{P}^{1_2} = \frac{1}{2}(1 - \mathbf{i}_3) = \mathbf{P}^y$  split ket  $|\mathbf{r}\rangle = \mathbf{r}|1\rangle$  or bra  $\langle \mathbf{r}| = \langle 1|\mathbf{r}^\dagger$  into  $\pm$  coset sums

$$\left[ \mathbf{P}^{n_2} |1\rangle = \frac{1}{2} (|1\rangle \pm |\mathbf{i}_3\rangle), \quad \right] = \left[ |\mathbf{r}_n^0\rangle \quad , \quad \right] \text{basis of } d^{n_2} \uparrow D_3$$

$$\left[ \langle 1| \mathbf{P}^{n_2} = \frac{1}{2} (\langle 1| \pm \langle \mathbf{i}_3|), \quad \right] = \left[ \langle \mathbf{r}_n^0 | \quad , \quad \right] \text{basis of } d^{n_2} \uparrow D_3$$




# $D_3$ - $C_2$ Coset structure of $d^{m_2}(C_2) \uparrow D^3$ induced representation basis

Left cosets [ $1C_2 = (1, \mathbf{i}_3)$ ,  $\mathbf{r}^1 C_2 = (\mathbf{r}^1, \mathbf{i}_2)$ ,  $\mathbf{r}^2 C_2 = (\mathbf{r}^2, \mathbf{i}_1)$ ] relate to sets of  $\mathbf{r}^{\text{p}}$ -transformed kets  
 $[|1\rangle, |\mathbf{i}_3\rangle] = (|1\rangle, |\mathbf{i}_3\rangle)$ ,  $\mathbf{r}^1(|1\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{r}^1\rangle, |\mathbf{i}_2\rangle)$ ,  $\mathbf{r}^2(|1\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{r}^2\rangle, |\mathbf{i}_1\rangle)]$

Right cosets [ $C_2 = (1, \mathbf{i}_3)$ ,  $C_2 \mathbf{r}^2 = (\mathbf{r}^2, \mathbf{i}_2)$ ,  $C_2 \mathbf{r} = (\mathbf{r}, \mathbf{i}_1)$ ] relate to sets of bras



$$[(\langle 1 |, \langle \mathbf{i}_3 |) \mathbf{1} = (\langle 1 |, \langle \mathbf{i}_3 |), \quad (\langle 1 |, \langle \mathbf{i}_3 |) \mathbf{r}^2 = (\langle \mathbf{r}^1 |, \langle \mathbf{i}_2 |), \quad (\langle 1 |, \langle \mathbf{i}_3 |) \mathbf{r}^1 = (\langle \mathbf{r}^2 |, \langle \mathbf{i}_1 |)]$$



$C_2$  projectors  $\mathbf{P}^{0_2} = \frac{1}{2}(1 + \mathbf{i}_3) = \mathbf{P}^x$  and  $\mathbf{P}^{1_2} = \frac{1}{2}(1 - \mathbf{i}_3) = \mathbf{P}^y$  split ket  $|\mathbf{r}\rangle = \mathbf{r}|1\rangle$  or bra  $\langle \mathbf{r}| = \langle 1|\mathbf{r}^\dagger$  into  $\pm$  coset sums

$$\left[ \mathbf{P}^{n_2} |1\rangle = \frac{1}{2} (|1\rangle \pm |\mathbf{i}_3\rangle), \quad \right] = \left[ |\mathbf{r}_n^0\rangle \right], \quad \text{basis of } d^{n_2} \uparrow D_3$$

$$\left[ \langle 1| \mathbf{P}^{n_2} = \frac{1}{2} (\langle 1| \pm \langle \mathbf{i}_3|), \quad \right] = \left[ \langle \mathbf{r}_n^0 | \right], \quad \text{basis of } d^{n_2} \uparrow D_3$$




# $D_3$ - $C_2$ Coset structure of $d^{m_2}(C_2) \uparrow D^3$ induced representation basis

Left cosets [ $1C_2 = (1, \mathbf{i}_3)$ ,  $\mathbf{r}^1 C_2 = (\mathbf{r}^1, \mathbf{i}_2)$ ,  $\mathbf{r}^2 C_2 = (\mathbf{r}^2, \mathbf{i}_1)$ ] relate to sets of  $\mathbf{r}^{\text{p}}$ -transformed kets  
 $[|1\rangle, |\mathbf{i}_3\rangle] = (|1\rangle, |\mathbf{i}_3\rangle)$ ,  $\mathbf{r}^1(|1\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{r}^1\rangle, |\mathbf{i}_2\rangle)$ ,  $\mathbf{r}^2(|1\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{r}^2\rangle, |\mathbf{i}_1\rangle)]$

Right cosets [ $C_2 = (1, \mathbf{i}_3)$ ,  $C_2 \mathbf{r}^2 = (\mathbf{r}^2, \mathbf{i}_2)$ ,  $C_2 \mathbf{r} = (\mathbf{r}, \mathbf{i}_1)$ ] relate to sets of bras



$$[(\langle 1 |, \langle \mathbf{i}_3 |) \mathbf{1} = (\langle 1 |, \langle \mathbf{i}_3 |), \quad (\langle 1 |, \langle \mathbf{i}_3 |) \mathbf{r}^2 = (\langle \mathbf{r}^1 |, \langle \mathbf{i}_2 |), \quad (\langle 1 |, \langle \mathbf{i}_3 |) \mathbf{r}^1 = (\langle \mathbf{r}^2 |, \langle \mathbf{i}_1 |)]$$



$C_2$  projectors  $\mathbf{P}^{0_2} = \frac{1}{2}(1 + \mathbf{i}_3) = \mathbf{P}^x$  and  $\mathbf{P}^{1_2} = \frac{1}{2}(1 - \mathbf{i}_3) = \mathbf{P}^y$  split ket  $|\mathbf{r}\rangle = \mathbf{r}|1\rangle$  or bra  $\langle \mathbf{r}| = \langle 1|\mathbf{r}^\dagger$  into  $\pm$  coset sums

$$\left[ , \mathbf{P}^{n_2} |\mathbf{r}^1\rangle = \frac{1}{2} (|\mathbf{r}^1\rangle \pm |\mathbf{i}_2\rangle), \quad \right] = \left[ , |\mathbf{r}_n^1\rangle, \right] \text{basis of } d^{n_2} \uparrow D_3$$

$$\left[ , \langle \mathbf{r}^1 | \mathbf{P}^{n_2} = \frac{1}{2} (\langle \mathbf{r}^1 | \pm \langle \mathbf{i}_2 |), \quad \right] = \left[ , \langle \mathbf{r}_n^1 |, \right] \text{basis of } d^{n_2} \uparrow D_3$$




# $D_3$ - $C_2$ Coset structure of $d^{m_2}(C_2) \uparrow D^3$ induced representation basis

Left cosets [ $1C_2 = (1, \mathbf{i}_3)$ ,  $\mathbf{r}^1 C_2 = (\mathbf{r}^1, \mathbf{i}_2)$ ,  $\mathbf{r}^2 C_2 = (\mathbf{r}^2, \mathbf{i}_1)$ ] relate to sets of  $\mathbf{r}$ -transformed kets  
 $[|1\rangle, |\mathbf{i}_3\rangle] = (|1\rangle, |\mathbf{i}_3\rangle)$ ,  $\mathbf{r}^1(|1\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{r}^1\rangle, |\mathbf{i}_2\rangle)$ ,  $\mathbf{r}^2(|1\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{r}^2\rangle, |\mathbf{i}_1\rangle)]$

Right cosets [ $C_2 = (1, \mathbf{i}_3)$ ,  $C_2 \mathbf{r}^2 = (\mathbf{r}^2, \mathbf{i}_2)$ ,  $C_2 \mathbf{r} = (\mathbf{r}, \mathbf{i}_1)$ ] relate to sets of bras



$$[(\langle 1 |, \langle \mathbf{i}_3 |) \mathbf{1} = (\langle 1 |, \langle \mathbf{i}_3 |), \quad (\langle 1 |, \langle \mathbf{i}_3 |) \mathbf{r}^2 = (\langle \mathbf{r}^1 |, \langle \mathbf{i}_2 |), \quad (\langle 1 |, \langle \mathbf{i}_3 |) \mathbf{r}^1 = (\langle \mathbf{r}^2 |, \langle \mathbf{i}_1 |)]$$



$C_2$  projectors  $\mathbf{P}^{0_2} = \frac{1}{2}(1 + \mathbf{i}_3) = \mathbf{P}^x$  and  $\mathbf{P}^{1_2} = \frac{1}{2}(1 - \mathbf{i}_3) = \mathbf{P}^y$  split ket  $|\mathbf{r}\rangle = \mathbf{r}|1\rangle$  or bra  $\langle \mathbf{r}| = \langle 1|\mathbf{r}^\dagger$  into  $\pm$  coset sums

$$\left[ \dots, \mathbf{P}^{n_2} |\mathbf{r}^2\rangle = \frac{1}{2} (|\mathbf{r}^2\rangle \pm |\mathbf{i}_1\rangle) \right] = \left[ \dots, |\mathbf{r}_n^2\rangle \right] \text{basis of } d^{n_2} \uparrow D_3$$

$$\left[ \dots, \langle \mathbf{r}^2 | \mathbf{P}^{n_2} = \frac{1}{2} (\langle \mathbf{r}^2 | \pm \langle \mathbf{i}_1 |) \right] = \left[ \dots, \langle \mathbf{r}_n^2 | \right] \text{basis of } d^{n_2} \uparrow D_3$$




# $D_3$ - $C_2$ Coset structure of $d^{m_2}(C_2) \uparrow D^3$ induced representation basis

Left cosets [ $1C_2 = (1, \mathbf{i}_3)$ ,  $\mathbf{r}^1 C_2 = (\mathbf{r}^1, \mathbf{i}_2)$ ,  $\mathbf{r}^2 C_2 = (\mathbf{r}^2, \mathbf{i}_1)$ ] relate to sets of  $\mathbf{r}^{\text{p}}$ -transformed kets

$$[|1\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{1}\rangle, |\mathbf{i}_3\rangle), \quad \mathbf{r}^1(|1\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{r}^1\rangle, |\mathbf{i}_2\rangle), \quad \mathbf{r}^2(|1\rangle, |\mathbf{i}_3\rangle) = (|\mathbf{r}^2\rangle, |\mathbf{i}_1\rangle)]$$

Right cosets [ $C_2 = (1, \mathbf{i}_3)$ ,  $C_2 \mathbf{r}^2 = (\mathbf{r}^2, \mathbf{i}_2)$ ,  $C_2 \mathbf{r} = (\mathbf{r}, \mathbf{i}_1)$ ] relate to sets of bras



$$[(\langle 1 |, \langle \mathbf{i}_3 |) \mathbf{1} = (\langle 1 |, \langle \mathbf{i}_3 |), \quad (\langle 1 |, \langle \mathbf{i}_3 |) \mathbf{r}^2 = (\langle \mathbf{r}^1 |, \langle \mathbf{i}_2 |), \quad (\langle 1 |, \langle \mathbf{i}_3 |) \mathbf{r}^1 = (\langle \mathbf{r}^2 |, \langle \mathbf{i}_1 |)]$$



$C_2$  projectors  $\mathbf{P}^{0_2} = \frac{1}{2}(1 + \mathbf{i}_3) = \mathbf{P}^x$  and  $\mathbf{P}^{1_2} = \frac{1}{2}(1 - \mathbf{i}_3) = \mathbf{P}^y$  split ket  $|\mathbf{r}\rangle = \mathbf{r}|1\rangle$  or bra  $\langle \mathbf{r}| = \langle 1|\mathbf{r}^\dagger$  into  $\pm$  coset sums

$$\left[ \mathbf{P}^{n_2} |1\rangle = \frac{1}{2} (|1\rangle \pm |\mathbf{i}_3\rangle), \quad \mathbf{P}^{n_2} |\mathbf{r}^1\rangle = \frac{1}{2} (|\mathbf{r}^1\rangle \pm |\mathbf{i}_2\rangle), \quad \mathbf{P}^{n_2} |\mathbf{r}^2\rangle = \frac{1}{2} (|\mathbf{r}^2\rangle \pm |\mathbf{i}_1\rangle) \right] = \left[ |\mathbf{r}_n^0\rangle, |\mathbf{r}_n^1\rangle, |\mathbf{r}_n^2\rangle \right] \text{basis of } d^{n_2} \uparrow D_3$$

$$\left[ \langle 1 | \mathbf{P}^{n_2} = \frac{1}{2} (\langle 1 | \pm \langle \mathbf{i}_3 |), \quad \langle \mathbf{r}^1 | \mathbf{P}^{n_2} = \frac{1}{2} (\langle \mathbf{r}^1 | \pm \langle \mathbf{i}_2 |), \quad \langle \mathbf{r}^2 | \mathbf{P}^{n_2} = \frac{1}{2} (\langle \mathbf{r}^2 | \pm \langle \mathbf{i}_1 |) \right] = \left[ \langle \mathbf{r}_n^0 |, \langle \mathbf{r}_n^1 |, \langle \mathbf{r}_n^2 | \right] \text{basis of } d^{n_2} \uparrow D_3$$



Review: Symmetry reduction and splitting: Subduced irep  $D^\alpha(D_3) \downarrow C_2 = d^{0_2} \oplus d^{1_2} \oplus \dots$  correlation

Symmetry induction and clustering: Induced rep  $d^a(C_2) \uparrow D_3 = D^\alpha \oplus D^\beta \oplus \dots$  correlation

*D<sub>3</sub>-C<sub>2</sub> Coset structure of d<sup>m<sub>2</sub></sup>(C<sub>2</sub>)↑D<sub>3</sub> induced representation basis*

*D<sub>3</sub>-Projection of d<sup>m<sub>2</sub></sup>(C<sub>2</sub>)↑D<sub>3</sub> induced representation basis*

→ *Derivation of Frobenius reciprocity*



*D<sub>6</sub> ⊃ D<sub>2</sub> ⊃ C<sub>2</sub> = D<sub>3</sub> × C<sub>2</sub> symmetry and outer product geometry*

*Irreducible characters*

*Irreducible representations*

*Correlations with D<sub>6</sub> characters:*


*...and C<sub>2</sub>(i<sub>3</sub>) characters.....and C<sub>6</sub>(1, h<sup>1</sup>, h<sup>2</sup>, ...) characters*

*D<sub>6</sub> symmetry and induced representation band structure*

*Introduction to octahedral tetrahedral symmetry O<sub>h</sub> ⊃ O ~ T<sub>d</sub> ⊃ T*

## *D<sub>3</sub>-Projection of d<sup>m<sub>2</sub></sup>(C<sub>2</sub>)↑D<sup>3</sup> induced representation basis*


D<sub>3</sub>⊇C<sub>2</sub> projectors  $\mathbf{P}_{0_20_2}^{A_1}$ ,  $\mathbf{P}_{1_21_2}^{A_2}$ ,  $\mathbf{P}_{0_20_2}^{E_1}$ ,  $\mathbf{P}_{0_21_2}^{E_1}$ ,  $\mathbf{P}_{1_20_2}^{E_1}$ ,  $\mathbf{P}_{1_21_2}^{E_1}$  must reduce induced representation  $d^{m_2}(C_2) \uparrow D_3$



## *D<sub>3</sub>-Projection of d<sup>m<sub>2</sub></sup>(C<sub>2</sub>)↑D<sup>3</sup> induced representation basis*

D<sub>3</sub>⊇C<sub>2</sub> projectors  $\mathbf{P}_{0_20_2}^{A_1}$ ,  $\mathbf{P}_{1_21_2}^{A_2}$ ,  $\mathbf{P}_{0_20_2}^{E_1}$ ,  $\mathbf{P}_{0_21_2}^{E_1}$ ,  $\mathbf{P}_{1_20_2}^{E_1}$ ,  $\mathbf{P}_{1_21_2}^{E_1}$  must reduce induced representation  $d^{m_2}(C_2) \uparrow D_3$

But, which D<sub>3</sub> projector  $\mathbf{P}_{j_2k_2}^{\mu}$  will work on base  $|\mathbf{r}_x^0\rangle = \mathbf{p}^{m_2}|1\rangle$  of induced representation  $d^{m_2}(C_2) \uparrow D_3$




## *D<sub>3</sub>-Projection of d<sup>m<sub>2</sub></sup>(C<sub>2</sub>)↑D<sup>3</sup> induced representation basis*

D<sub>3</sub>⊇C<sub>2</sub> projectors  $\mathbf{P}_{0_20_2}^{A_1}$ ,  $\mathbf{P}_{1_21_2}^{A_2}$ ,  $\mathbf{P}_{0_20_2}^{E_1}$ ,  $\mathbf{P}_{0_21_2}^{E_1}$ ,  $\mathbf{P}_{1_20_2}^{E_1}$ ,  $\mathbf{P}_{1_21_2}^{E_1}$  must reduce induced representation  $d^{m_2}(C_2) \uparrow D_3$

But, which D<sub>3</sub> projector  $\mathbf{P}_{j_2k_2}^{\mu}$  will work on base  $|\mathbf{r}_{m_2}^0\rangle = \mathbf{p}^{m_2}|1\rangle$  of induced representation  $d^{m_2}(C_2) \uparrow D_3$

$$\mathbf{P}_{j_2k_2}^{\mu} |\mathbf{r}_{m_2}^0\rangle = \mathbf{P}_{j_2k_2}^{\mu} \mathbf{p}^{m_2} |1\rangle = ?$$




## *D<sub>3</sub>-Projection of d<sup>m<sub>2</sub></sup>(C<sub>2</sub>)↑D<sup>3</sup> induced representation basis*

D<sub>3</sub>⊇C<sub>2</sub> projectors  $\mathbf{P}_{0_20_2}^{A_1}$ ,  $\mathbf{P}_{1_21_2}^{A_2}$ ,  $\mathbf{P}_{0_20_2}^{E_1}$ ,  $\mathbf{P}_{0_21_2}^{E_1}$ ,  $\mathbf{P}_{1_20_2}^{E_1}$ ,  $\mathbf{P}_{1_21_2}^{E_1}$  must reduce induced representation  $d^{m_2}(C_2) \uparrow D_3$

But, which D<sub>3</sub> projector  $\mathbf{P}_{j_2k_2}^{\mu}$  will work on base  $|\mathbf{r}_{m_2}^0\rangle = \mathbf{p}^{m_2}|1\rangle$  of induced representation  $d^{m_2}(C_2) \uparrow D_3$

$$\mathbf{P}_{j_2k_2}^{\mu} |\mathbf{r}_{m_2}^0\rangle = \mathbf{P}_{j_2k_2}^{\mu} \mathbf{p}^{m_2} |1\rangle = \delta_{k_2}^{m_2} \mathbf{P}_{j_2m_2}^{\mu} |1\rangle$$

Local symmetry  $k_2$  of  $\mathbf{P}_{j_2k_2}^{\mu}$  must match that of  $|\mathbf{r}_{m_2}^0\rangle$



## D<sub>3</sub>-Projection of $d^{m_2}(C_2) \uparrow D^3$ induced representation basis


$D_3 \supset C_2$  projectors  $\mathbf{P}_{0_2 0_2}^{A_1}$ ,  $\mathbf{P}_{1_2 1_2}^{A_2}$ ,  $\mathbf{P}_{0_2 0_2}^{E_1}$ ,  $\mathbf{P}_{0_2 1_2}^{E_1}$ ,  $\mathbf{P}_{1_2 0_2}^{E_1}$ ,  $\mathbf{P}_{1_2 1_2}^{E_1}$  must reduce induced representation  $d^{m_2}(C_2) \uparrow D_3$

But, which  $D_3$  projector  $\mathbf{P}_{j_2 k_2}^\mu$  will work on base  $|\mathbf{r}_{m_2}^0\rangle = \mathbf{p}^{m_2}|1\rangle$  of induced representation  $d^{m_2}(C_2) \uparrow D_3$

$$\mathbf{P}_{j_2 k_2}^\mu |\mathbf{r}_{m_2}^0\rangle = \mathbf{P}_{j_2 k_2}^\mu \mathbf{p}^{m_2} |1\rangle = \delta_{k_2}^{m_2} \mathbf{P}_{j_2 m_2}^\mu |1\rangle$$

Local symmetry  $k_2$  of  $\mathbf{P}_{j_2 k_2}^\mu$  must match that  $m_2$  of  $|\mathbf{r}_{m_2}^0\rangle$

For example, base  $|\mathbf{r}_x^0\rangle = |\mathbf{r}_{0_2}^0\rangle = \mathbf{p}^{0_2}|1\rangle$  of  $d^{0_2}(C_2) \uparrow D_3$  gives zero for all  $\mathbf{P}_{j_2 k_2}^\mu$  except  $\mathbf{P}_{0_2 0_2}^{A_1}$ ,  $\mathbf{P}_{0_2 0_2}^{E_1}$ , and  $\mathbf{P}_{1_2 0_2}^{E_1}$ ,  $D_3$  projectors:  $\mathbf{P}_{0_2 0_2}^{A_1}$ ,  $\cancel{\mathbf{P}_{1_2 1_2}^{A_2}}$ ,  $\cancel{\mathbf{P}_{0_2 0_2}^{E_1}}$ ,  $\cancel{\mathbf{P}_{0_2 1_2}^{E_1}}$ ,  $\cancel{\mathbf{P}_{1_2 0_2}^{E_1}}$ ,  $\cancel{\mathbf{P}_{1_2 1_2}^{E_1}}$



## D<sub>3</sub>-Projection of $d^{m_2}(C_2) \uparrow D^3$ induced representation basis

$D_3 \supset C_2$  projectors  $\mathbf{P}_{0_20_2}^{A_1}$ ,  $\mathbf{P}_{1_21_2}^{A_2}$ ,  $\mathbf{P}_{0_20_2}^{E_1}$ ,  $\mathbf{P}_{0_21_2}^{E_1}$ ,  $\mathbf{P}_{1_20_2}^{E_1}$ ,  $\mathbf{P}_{1_21_2}^{E_1}$  must reduce induced representation  $d^{m_2}(C_2) \uparrow D_3$



But, which  $D_3$  projector  $\mathbf{P}_{j_2k_2}^\mu$  will work on base  $|\mathbf{r}_{m_2}^0\rangle = \mathbf{p}^{m_2}|1\rangle$  of induced representation  $d^{m_2}(C_2) \uparrow D_3$

$$\mathbf{P}_{j_2k_2}^\mu |\mathbf{r}_{m_2}^0\rangle = \mathbf{P}_{j_2k_2}^\mu \mathbf{p}^{m_2}|1\rangle = \delta_{k_2}^{m_2} \mathbf{P}_{j_2m_2}^\mu |1\rangle$$

Local symmetry  $k_2$  of  $\mathbf{P}_{j_2k_2}^\mu$  must match that  $m_2$  of  $|\mathbf{r}_{m_2}^0\rangle$

For example, base  $|\mathbf{r}_x^0\rangle = |\mathbf{r}_{0_2}^0\rangle = \mathbf{p}^{0_2}|1\rangle$  of  $d^{0_2}(C_2) \uparrow D_3$  gives zero for all  $\mathbf{P}_{j_2k_2}^\mu$  except  $\mathbf{P}_{0_20_2}^{A_1}$ ,  $\mathbf{P}_{0_20_2}^{E_1}$ , and  $\mathbf{P}_{1_20_2}^{E_1}$ ,

$D_3$  projectors:  $\mathbf{P}_{0_20_2}^{A_1}$ ,  $\cancel{\mathbf{P}_{1_21_2}^{A_2}}$ ,  $\cancel{\mathbf{P}_{0_20_2}^{E_1}}$ ,  $\cancel{\mathbf{P}_{0_21_2}^{E_1}}$ ,  $\cancel{\mathbf{P}_{1_20_2}^{E_1}}$ ,  $\cancel{\mathbf{P}_{1_21_2}^{E_1}}$   
 $\mathbf{P}_{xx}^{A_1}$ ,  $\cancel{\mathbf{P}_{yy}^{A_2}}$ ,  $\cancel{\mathbf{P}_{xx}^{E_1}}$ ,  $\cancel{\mathbf{P}_{xy}^{E_1}}$ ,  $\mathbf{P}_{yx}^{E_1}$ ,  $\cancel{\mathbf{P}_{yy}^{E_1}}$



## D<sub>3</sub>-Projection of $d^{m_2}(C_2) \uparrow D^3$ induced representation basis

$D_3 \supset C_2$  projectors  $\mathbf{P}_{0_20_2}^{A_1}$ ,  $\mathbf{P}_{1_21_2}^{A_2}$ ,  $\mathbf{P}_{0_20_2}^{E_1}$ ,  $\mathbf{P}_{0_21_2}^{E_1}$ ,  $\mathbf{P}_{1_20_2}^{E_1}$ ,  $\mathbf{P}_{1_21_2}^{E_1}$  must reduce induced representation  $d^{m_2}(C_2) \uparrow D_3$


But, which  $D_3$  projector  $\mathbf{P}_{j_2k_2}^\mu$  will work on base  $|\mathbf{r}_{m_2}^0\rangle = \mathbf{p}^{m_2}|1\rangle$  of induced representation  $d^{m_2}(C_2) \uparrow D_3$

$$\mathbf{P}_{j_2k_2}^\mu |\mathbf{r}_{m_2}^0\rangle = \mathbf{P}_{j_2k_2}^\mu \mathbf{p}^{m_2}|1\rangle = \delta_{k_2}^{m_2} \mathbf{P}_{j_2m_2}^\mu |1\rangle$$


Local symmetry  $k_2$  of  $\mathbf{P}_{j_2k_2}^\mu$  must match that  $m_2$  of  $|\mathbf{r}_{m_2}^0\rangle$

For example, base  $|\mathbf{r}_y^0\rangle = |\mathbf{r}_{1_2}^0\rangle = \mathbf{p}^{1_2}|1\rangle$  of  $d^{1_2}(C_2) \uparrow D_3$  gives zero for all  $\mathbf{P}_{j_2k_2}^\mu$  except  $\mathbf{P}_{1_21_2}^{A_2}$ ,  $\mathbf{P}_{0_21_2}^{E_1}$ , and  $\mathbf{P}_{1_21_2}^{E_1}$ ,

$D_3$  projectors:  $\mathbf{P}_{0_20_2}^{A_1}$ ,  $\mathbf{P}_{1_21_2}^{A_2}$ ,  $\mathbf{P}_{0_20_2}^{E_1}$ ,  $\mathbf{P}_{0_21_2}^{E_1}$ ,  $\mathbf{P}_{1_20_2}^{E_1}$ ,  $\mathbf{P}_{1_21_2}^{E_1}$   
 $\mathbf{P}_{xx}^{A_1}$ ,  $\mathbf{P}_{yy}^{A_2}$ ,  $\mathbf{P}_{xx}^{E_1}$ ,  $\mathbf{P}_{xy}^{E_1}$ ,  $\mathbf{P}_{yx}^{E_1}$ ,  $\mathbf{P}_{yy}^{E_1}$



These give the "y-band"



### *Frobenius Reciprocity Theorem*

$$\text{Number of } D^\alpha \text{ in } d^k(K) \uparrow G = \text{Number of } d^k \text{ in } D^\alpha(G) \downarrow K$$

## *Frobenius Reciprocity Theorem*

$$\text{Number of } D^\alpha \text{ in } d^k(K) \uparrow G = \text{Number of } d^k \text{ in } D^\alpha(G) \downarrow K$$

*..and regular representation*

$D_3 \supset C_1$	$0_1 = 1_1$
$A_1$	1
$A_2$	1
$E_1$	2

## Frobenius Reciprocity Theorem

$$\text{Number of } D^\alpha \text{ in } d^k(K) \uparrow G = \text{Number of } d^k \text{ in } D^\alpha(G) \downarrow K$$

..and regular representation

$D_3 \supset C_1$	$0_1 = 1_1$
$A_1$	1
$A_2$	1
$E_1$	2

$D_3 \supset C_2$	$0_2$	$1_2$
$A_1$	1	.
$A_2$	.	1
$E_1$	1	1

$D_3 \supset C_3$	$0_3$	$1_3$	$2_3$
$A_1$	1	.	.
$A_2$	1	.	.
$E_1$	.	1	1

Review: Symmetry reduction and splitting: Subduced irep  $D^\alpha(D_3) \downarrow C_2 = d^{0_2} \oplus d^{1_2} \oplus \dots$  correlation  
Symmetry induction and clustering: Induced rep  $d^a(C_2) \uparrow D_3 = D^\alpha \oplus D^\beta \oplus \dots$  correlation

$D_3$ - $C_2$  Coset structure of  $d^{m_2}(C_2) \uparrow D_3$  induced representation basis

$D_3$ -Projection of  $d^{m_2}(C_2) \uparrow D_3$  induced representation basis

Derivation of Frobenius reciprocity

→  $D_6 \supset D_2 \supset C_2 = D_3 \times C_2$  symmetry and outer product geometry ←

Irreducible characters

Irreducible representations

Correlations with  $D_6$  characters:

...and  $C_2(\mathbf{i}_3)$  characters.....and  $C_6(\mathbf{1}, \mathbf{h}^1, \mathbf{h}^2, \dots)$  characters



$D_6$  symmetry and induced representation band structure

Introduction to octahedral tetrahedral symmetry  $O_h \supset O \sim T_d \supset T$

$D_6 \supset D_2 \supset C_2 = D_3 \times C_2$  symmetry and outer product geometry

$D_6$  is the *outer product* ( $\times$ ) product  $D_3 \times C_2$  of  $D_3$  and  $C_2$ . (Requires  $C_2$  to commute with all of  $D_3$ .)

$$D_6 = D_3 \times C_2 = \{1, \mathbf{r}, \mathbf{r}^2, \mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3\} \times \{1, \mathbf{R}_z\}$$





$D_6 \supset D_2 \supset C_2 = D_3 \times C_2$  symmetry and outer product geometry

$D_6$  is the *outer product* ( $\times$ ) product  $D_3 \times C_2$  of  $D_3$  and  $C_2$ . (Requires  $C_2$  to commute with all of  $D_3$ .)

$$D_6 = D_3 \times C_2 = \{1, \mathbf{r}, \mathbf{r}^2, \mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3\} \times \{1, \mathbf{R}_z\}$$

$\times$  product and  $D_6$  operators. Define *hexagonal generator*  $\mathbf{h}_{(60^\circ)}$  of subgroup  $C_6 = \{1, \mathbf{h}, \mathbf{h}^2, \mathbf{h}^3, \mathbf{h}^4, \mathbf{h}^5\}$

$$D_6 = D_3 \times C_2 = \{1, \mathbf{r}, \mathbf{r}^2, \mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3, 1 \cdot \mathbf{R}_z, \mathbf{r} \cdot \mathbf{R}_z, \mathbf{r}^2 \cdot \mathbf{R}_z, \mathbf{i}_1 \cdot \mathbf{R}_z, \mathbf{i}_2 \cdot \mathbf{R}_z, \mathbf{i}_3 \cdot \mathbf{R}_z\}$$



$D_6 \supset D_2 \supset C_2 = D_3 \times C_2$  symmetry and outer product geometry



$D_6$  is the *outer product* ( $\times$ ) product  $D_3 \times C_2$  of  $D_3$  and  $C_2$ . (Requires  $C_2$  to commute with all of  $D_3$ .)

$$D_6 = D_3 \times C_2 = \{1, \mathbf{r}, \mathbf{r}^2, \mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3\} \times \{1, \mathbf{R}_z\}$$

$\times$  product and  $D_6$  operators. Define *hexagonal generator*  $\mathbf{h}_{(60^\circ)}$  of subgroup  $C_6 = \{1, \mathbf{h}, \mathbf{h}^2, \mathbf{h}^3, \mathbf{h}^4, \mathbf{h}^5\}$

$$D_6 = D_3 \times C_2 = \{1, \mathbf{r}, \mathbf{r}^2, \mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3, 1 \cdot \mathbf{R}_z, \mathbf{r} \cdot \mathbf{R}_z, \mathbf{r}^2 \cdot \mathbf{R}_z, \mathbf{i}_1 \cdot \mathbf{R}_z, \mathbf{i}_2 \cdot \mathbf{R}_z, \mathbf{i}_3 \cdot \mathbf{R}_z\}$$

$$D_6 = D_3 \times C_2 = \{1, \mathbf{h}^2, \mathbf{h}^4, \mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3, \mathbf{h}^3, \mathbf{h}^5, \mathbf{h}, \mathbf{j}_1, \mathbf{j}_2, \mathbf{j}_3\}$$



$D_6$

$D_6 \supset D_2 \supset C_2 = D_3 \times C_2$  symmetry and outer product geometry

$D_6$  is the *outer product* ( $\times$ ) product  $D_3 \times C_2$  of  $D_3$  and  $C_2$ . (Requires  $C_2$  to commute with all of  $D_3$ .)



$$D_6 = D_3 \times C_2 = \{1, \mathbf{r}, \mathbf{r}^2, \mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3\} \times \{1, \mathbf{R}_z\}$$

$\times$  product and  $D_6$  operators. Define *hexagonal generator*  $\mathbf{h}_{(60^\circ)}$  of subgroup  $C_6 = \{1, \mathbf{h}, \mathbf{h}^2, \mathbf{h}^3, \mathbf{h}^4, \mathbf{h}^5\}$

$$D_6 = D_3 \times C_2 = \{1, \mathbf{r}, \mathbf{r}^2, \mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3, 1 \cdot \mathbf{R}_z, \mathbf{r} \cdot \mathbf{R}_z, \mathbf{r}^2 \cdot \mathbf{R}_z, \mathbf{i}_1 \cdot \mathbf{R}_z, \mathbf{i}_2 \cdot \mathbf{R}_z, \mathbf{i}_3 \cdot \mathbf{R}_z\}$$

$$D_6 = D_3 \times C_2 = \{1, \mathbf{h}^2, \mathbf{h}^4, \mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3, \mathbf{h}^3, \mathbf{h}^5, \mathbf{h}, \mathbf{j}_1, \mathbf{j}_2, \mathbf{j}_3\}$$

Note:  $\mathbf{h}^2 = \mathbf{r}_{(120^\circ)}$  and  $\mathbf{h}^3 = \mathbf{R}_z(180^\circ)$  and  $\mathbf{h}^4 = \mathbf{r}^2$  and  $\mathbf{h}^5 = \mathbf{r} \cdot \mathbf{R}_z$



$D_6$

$D_6 \supset D_2 \supset C_2 = D_3 \times C_2$  symmetry and outer product geometry

$D_6$  is the *outer product* ( $\times$ ) product  $D_3 \times C_2$  of  $D_3$  and  $C_2$ . (Requires  $C_2$  to commute with all of  $D_3$ .)


$$D_6 = D_3 \times C_2 = \{1, \mathbf{r}, \mathbf{r}^2, \mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3\} \times \{1, \mathbf{R}_z\}$$

$\times$  product and  $D_6$  operators. Define *hexagonal generator*  $\mathbf{h}_{(60^\circ)}$  of subgroup  $C_6 = \{1, \mathbf{h}, \mathbf{h}^2, \mathbf{h}^3, \mathbf{h}^4, \mathbf{h}^5\}$


$$D_6 = D_3 \times C_2 = \{1, \mathbf{r}, \mathbf{r}^2, \mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3, 1 \cdot \mathbf{R}_z, \mathbf{r} \cdot \mathbf{R}_z, \mathbf{r}^2 \cdot \mathbf{R}_z, \mathbf{i}_1 \cdot \mathbf{R}_z, \mathbf{i}_2 \cdot \mathbf{R}_z, \mathbf{i}_3 \cdot \mathbf{R}_z\}$$

$$D_6 = D_3 \times C_2 = \{1, \mathbf{h}^2, \mathbf{h}^4, \mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3, \mathbf{h}^3, \mathbf{h}^5, \mathbf{h}, \mathbf{j}_1, \mathbf{j}_2, \mathbf{j}_3\}$$

Note:  $\mathbf{h}^2 = \mathbf{r}_{(120^\circ)}$  and  $\mathbf{h}^3 = \mathbf{R}_z(180^\circ)$  and  $\mathbf{h}^4 = \mathbf{r}^2$  and  $\mathbf{h}^5 = \mathbf{r} \cdot \mathbf{R}_z$



NOTE:  
The  $\mathbf{i}_a$  and  $\mathbf{j}_b$  do not flip over the potential plot.



Electrostatic potential  $V(\phi)$  doesn't care which way is "up." Wells remain wells, and barriers remain barriers under all  $D_6$  operations.

Review: Symmetry reduction and splitting: Subduced irep  $D^\alpha(D_3) \downarrow C_2 = d^{0_2} \oplus d^{1_2} \oplus \dots$  correlation  
Symmetry induction and clustering: Induced rep  $d^a(C_2) \uparrow D_3 = D^\alpha \oplus D^\beta \oplus \dots$  correlation

$D_3$ - $C_2$  Coset structure of  $d^{m_2}(C_2) \uparrow D_3$  induced representation basis

$D_3$ -Projection of  $d^{m_2}(C_2) \uparrow D_3$  induced representation basis

Derivation of Frobenius reciprocity

$D_6 \supset D_2 \supset C_2 = D_3 \times C_2$  symmetry and outer product geometry

→ Irreducible characters ←

Irreducible representations

Correlations with  $D_6$  characters:

...and  $C_2(\mathbf{i}_3)$  characters.....and  $C_6(\mathbf{1}, \mathbf{h}^1, \mathbf{h}^2, \dots)$  characters

$D_6$  symmetry and induced representation band structure

Introduction to octahedral tetrahedral symmetry  $O_h \supset O \sim T_d \supset T$

*D<sub>6</sub> ⊃ D<sub>2</sub> ⊃ C<sub>2</sub> = D<sub>3</sub> × C<sub>2</sub> Irreducible characters*

D <sub>3</sub>	1	{r, r <sup>2</sup> }	{i <sub>1</sub> , i <sub>2</sub> , i <sub>3</sub> }
χ <sup>A<sub>1</sub></sup> (g)	1	1	1
χ <sup>A<sub>2</sub></sup> (g)	1	1	-1
χ <sup>E<sub>1</sub></sup> (g)	2	-1	0

C <sub>2</sub> <sup>Z</sup>	1	R <sub>z</sub>
(A)	1	1
(B)	1	-1

$$\times \begin{array}{|c|cc|} \hline & 1 & R_z \\ \hline (A) & 1 & 1 \\ (B) & 1 & -1 \\ \hline \end{array}$$


D <sub>3</sub> × C <sub>2</sub> <sup>Z</sup>	1	{r, r <sup>2</sup> }	{i <sub>1</sub> , i <sub>2</sub> , i <sub>3</sub> }	1 · R <sub>z</sub>	{r, r <sup>2</sup> } · R <sub>z</sub>	{i <sub>1</sub> , i <sub>2</sub> , i <sub>3</sub> } · R <sub>z</sub>
A <sub>1</sub> · (A)	1·1	1·1	1·1	1·1	1·1	1·1
A <sub>2</sub> · (A)	1·1	1·1	-1·1	1·1	1·1	-1·1
E <sub>1</sub> · (A)	2·1	-1·1	0·1	2·1	-1·1	0·1
A <sub>1</sub> · (B)	1·1	1·1	1·1	1·(-1)	1·(-1)	1·(-1)
A <sub>2</sub> · (B)	1·1	1·1	-1·1	1·(-1)	1·(-1)	-1·(-1)
E <sub>1</sub> · (B)	2·1	-1·1	0·1	2·(-1)	-1·(-1)	0·(-1)

$D_6 \supset D_2 \supset C_2 = D_3 \times C_2$  Irreducible characters

$D_3$	1	$\{r, r^2\}$	$\{i_1, i_2, i_3\}$
$\chi^{A_1}(g)$	1	1	1
$\chi^{A_2}(g)$	1	1	-1
$\chi^{E_1}(g)$	2	-1	0

$C_2$	1	$R_z$
(A)	1	1
(B)	1	-1

$D_3 \times C_2$	1	$\{r, r^2\}$	$\{i_1, i_2, i_3\}$	$1 \cdot R_z$	$\{r, r^2\} \cdot R_z$	$\{i_1, i_2, i_3\} \cdot R_z$
$A_1 \cdot (A)$	1·1	1·1	1·1	1·1	1·1	1·1
$A_2 \cdot (A)$	1·1	1·1	-1·1	1·1	1·1	-1·1
$E_1 \cdot (A)$	2·1	-1·1	0·1	2·1	-1·1	0·1
$A_1 \cdot (B)$	1·1	1·1	1·1	1·(-1)	1·(-1)	1·(-1)
$A_2 \cdot (B)$	1·1	1·1	-1·1	1·(-1)	1·(-1)	-1·(-1)
$E_1 \cdot (B)$	2·1	-1·1	0·1	2·(-1)	-1·(-1)	0·(-1)



$D_6 \supset D_2 \supset C_2 = D_3 \times C_2$  Irreducible characters

$D_3$	1	$\{r, r^2\}$	$\{i_1, i_2, i_3\}$
$\chi^{A_1}(g)$	1	1	1
$\chi^{A_2}(g)$	1	1	-1
$\chi^{E_1}(g)$	2	-1	0


$C_2$	1	$R_z$
(A)	1	1
(B)	1	-1

$D_3 \times C_2$	1	$\{r, r^2\}$	$\{i_1, i_2, i_3\}$	$1 \cdot R_z$	$\{r, r^2\} \cdot R_z$	$\{i_1, i_2, i_3\} \cdot R_z$
$A_1 \cdot (A)$	1·1	1·1	1·1	1·1	1·1	1·1
$A_2 \cdot (A)$	1·1	1·1	-1·1	1·1	1·1	-1·1
$E_1 \cdot (A)$	2·1	-1·1	0·1	2·1	-1·1	0·1
$A_1 \cdot (B)$	1·1	1·1	1·1	1·(-1)	1·(-1)	1·(-1)
$A_2 \cdot (B)$	1·1	1·1	-1·1	1·(-1)	1·(-1)	-1·(-1)
$E_1 \cdot (B)$	2·1	-1·1	0·1	2·(-1)	-1·(-1)	0·(-1)

Recall  $C_2 \times C_2 = D_2 = \{1, R_x, R_z, R_y\}$  characters


(Lect.12 p.50-60)

$D_6$  has  $D_2 = \{1, i_3, h^3, j_3\}$  subgroup



$D_2$

1	$R_x$	$R_z$	$R_y$
$R_x$	1	$R_y$	$R_z$
$R_z$	$R_y$	1	$R_x$
$R_y$	$R_z$	$R_x$	1



$C_2^X$	1	$R_X$	$C_2^Z$	1	$R_Z$	$C_2^Y \times C_2^Z$	1·1	$R_X \cdot 1$	$1 \cdot R_Z$	$R_X \cdot R_Z$
$+ = 1$	1	1	$+ = A$	1	1	$+ \cdot + = A_1$	1·1	1·1	1·1	1·1
$- = 2$	1	-1	$- = B$	1	-1	$- \cdot + = A_2$	1·1	-1·1	1·1	-1·1
						$+ \cdot - = B_1$	1·1	1·1	k(-1)	k(-1)
						$- \cdot - = B_2$	1·1	-1·1	1(-1)	-1(-1)

## \$D\_6 \supset D\_2 \supset C\_2 = D\_3 \times C\_2\$ Irreducible characters

\$D_3\$	1	\$\{r, r^2\}\$	\$\{i_1, i_2, i_3\}\$
\$\chi^{A_1}(g)\$	1	1	1
\$\chi^{A_2}(g)\$	1	1	-1
\$\chi^{E_1}(g)\$	2	-1	0

\$C_2^Z\$	1	\$R_z\$
(A)	1	1
(B)	1	-1

	1	\$R_z\$
(A)	1	1
(B)	1	-1

\$D_3 \times C_2^Z\$	1	\$\{r, r^2\}\$	\$\{i_1, i_2, i_3\}\$	\$1 \cdot R_z\$	\$\{r, r^2\} \cdot R_z\$	\$\{i_1, i_2, i_3\} \cdot R_z\$
\$A_1 \cdot (A)\$	1·1	1·1	1·1	1·1	1·1	1·1
\$A_2 \cdot (A)\$	1·1	1·1	-1·1	1·1	1·1	-1·1
\$E_1 \cdot (A)\$	2·1	-1·1	0·1	2·1	-1·1	0·1
\$A_1 \cdot (B)\$	1·1	1·1	1·1	1·(-1)	1·(-1)	1·(-1)
\$A_2 \cdot (B)\$	1·1	1·1	-1·1	1·(-1)	1·(-1)	-1·(-1)
\$E_1 \cdot (B)\$	2·1	-1·1	0·1	2·(-1)	-1·(-1)	0·(-1)

Recall \$C\_2 \times C\_2 = D\_2 = \{1, R\_x, R\_z, R\_y\}\$ characters


(Lect.12 p.50-60)

\$D\_6\$ has \$D\_2 = \{1, i\_3, h^3, j\_3\}\$ subgroup

\$R\_z(180^\circ)\$

\$D\_2\$

1	\$R_x\$	\$R_z\$	\$R_y\$
\$R_x\$	1	\$R_y\$	\$R_z\$
\$R_z\$	\$R_y\$	1	\$R_x\$
\$R_y\$	\$R_z\$	\$R_x\$	1



$$\chi_g^\mu(D_6) =$$

\$D_3 \times C_2^z\$	1	\$\{h^2, h^4\}\$	\$\{i_1, i_2, i_3\}\$	\$h^3\$	\$\{h, h^5\}\$	\$\{j_1, j_2, j_3\}\$
\$A_1\$	1	1	1	1	1	1
\$A_2\$	1	1	-1	1	1	-1
\$E_2\$	2	-1	0	2	-1	0
\$B_1\$	1	1	1	-1	-1	-1
\$B_2\$	1	1	-1	-1	-1	1
\$E_1\$	2	-1	0	-2	1	0

\$C_2^X\$	1	\$R_X\$
+\$=1\$	1	1
-\$=2\$	1	-1

\$C_2^Z\$	1	\$R_Z\$
+\$=A\$	1	1
-\$=B\$	1	-1

\$C_2^X \times C_2^Z\$	1·1	\$R_X \cdot 1\$	\$1 \cdot R_Z\$	\$R_X \cdot R_Z\$
+\$\cdot+\$ = \$A_1\$	1·1	1·1	1·1	1·1
-\$\cdot+\$ = \$A_2\$	1·1	-1·1	1·1	-1·1
+\$\cdot-\$ = \$B_1\$	1·1	1·1	1(-1)	1(-1)
-\$\cdot-\$ = \$B_2\$	1·1	-1·1	1(-1)	-1(-1)

## \$D\_6 \supset D\_2 \supset C\_2 = D\_3 \times C\_2\$ Irreducible characters

\$D_3\$	1	\$\{r, r^2\}\$	\$\{i_1, i_2, i_3\}\$
\$\chi^{A_1}(g)\$	1	1	1
\$\chi^{A_2}(g)\$	1	1	-1
\$\chi^{E_1}(g)\$	2	-1	0

\$C_2^Z\$	1	\$R_z\$
(A)	1	1
(B)	1	-1

\$D_3 \times C_2^Z\$	1	\$\{r, r^2\}\$	\$\{i_1, i_2, i_3\}\$	\$1 \cdot R_z\$	\$\{r, r^2\} \cdot R_z\$	\$\{i_1, i_2, i_3\} \cdot R_z\$
\$A_1 \cdot (A)\$	1·1	1·1	1·1	1·1	1·1	1·1
\$A_2 \cdot (A)\$	1·1	1·1	-1·1	1·1	1·1	-1·1
\$E_1 \cdot (A)\$	2·1	-1·1	0·1	2·1	-1·1	0·1
\$A_1 \cdot (B)\$	1·1	1·1	1·1	1·(-1)	1·(-1)	1·(-1)
\$A_2 \cdot (B)\$	1·1	1·1	-1·1	1·(-1)	1·(-1)	-1·(-1)
\$E_1 \cdot (B)\$	2·1	-1·1	0·1	2·(-1)	-1·(-1)	0·(-1)

Recall \$C\_2 \times C\_2 = D\_2 = \{1, R\_x, R\_z, R\_y\}\$ characters


(Lect.12 p.50-60)

\$D\_6\$ has \$D\_2 = \{1, i\_3, h^3, j\_3\}\$ subgroup

\$R\_z(180^\circ)\$

\$D\_2\$

1	\$R_x\$	\$R_z\$	\$R_y\$
\$R_x\$	1	\$R_y\$	\$R_z\$
\$R_z\$	\$R_y\$	1	\$R_x\$
\$R_y\$	\$R_z\$	\$R_x\$	1



$$\chi_g^\mu(D_6) =$$

\$D_3 \times C_2^z\$	1	\$\{h^2, h^4\}\$	\$\{i_1, i_2, i_3\}\$	\$h^3\$	\$\{h, h^5\}\$	\$\{j_1, j_2, j_3\}\$
\$A_1\$	1	1	1	1	1	1
\$A_2\$	1	1	-1	1	1	-1
\$E_2\$	2	-1	0	2	-1	0
\$B_1\$	1	1	1	-1	-1	-1
\$B_2\$	1	1	-1	-1	-1	1
\$E_1\$	2	-1	0	-2	1	0

Let \$X\$-rotation

or  
180° \$X\$-flip \$i\_3\$  
determine

\$A\_1\$ or \$B\_1\$ vs \$A\_2\$ or \$B\_2\$  
(+1) vs (-1)

Let unit translation

or  
60° hex-Z rotation \$h\$  
determine  
\$A\_p\$ vs \$B\_p\$  
(+1) vs (-1)

So also does:  
180° \$h^3\$

\$C_2^X\$	1	\$R_X\$	\$C_2^Z\$	1	\$R_X \cdot 1\$	\$1 \cdot R_Z\$	\$R_X \cdot R_Z\$
+\$=1\$	1	1	+\$=A\$	1	1·1	1·1	1·1
-\$=2\$	1	-1	-\$=B\$	1	1·1	-1·1	-1·1

+\$+\cdot= A_1\$	1·1	1·1	1·1	1·1
-\$+\cdot= A_2\$	1·1	-1·1	1·1	-1·1
+\$-\cdot= B_1\$	1·1	1·1	1(-1)	1(-1)
-\$-\cdot= B_2\$	1·1	-1·1	1(-1)	-1(-1)

Review: Symmetry reduction and splitting: Subduced irep  $D^\alpha(D_3) \downarrow C_2 = d^{0_2} \oplus d^{1_2} \oplus \dots$  correlation  
Symmetry induction and clustering: Induced rep  $d^a(C_2) \uparrow D_3 = D^\alpha \oplus D^\beta \oplus \dots$  correlation

$D_3$ - $C_2$  Coset structure of  $d^{m_2}(C_2) \uparrow D_3$  induced representation basis

$D_3$ -Projection of  $d^{m_2}(C_2) \uparrow D_3$  induced representation basis

Derivation of Frobenius reciprocity

$D_6 \supset D_2 \supset C_2 = D_3 \times C_2$  symmetry and outer product geometry

→ Irreducible characters

Irreducible representations

Correlations with  $D_6$  characters:


...and  $C_2(\mathbf{i}_3)$  characters.....and  $C_6(\mathbf{1}, \mathbf{h}^1, \mathbf{h}^2, \dots)$  characters

$D_6$  symmetry and induced representation band structure




Introduction to octahedral tetrahedral symmetry  $O_h \supset O \sim T_d \supset T$

$D_6 \supset D_2 \supset C_2 = D_3 \times C_2$  Irreducible representations



$D_6 \supset D_2 \supset C_2 = D_3 \times C_2$  Irreducible representations



$$g = \mathbf{1}, \mathbf{r} = \mathbf{h}^2, \mathbf{r}^2 = \mathbf{h}^4, \mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3, \mathbf{h}^3, \mathbf{h}^3\mathbf{r} = \mathbf{h}^5, \mathbf{h}^3\mathbf{r}^2 = \mathbf{h}^1, \mathbf{h}^3\mathbf{i}_1 = \mathbf{j}_1, \mathbf{h}^3\mathbf{i}_2 = \mathbf{j}_2, \mathbf{h}^3\mathbf{i}_3 = \mathbf{j}_3$$

$D^{A_1}(g) =$	1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1
$D^{A_2}(g) =$	1, 1, 1, -1, -1, -1, 1, 1, 1, -1, -1, -1
$D^{E_2}(g) =$	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{-1}{2} \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{-\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{-\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{-1}{2} \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{-\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{-\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$
$D^{B_1}(g) =$	1, 1, 1, 1, 1, 1, -1, -1, -1, -1, -1, -1
$D^{B_2}(g) =$	1, 1, 1, -1, -1, -1, -1, -1, -1, 1, 1, 1
$D^{E_1}(g) =$	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{-1}{2} \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{-\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{-\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} \frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{-\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}, \begin{pmatrix} \frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}, \begin{pmatrix} \frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{-\sqrt{3}}{2} & \frac{-1}{2} \end{pmatrix}, \begin{pmatrix} \frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{-1}{2} \end{pmatrix}, \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$


Let  $X$ -rotation

or  
180°  $X$ -flip  $i_3$   
determines

$A_1$  or  $B_1$  vs  $A_2$  or  $B_2$

(+1) vs (-1)

$D_6 \supset D_2 \supset C_2 = D_3 \times C_2$  Irreducible representations



$D_6$

$$g = 1, r = h^2, r^2 = h^4, i_1, i_2, i_3, h^3, h^3r = h^5, h^3r^2 = h^1, h^3i_1 = j_1, h^3i_2 = j_2, h^3i_3 = j_3$$

$$D^{A_1}(g) = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & -1 & -1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 \end{pmatrix}$$

$$D^{A_2}(g) = \begin{pmatrix} 1 & 1 & 1 & -1 & -1 & -1 & 1 & 1 & 1 & -1 & -1 & -1 \end{pmatrix}$$

$$D^{E_2}(g) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{-1}{2} \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{-\sqrt{3}}{2} & \frac{-1}{2} \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{-\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{-1}{2} \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{-\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$D^{B_1}(g) = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & -1 & -1 & -1 & -1 & -1 & -1 \end{pmatrix}$$


$$D^{B_2}(g) = \begin{pmatrix} 1 & 1 & 1 & -1 & -1 & -1 & -1 & -1 & -1 & 1 & 1 & 1 \end{pmatrix}$$

$$D^{E_1}(g) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{-1}{2} \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{-\sqrt{3}}{2} & \frac{-1}{2} \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}, \begin{pmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{-\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} \frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{-\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}, \begin{pmatrix} \frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}, \begin{pmatrix} \frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{-\sqrt{3}}{2} & \frac{-1}{2} \end{pmatrix}, \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$$

Let  $X$ -rotation  
or  
 $180^\circ$   $X$ -flip  $i_3$   
determines  
 $A_1$  or  $B_1$  vs  $A_2$  or  $B_2$   
 $(+1)$  vs  $(-1)$

Let unit translation  
or  
 $60^\circ$  hex-Z rotation  $h$   
determine  
 $A_p$  vs  $B_p$   
 $(+1)$  vs  $(-1)$   
So also does:  
 $180^\circ h^3$

$D_6 \supset D_2 \supset C_2 = D_3 \times C_2$  Irreducible representations



$g =$	$1$	$r = h^2$	$r^2 = h^4$	$i_1$	$i_2$	$i_3$	$h^3$	$h^3 r = h^5$	$h^3 r^2 = h^1$	$h^3 i_1 = j_1$	$h^3 i_2 = j_2$	$h^3 i_3 = j_3$
$D^{A_1}(g) =$	1	1	1	1	1	1	1	1	1	1	1	1
$D^{A_2}(g) =$	1	1	1	-1	-1	-1	-1	1	1	-1	-1	-1
$D^{E_2}(g) =$	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$\begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{-1}{2} \end{pmatrix}$	$\begin{pmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{-\sqrt{3}}{2} & \frac{-1}{2} \end{pmatrix}$	$\begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}$	$\begin{pmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{-\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}$	$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$	$\begin{pmatrix} 1 & 0 \\ \frac{\sqrt{3}}{2} & \frac{-1}{2} \end{pmatrix}$	$\begin{pmatrix} -1 & 0 \\ -\frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}$	$\begin{pmatrix} -1 & 0 \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}$	$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$
$D^{B_1}(g) =$	1	1	1	1	1	1	1	-1	-1	-1	-1	-1
$D^{B_2}(g) =$	1	1	1	-1	-1	-1	-1	-1	-1	1	1	1
$D^{E_1}(g) =$	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$\begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{-1}{2} \end{pmatrix}$	$\begin{pmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{-\sqrt{3}}{2} & \frac{-1}{2} \end{pmatrix}$	$\begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}$	$\begin{pmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{-\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}$	$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$	$\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$	$\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$	$\begin{pmatrix} 1 & 0 \\ \frac{1}{2} & \frac{\sqrt{3}}{2} \end{pmatrix}$	$\begin{pmatrix} -1 & 0 \\ -\frac{1}{2} & \frac{1}{2} \end{pmatrix}$	$\begin{pmatrix} 1 & 0 \\ \frac{1}{2} & -\frac{\sqrt{3}}{2} \end{pmatrix}$	$\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$

Let  $X$ -rotation  
or  
 $180^\circ$   $X$ -flip  $i_3$   
determines  
 $A_1$  or  $B_1$  vs  $A_2$  or  $B_2$   
 $(+1)$  vs  $(-1)$

Let unit translation  
or  
 $60^\circ$  hex-Z rotation  $h$   
determine  
 $A_p$  vs  $B_p$   
 $(+1)$  vs  $(-1)$   
So also does:  
 $180^\circ h^3$

$Y$ -rotation  
or  
 $180^\circ$  flip  $j_3$   
is product  
 $i_3 h^3 = h^3 i_3$

*Review: Symmetry reduction and splitting: Subduced irep  $D^\alpha(D_3) \downarrow C_2 = d^{0_2} \oplus d^{1_2} \oplus \dots$  correlation  
Symmetry induction and clustering: Induced rep  $d^a(C_2) \uparrow D_3 = D^\alpha \oplus D^\beta \oplus \dots$  correlation*

*$D_3$ - $C_2$  Coset structure of  $d^{m_2}(C_2) \uparrow D_3$  induced representation basis*

*$D_3$ -Projection of  $d^{m_2}(C_2) \uparrow D_3$  induced representation basis*

*Derivation of Frobenius reciprocity*

*$D_6 \supset D_2 \supset C_2 = D_3 \times C_2$  symmetry and outer product geometry*

*Irreducible characters*

*Irreducible representations*

*Correlations with  $D_6$  characters:*

*...and  $C_2(\mathbf{i}_3)$  characters.....and  $C_6(\mathbf{1}, \mathbf{h}^1, \mathbf{h}^2, \dots)$  characters*

*$D_6$  symmetry and induced representation band structure*



*Introduction to octahedral tetrahedral symmetry  $O_h \supset O \sim T_d \supset T$*

*Correlations with  $D_6$  characters:*  $\chi_g^\mu(D_6) =$

*...and  $C_2(\mathbf{i}_3)$  characters:*

$C_2^X$	1	$\mathbf{i}_3$
$0_2$	1	1
$1_2$	1	-1

$D_3 \times C_2^z$	1	$\{\mathbf{h}^2, \mathbf{h}^4\}$	$\{\mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3\}$	$\mathbf{h}^3$	$\{\mathbf{h}, \mathbf{h}^5\}$	$\{\mathbf{j}_1, \mathbf{j}_2, \mathbf{j}_3\}$
$A_1$	1	1	1	1	1	1
$A_2$	1	1	-1	1	1	-1
$E_2$	2	-1	0	2	-1	0
$B_1$	1	1	1	-1	-1	-1
$B_2$	1	1	-1	-1	-1	1
$E_1$	2	-1	0	-2	1	0

*Let X-rotation*

*or  
180° X-flip  $\mathbf{i}_3$   
determine*

$A_1$  or  $B_1$  vs  $A_2$  or  $B_2$   
 $(+1)$  vs  $(-1)$

$D_3 \supset C_2(\mathbf{i}_3)$	$0_2$	$1_2$
$A_1$	1	.
$A_2$	.	1
$E_2$	1	1
$B_1$	1	.
$B_2$	.	1
$E_1$	1	1

Correlations with  $D_6$  characters:  $\chi_g^\mu(D_6) =$

$D_3 \times C_2^z$	1	$\{\mathbf{h}^2, \mathbf{h}^4\}$	$\{\mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3\}$	$\mathbf{h}^3$	$\{\mathbf{h}, \mathbf{h}^5\}$	$\{\mathbf{j}_1, \mathbf{j}_2, \mathbf{j}_3\}$
$A_1$	1	1	1	1	1	1
$A_2$	1	1	-1	1	1	-1
$E_2$	2	-1	0	2	-1	0
$B_1$	1	1	1	-1	-1	-1
$B_2$	1	1	-1	-1	-1	1
$E_1$	2	-1	0	-2	1	0

...and  $C_2(\mathbf{i}_3)$  characters:

$C_2$	1	$\mathbf{i}_3$
$0_2$	1	1
$1_2$	1	-1

...and  $C_6(1, \mathbf{h}^1, \mathbf{h}^2, \dots)$  characters:

$C_6$	$p=0$	1	2	3	4	5	$C_6$
$0_6$	1	$\mathbf{h}^1$	$\mathbf{h}^2$	$\mathbf{h}^3$	$\mathbf{h}^{-2}$	$\mathbf{h}^{-1}$	$0_6$
$1_6$	1	$\epsilon^1$	$\epsilon^2$	-1	$\epsilon^{-2}$	$\epsilon^{-1}$	$1_6$
$2_6$	1	$\epsilon^2$	$\epsilon^4$	1	$\epsilon^{-4}$	$\epsilon^{-2}$	$2_6$
$3_6$	1	-1	1	-1	1	-1	$3_6$
$-2_6$	1	$\epsilon^{-2}$	$\epsilon^{-4}$	-1	$\epsilon^4$	$\epsilon^2$	$-2_6$
$-1_6$	1	$\epsilon^{-1}$	$\epsilon^{-2}$	$\epsilon^{-3}$	$\epsilon^{-4}$	$\epsilon^{-5}$	$-1_6$

$(\epsilon = e^{\pi i/3})$

Let X-rotation  
or  
180° X-flip  $\mathbf{i}_3$   
determine  
 $A_1$  or  $B_1$  vs  $A_2$  or  $B_2$   
(+1) vs (-1)

Let unit translation  
or  
60° hex-Z rotation  $\mathbf{h}$   
determine  
 $A_p$  vs  $B_p$   
(+1) vs (-1)  
So also does:  
180°  $\mathbf{h}^3$

Y-rotation  
or  
180° flip  $\mathbf{j}_3$   
is product  
 $\mathbf{i}_3 \mathbf{h}^3 = \mathbf{h}^3 \mathbf{i}_3$

Correlations with  $D_6$  characters:  $\chi_g^\mu(D_6) =$

$D_3 \times C_2^z$	1	$\{\mathbf{h}^2, \mathbf{h}^4\}$	$\{\mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3\}$	$\mathbf{h}^3$	$\{\mathbf{h}, \mathbf{h}^5\}$	$\{\mathbf{j}_1, \mathbf{j}_2, \mathbf{j}_3\}$
$A_1$	1	1	1	1	1	1
$A_2$	1	1	-1	1	1	-1
$E_2$	2	-1	0	2	-1	0
$B_1$	1	1	1	-1	-1	-1
$B_2$	1	1	-1	-1	-1	1
$E_1$	2	-1	0	-2	1	0

...and  $C_2(\mathbf{i}_3)$  characters:

$C_2^X$	1	$\mathbf{i}_3$
$0_2$	1	1
$1_2$	1	-1

...and  $C_6(1, \mathbf{h}^1, \mathbf{h}^2, \dots)$  characters:

$C_6$	$p=0$	1	2	3	4	5	$C_6$	1	$\mathbf{h}^1$	$\mathbf{h}^2$	$\mathbf{h}^3$	$\mathbf{h}^{-2}$	$\mathbf{h}^{-1}$
$0_6$	1	1	1	1	1	1	$0_6$	1	1	1	1	1	1
$1_6$	1	$\epsilon^1$	$\epsilon^2$	-1	$\epsilon^{-2}$	$\epsilon^{-1}$	$1_6$	1	$\epsilon^1$	$\epsilon^2$	-1	$\epsilon^{-2}$	$\epsilon^{-1}$
$2_6$	1	$\epsilon^2$	$\epsilon^4$	1	$\epsilon^{-4}$	$\epsilon^{-2}$	$2_6$	1	$\epsilon^2$	$\epsilon^4$	1	$\epsilon^{-4}$	$\epsilon^{-2}$
$3_6$	1	-1	1	-1	1	-1	$3_6$	1	-1	1	-1	1	-1
$-2_6$	1	$\epsilon^{-2}$	$\epsilon^{-4}$	-1	$\epsilon^4$	$\epsilon^2$	$-2_6$	1	$\epsilon^{-2}$	$\epsilon^{-4}$	-1	$\epsilon^4$	$\epsilon^2$
$-1_6$	1	$\epsilon^{-1}$	$\epsilon^{-2}$	$\epsilon^{-3}$	$\epsilon^{-4}$	$\epsilon^{-5}$	$-1_6$	1	$\epsilon^{-1}$	$\epsilon^{-2}$	$\epsilon^{-3}$	$\epsilon^{-4}$	$\epsilon^{-5}$

$(\epsilon = e^{\pi i/3})$

Let X-rotation  
or  
180° X-flip  $\mathbf{i}_3$   
determine  
 $A_1$  or  $B_1$  vs  $A_2$  or  $B_2$   
(+1) vs (-1)

Let unit translation  
or  
60° hex-Z rotation  $\mathbf{h}$   
determine  
 $A_p$  vs  $B_p$   
(+1) vs (-1)  
So also does:  
180°  $\mathbf{h}^3$

Y-rotation  
or  
180° flip  $\mathbf{j}_3$   
is product  
 $\mathbf{i}_3 \mathbf{h}^3 = \mathbf{h}^3 \mathbf{i}_3$

$D_3 \supseteq C_2^X(\mathbf{i}_3)$	$0_2$	$1_2$
$A_1$	1	.
$A_2$	.	1
$E_2$	1	1
$B_1$	1	.
$B_2$	.	1
$E_1$	1	1

$D_6 \supseteq C_6(\mathbf{h})$	$0_6$	$1_6$	$2_6$	$3_6$	$4_6$	$5_6$
$A_1$	1	.	.	.	.	.
$A_2$	1	.	.	.	.	.
$E_2$	.	.	1	.	1	.
$B_2$	.	.	.	1	.	.
$B_1$	.	.	.	1	.	.
$E_1$	.	1	.	.	.	1

Review: Symmetry reduction and splitting: Subduced irep  $D^\alpha(D_3) \downarrow C_2 = d^{0_2} \oplus d^{1_2} \oplus \dots$  correlation

Symmetry induction and clustering: Induced rep  $d^a(C_2) \uparrow D_3 = D^\alpha \oplus D^\beta \oplus \dots$  correlation

$D_3$ - $C_2$  Coset structure of  $d^{m_2}(C_2) \uparrow D_3$  induced representation basis

$D_3$ -Projection of  $d^{m_2}(C_2) \uparrow D_3$  induced representation basis

Derivation of Frobenius reciprocity

$D_6 \supset D_2 \supset C_2 = D_3 \times C_2$  symmetry and outer product geometry

Irreducible characters

Irreducible representations

Correlations with  $D_6$  characters:

...and  $C_2(\mathbf{i}_3)$  characters.....and  $C_6(\mathbf{1}, \mathbf{h}^1, \mathbf{h}^2, \dots)$  characters


$\rightarrow D_6$  symmetry and induced representation band structure  $\leftarrow$

Introduction to octahedral tetrahedral symmetry  $O_h \supset O \sim T_d \supset T$

# $D_6$ symmetry and induced representation band structure

$D_6$  Band structure and related induced representations

For high energy above potential barriers local  $C_2$  symmetry is replaced by global  $C_6$  angular momentum doublets such as  $E_{\pm m}$ ,  $A_1A_2$ , and  $B_1B_2$




For low energy deep in potential local  $C_2$  symmetry dominates and the bands  $A_1E_1E_2B_1$  and  $B_2E_2E_1A_2$  that become tight clusters

$D_3 \supset C_2(j_3)$	$0_2$	$1_2$
$A_1$	1	.
$A_2$	.	1
$E_2$	1	1
$B_2$	.	1
$B_1$	1	.
$E_1$	1	1

$D_6 \supset C_3(h)$	$0_6$	$1_6$	$2_6$	$3_6$	$4_6$	$5_6$
$A_1$	1	.	.	.	.	.
$A_2$	1	.	.	.	.	.
$E_2$	.	.	1	.	1	.
$B_2$	.	.	.	1	.	.
$B_1$	.	.	.	1	.	.
$E_1$	.	1	.	.	.	1

# $D_6$ symmetry and induced representation band structure



For high energy above potential barriers local  $C_2$  symmetry is replaced by global  $C_6$  angular momentum doublets such as  $E_{\pm m}$ ,  $A_1A_2$ , and  $B_1B_2$

$D_6 \supset C_3(h)$	$0_6$	$1_6$	$2_6$	$3_6$	$4_6$	$5_6$
$A_1$	1	.	.	.	.	.
$A_2$	1	.	.	.	.	.
$E_2$	.	.	1	.	1	.
$B_2$	.	.	.	1	.	.
$B_1$	.	.	.	1	.	.
$E_1$	.	1	.	.	.	1

For low energy deep in potential local  $C_2$  symmetry dominates and the bands  $A_1E_1E_2B_1$  and  $B_2E_2E_1A_2$  then become tight clusters

$D_3 \supset C_2(j_3)$	$0_2$	$1_2$
$A_1$	1	.
$A_2$	.	1
$E_2$	1	1
$B_2$	.	1
$B_1$	1	.
$E_1$	1	1

*Review: Symmetry reduction and splitting: Subduced irep  $D^\alpha(D_3) \downarrow C_2 = d^{0_2} \oplus d^{1_2} \oplus \dots$  correlation  
Symmetry induction and clustering: Induced rep  $d^a(C_2) \uparrow D_3 = D^\alpha \oplus D^\beta \oplus \dots$  correlation*

*$D_3$ - $C_2$  Coset structure of  $d^{m_2}(C_2) \uparrow D_3$  induced representation basis*

*$D_3$ -Projection of  $d^{m_2}(C_2) \uparrow D_3$  induced representation basis*

*Derivation of Frobenius reciprocity*

*$D_6 \supset D_2 \supset C_2 = D_3 \times C_2$  symmetry and outer product geometry*

*Irreducible characters*

*Irreducible representations*

*Correlations with  $D_6$  characters:*

*...and  $C_2(\mathbf{i}_3)$  characters.....and  $C_6(\mathbf{1}, \mathbf{h}^1, \mathbf{h}^2, \dots)$  characters*


*$D_6$  symmetry and induced representation band structure*

**→ Introduction to octahedral/tetrahedral symmetry  $O_h \supset O \sim T_d \supset T$**



# *Introduction to octahedral/ tetrahedral symmetry $O_h \supset O \sim T_d \supset T$*


## *Octahedral-cubic $O$ symmetry*



*Order  $^oO = 6$  hexahedron squares  $\cdot 4$  pts  $= 24$   
 $= 8$  octahedron triangles  $\cdot 3$  pts  $= 24$   
 $= 12$  lines  $\cdot 2$  pts  $= 24$  positions*


# Introduction to octahedral/ tetrahedral symmetry $O_h \supset O \sim T_d \supset T$

## Octahedral-cubic $O$ symmetry




Order  $^oO = 6$  hexahedron squares  $\cdot 4$  pts  $= 24$   
 $= 8$  octahedron triangles  $\cdot 3$  pts  $= 24$   
 $= 12$  lines  $\cdot 2$  pts  $= 24$  positions

## Octahedral group $O$ operations





# Introduction to octahedral/ tetrahedral symmetry $O_h \supset O \sim T_d \supset T$

## Octahedral-cubic $O$ symmetry




Order  $^{\circ}O = 6$  hexahedron squares  $\cdot 4$  pts  $= 24$   
 $= 8$  octahedron triangles  $\cdot 3$  pts  $= 24$   
 $= 12$  lines  $\cdot 2$  pts  $= 24$  positions

## Octahedral group $O$ operations




# Introduction to octahedral/ tetrahedral symmetry $O_h \supset O \sim T_d \supset T$

## Octahedral-cubic $O$ symmetry





Order  $^{\circ}O = 6 \text{ hexahedron squares} \cdot 4 \text{ pts} = 24$   
 $= 8 \text{ octahedron triangles} \cdot 3 \text{ pts} = 24$   
 $= 12 \text{ lines} \cdot 2 \text{ pts} = 24 \text{ positions}$

## Octahedral group $O$ operations




## Tetrahedral symmetry becomes Icosahedral



# Introduction to octahedral tetrahedral symmetry $O_h \supset O \sim T_d \supset T$



## Octahedral groups $O_h \supset O \sim T_d \supset T$



**Figure 4.1.5** The full octahedral group ( $O_h$ ) and four non-Abelian subgroups  $T$ ,  $T_h$ ,  $T_d$ , and  $O$ . The Abelian  $D_2$  subgroup of  $T$  is indicated also.

# Introduction to octahedral tetrahedral symmetry $O_h \supset O \sim T_d \supset T$



## Octahedral groups $O_h \supset O \sim T_d \supset T$



**Figure 4.1.5** The full octahedral group ( $O_h$ ) and four non-Abelian subgroups  $T$ ,  $T_h$ ,  $T_d$ , and  $O$ . The Abelian  $D_2$  subgroup of  $T$  is indicated also.

# Introduction to octahedral tetrahedral symmetry $O_h \supset O \sim T_d \supset T$

## Octahedral groups $O_h \supset O \sim T_d \supset T$



**Figure 4.1.5** The full octahedral group ( $O_h$ ) and four non-Abelian subgroups  $T$ ,  $T_h$ ,  $T_d$ , and  $O$ . The Abelian  $D_2$  subgroup of  $T$  is indicated also.






TABLE F.2.1 *O*-Group Table

1	$r_1$	$r_2$	$r_3$	$r_4$	$r_1^2$	$r_2^2$	$r_3^2$	$r_4^2$	$R_1^2$	$R_2^2$	$R_3^2$	$R_1$	$R_2$	$R_3$	$R_1^3$	$R_2^3$	$R_3^3$	$i_1$	$i_2$	$i_3$	$i_4$	$i_5$	$i_6$
$r_1$	$r_1^2$	$-r_4^2$	$-r_2^2$	$-r_3^2$	-1	$-R_2^2$	$-R_3^2$	$-R_1^2$	$-r_2$	$-r_3$	$-r_4$	$i_3$	$i_6$	$i_1$	$-R_3$	$-R_1$	$-R_2$	$R_1^3$	$i_5$	$R_2^3$	$i_2$	$-i_4$	$R_3^3$
$r_2$	$-r_3^2$	$r_2^2$	$-r_4^2$	$-r_1^2$	$R_2^2$	-1	$R_1^2$	$-R_3^2$	$r_1$	$r_4$	$-r_3$	$R_3$	$-R_1^3$	$i_2$	$i_3$	$-i_5$	$R_2^3$	$i_6$	$-R_1$	$R_2$	$-i_1$	$R_3^3$	$i_4$
$r_3$	$-r_4^2$	$-r_1^2$	$r_3^2$	$-r_2^2$	$R_3^2$	$-R_1^2$	-1	$R_2^2$	$-r_4$	$r_1$	$r_2$	$-i_4$	$R_1$	$-R_2^3$	$R_3^3$	$i_6$	$i_2$	$i_5$	$-R_1^3$	$i_1$	$R_2$	$-i_3$	$R_3$
$r_4$	$-r_2^2$	$-r_3^2$	$-r_1^2$	$r_4^2$	$R_1^2$	$R_3^2$	$-R_2^2$	-1	$r_3$	$-r_2$	$r_1$	$-R_3^3$	$-i_5$	$R_2$	$-i_4$	$R_1^3$	$i_1$	$R_1$	$i_6$	$-i_2$	$R_2^3$	$R_3$	$i_3$
$r_1^2$	-1	$R_1^2$	$R_2^2$	$R_3^2$	$-r_1$	$r_3$	$r_4$	$r_2$	$r_4^2$	$r_2^2$	$r_3^2$	$R_2^3$	$R_3^3$	$R_1^3$	$-i_1$	$-i_3$	$-i_6$	$-R_3$	$-i_4$	$-R_1$	$i_5$	$-i_2$	$-R_2$
$r_2^2$	$-R_1^2$	-1	$R_3^2$	$-R_2^2$	$r_4$	$-r_2$	$r_1$	$r_3$	$-r_3^2$	$-r_1^2$	$r_4^2$	$i_2$	$-i_3$	$-R_1$	$R_2$	$-R_3^3$	$-i_5$	$i_4$	$-R_3$	$-R_1^3$	$-i_6$	$R_2^3$	$-i_1$
$r_3^2$	$-R_2^2$	$-R_3^2$	-1	$R_1^2$	$r_2$	$r_4$	$-r_3$	$r_1$	$r_2^2$	$-r_4^2$	$-r_1^2$	$-R_2$	$-i_4$	$-i_6$	$i_2$	$R_3$	$-R_1^3$	$-i_3$	$i_5$	$R_1$	$-i_1$	$-R_2^3$	
$r_4^2$	$-R_3^2$	$R_2^2$	$-R_1^2$	-1	$r_3$	$r_1$	$r_2$	$-r_4$	$-r_1^2$	$r_3^2$	$-r_2^2$	$-i_1$	$-R_3$	$-i_5$	$-R_2^3$	$-i_4$	$R_1$	$-R_3$	$i_3$	$-i_6$	$R_1^3$	$R_2$	$-i_2$
$R_1^2$	$-r_4$	$r_3$	$-r_2$	$r_1$	$r_2^2$	$-r_1^2$	$r_4^2$	$-r_3^2$	-1	$R_3^2$	$-R_2^2$	$R_1^3$	$i_1$	$-i_4$	$-R_1$	$i_2$	$-i_3$	$-R_2$	$-R_3^3$	$R_3^3$	$R_3$	$-i_6$	$i_5$
$R_2^2$	$-r_2$	$r_1$	$r_4$	$-r_3$	$r_3^2$	$-r_4^2$	$-r_1^2$	$r_2^2$	$-R_3^2$	-1	$R_1^2$	$-i_5$	$R_2^3$	$i_3$	$-i_6$	$-R_2$	$-i_4$	$-i_2$	$i_1$	$-R_3$	$R_3^3$	$R_1$	$R_1^3$
$R_3^2$	$-r_3$	$-r_4$	$r_1$	$r_2$	$r_4^2$	$r_3^2$	$-r_2^2$	$-r_1^2$	$R_2^2$	$-R_1^2$	-1	$i_6$	$i_2$	$R_3^3$	$-i_5$	$-i_1$	$-R_3$	$R_2^3$	$-R_2$	$i_4$	$-i_3$	$R_1^3$	$-R_1$
$R_1$	$i_1$	$-R_2^3$	$-i_2$	$R_2$	$R_3^3$	$-i_3$	$-R_3$	$i_4$	$R_1^3$	$i_6$	$i_5$	$R_2^3$	$r_1$	$-r_4^2$	-1	$-r_3$	$r_2^2$	$-r_4$	$r_2$	$r_1^2$	$-r_3^2$	$-R_2^2$	$R_3^2$
$R_2$	$i_3$	$R_3$	$-R_3^3$	$i_4$	$R_1^3$	$i_5$	$-i_6$	$-R_1$	$-i_2$	$R_2^3$	$i_1$	$-r_2^2$	$R_2^2$	$r_1$	$r_3^2$	-1	$-r_4$	$R_1^2$	$R_3^2$	$-r_2$	$-r_3$	$-r_4^2$	$r_1^2$
$R_3$	$i_6$	$i_5$	$R_1$	$-R_1^3$	$R_2^3$	$-R_2$	$-i_2$	$-i_1$	$i_3$	$i_4$	$R_3^3$	$r_1$	$-r_3^2$	$R_3^2$	$-r_2$	$r_4^2$	-1	$r_1^2$	$r_2^2$	$R_2^2$	$-R_1^2$	$-r_4$	$-r_3$
$R_1^3$	$-R_2$	$-i_2$	$R_2^3$	$i_1$	$-i_3$	$-R_3^3$	$i_4$	$R_3$	$-R_1$	$i_5$	$-i_6$	-1	$-r_4$	$r_3^2$	$-R_1^2$	$r_2$	$-r_1^2$	$r_1$	$r_3$	$r_2^2$	$-R_3^2$	$-R_2^2$	
$R_2^3$	$-R_3$	$i_3$	$i_4$	$R_3^3$	$-i_6$	$R_1$	$-R_1^3$	$i_5$	$-i_1$	$-R_2$	$-i_2$	$-i_2$	$r_4^2$	-1	$-r_2$	$-r_1^2$	$-R_2^2$	$r_3$	$-R_1^2$	$-r_4$	$r_2^2$	$r_3^2$	
$R_3^3$	$-R_1$	$R_1^3$	$i_6$	$i_5$	$-i_1$	$-i_2$	$R_2$	$-R_2^3$	$i_4$	$-i_3$	$-R_3$	$-r_3$	$r_2^2$	-1	$r_4$	$-r_1^2$	$-R_3^2$	$r_2$	$R_1^2$	$-R_2^2$	$-r_2$	$-r_1$	
$i_1$	$R_3^3$	$-i_4$	$i_3$	$R_3$	$-R_1$	$-i_6$	$-i_5$	$-R_1^3$	$R_2^3$	$i_2$	$-R_2$	$r_1^2$	$R_3^2$	$-r_4$	$r_4^2$	$-R_1^2$	$-r_1$	-1	$-R_2^2$	$-r_3$	$r_2$	$r_3^2$	$r_2^2$
$i_2$	$i_4$	$R_3^3$	$R_3$	$-i_3$	$-i_5$	$R_1^3$	$R_1$	$-i_6$	$R_2$	$-i_1$	$R_2^3$	$-r_3^2$	$-R_1^2$	$-r_3$	$-r_2^2$	$-R_3^2$	$-r_2$	$R_2^2$	-1	$r_4$	$-r_1$	$r_1^2$	$r_4^2$
$i_3$	$R_1^3$	$R_1$	$-i_5$	$i_6$	$-R_2$	$-R_2^3$	$-i_1$	$i_2$	$-R_3$	$R_3^3$	$-i_4$	$-r_2$	$r_1^2$	$R_1^2$	$-r_1$	$r_2^2$	$-R_2^2$	$r_3$	$-r_4^2$	$r_3^2$	$-R_3^2$	$r_4$	
$i_4$	$-i_5$	$i_6$	$-R_1^3$	$-R_1$	$-i_2$	$i_1$	$-R_2^3$	$-R_2$	$-R_3^3$	$-R_3$	$i_3$	$r_4$	$r_4^2$	$R_2^2$	$r_3$	$r_3^2$	$R_1^2$	$-r_2^2$	$r_1^2$	$-R_3^3$	$-1$	$r_1$	$-r_2$
$i_5$	$i_2$	$-R_2$	$i_1$	$-R_2^3$	$i_4$	$-R_3$	$i_3$	$-R_3^3$	$i_6$	$-R_1$	$R_2^3$	$R_2^2$	$r_2^2$	$R_2^2$	$r_4$	$r_4^2$	$R_1^2$	$-r_3$	$-r_2^2$	$-r_1^2$	$-1$	$-R_2^2$	
$i_6$	$R_2^3$	$i_1$	$R_2$	$i_2$	$-R_3$	$-i_4$	$-R_3^3$	$-i_3$	$-i_5$	$-R_1$	$R_1^3$	$R_2^2$	$-r_3$	$R_2^2$	$-r_1$	$R_3^2$	$r_3$	$-r_2$	$r_2^2$	$R_1^2$	$-1$		





THE 180° CLASS

$i_1 \ i_2 \ i_3 \ i_4 \ i_5 \ i_6$

THE 120° CLASS

$r_1 \ r_2 \ r_3 \ r_4$   
 $r_1^2 \ r_2^2 \ r_3^2 \ r_4^2$


THE 90° CLASS

$R_1 \ R_2 \ R_3$   
 $R_1^3 \ R_2^3 \ R_3^3$


THE 180° C

$R_1^2 \ R_2^2 \ R_3^2$

(a)



(b)



$\ell^{A_1} = 1$

$\ell^{A_2} = 1$

$\ell^E = 2$

$\ell^{T_1} = 3$

$\ell^{T_2} = 3$

*Example: G=O Centrum:  $\kappa(O) = \Sigma_{(\alpha)} (\ell^\alpha)^0 = 1^0 + 1^0 + 2^0 + 3^0 + 3^0 = 5$*

*Cubic-Octahedral Group O*

$\text{Rank: } \rho(O) = \Sigma_{(\alpha)} (\ell^\alpha)^1 = 1^1 + 1^1 + 2^1 + 3^1 + 3^1 = 10$


$\text{Order: } o(O) = \Sigma_{(\alpha)} (\ell^\alpha)^0 = 1^2 + 1^2 + 2^2 + 3^2 + 3^2 = 24$

<i>O group</i> $\chi_{\kappa_g}^\alpha$	$g = 1$	$r_{1-4}$ $\tilde{r}_{1-4}$	$\rho_{xyz}$	$R_{xyz}$ $\tilde{R}_{xyz}$	$i_{1-6}$
<i>s-orbital <math>r^2</math></i> $\rightarrow \alpha = A_1$	1	1	1	1	1
<i>d-orbitals</i> $\{x^2+y^2-2z^2, x^2-y^2\}$ $\rightarrow A_2$	1	1	1	-1	-1
<i>p-orbitals <math>\{x, y, z\}</math></i> $\rightarrow E$	2	-1	2	0	0
<i>p-orbitals <math>\{x, y, z\}</math></i> $\rightarrow T_1$	3	0	-1	1	-1
<i>d-orbitals</i> $\{xz, yz, xy\}$ $\rightarrow T_2$	3	0	-1	-1	1


$O \supset C_4 (0)_4 (1)_4 (2)_4 (3)_4 = (-1)_4$ 
 $O \supset C_3 (0)_3 (1)_3 (2)_3 = (-1)_3$

A <sub>1</sub>	1	•	•	•
A <sub>2</sub>	•	•	1	•
E	1	•	1	•
T <sub>1</sub>	1	1	•	1
T <sub>2</sub>	•	1	1	1

A <sub>1</sub>	1	•	•
A <sub>2</sub>	1	•	•
E	•	1	1
T <sub>1</sub>	1	1	1
T <sub>2</sub>	1	1	1



(a)  $SF_6 \nu_4$  Rotational Structure




FT IR and Laser Diode Spectra  
K.C. Kim, W.B. Person, D. Seitz, and B.J. Krohn  
*J. Mol. Spectrosc.* **76**, 322 (1979).


Primary AET species mixing  
increases with distance from  
"separatrix"

(b) P(88) Fine Structure (Rotational anisotropy effects)

$SF_6 \nu_3 P(88) \sim 16m$



(c) Superfine Structure (Rotational axis tunneling)



Observed repeating sequence(s)...  $A_1 T_1 E T_2 T_1 ET_2 A_2 T_2 T_1 A_1 T_1 ET_2 T_1 ET_2 A_2 T_2 T_1 A_1 \dots$

$$O \supset C_4 (0)_4 (1)_4 (2)_4 (3)_4 = (-1)_4$$

	$A_1$	$\cdot$	$\cdot$	$\cdot$
$A_2$	$\cdot$	$\cdot$	$1$	$\cdot$
$E$	$1$	$\cdot$	$1$	$\cdot$
$T_1$	$1$	$1$	$\cdot$	$1$
$T_2$	$\cdot$	$1$	$1$	$1$

$$O \supset C_3 (0)_3 (1)_3 (2)_3 = (-1)_3$$

	$A_1$	$1$	$\cdot$	$\cdot$
$A_2$	$1$	$\cdot$	$\cdot$	$\cdot$
$E$	$\cdot$	$1$	$1$	$\cdot$
$T_1$	$1$	$1$	$1$	$1$
$T_2$	$1$	$1$	$1$	$1$

Local correlations explain clustering...

... but what about spacing and ordering?...

...and physical consequences?









Deriving  $D_3 \sim C_{3v}$  products - By group definition  $|g\rangle = g|1\rangle$  of position ket  $|g\rangle$

