

Group Theory in Quantum Mechanics

Lecture 20 (4.12.17)

Octahedral-tetrahedral $O \sim T_d$ representations and spectra

(Int.J.Mol.Sci, 14, 714(2013) p.755-774 , QTCA Unit 5 Ch. 15)

(PSDS - Ch. 4)

Review Octahedral $O_h \supset O$ group operator structure

Review Octahedral $O_h \supset O \supset D_4 \supset C_4$ subgroup chain correlations

Comparison of $O \supset D_4 \supset C_4$ and $O \supset D_4 \supset D_2$ correlations and level/projector splitting

$O \supset D_4 \supset C_4$ subgroup chain splitting

$O \supset D_4 \supset D_2$ subgroup chain splitting (normal D_2 vs. unnormal D_2)

$O_h \supset O \supset D_4 \supset C_{4v}$ and $O_h \supset O \supset D_4 \supset C_{4v} \supset C_{2v}$ subgroup splitting

Splitting O class projectors \mathbf{P}^μ into irreducible projectors $\mathbf{P}^\mu_{m_4 m_4}$ for $O \supset C_4$

Development of irreducible projectors $\mathbf{P}^\mu_{m_4 m_4}$ and representations $D^\mu_{m_4 m_4}$

Calculating $\mathbf{P}^E_{0_4 0_4}$, $\mathbf{P}^E_{2_4 2_4}$, $\mathbf{P}^{T_1}_{0_4 0_4}$, $\mathbf{P}^{T_1}_{1_4 1_4}$, $\mathbf{P}^{T_2}_{2_4 2_4}$, $\mathbf{P}^{T_2}_{1_4 1_4}$,


$O \supset C_4$ induced representation $0_4(C_4) \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples

Elementary induced representation $0_4(C_4) \uparrow O$


Projection reduction of induced representation $0_4(C_4) \uparrow O$

Introduction to ortho-complete eigenvalue-parameter relations

Examples in SF_6 spectroscopy



Review Octahedral $O_h \supset O$ group operator structure



Review Octahedral $O_h \supset O \supset D_4 \supset C_4$ subgroup chain correlations

Comparison of $O \supset D_4 \supset C_4$ and $O \supset D_4 \supset D_2$ correlations and level/projector splitting

$O \supset D_4 \supset C_4$ subgroup chain splitting

$O \supset D_4 \supset D_2$ subgroup chain splitting (normal D_2 vs. unnormal D_2)

$O_h \supset O \supset D_4 \supset C_{4v}$ and $O_h \supset O \supset D_4 \supset C_{4v} \supset C_{2v}$ subgroup splitting

Splitting O class projectors P^μ into irreducible projectors $P^\mu_{m+m_4}$ for $O \supset C_4$

Development of irreducible projectors $P^\mu_{m+m_4}$ and representations $D^\mu_{m+m_4}$

Calculating $P^{E_{0404}}$, $P^{E_{2424}}$, $P^{T_{10404}}$, $P^{T_{11414}}$, $P^{T_{22424}}$, $P^{T_{21414}}$,

$O \supset C_4$ induced representation $0_4(C_4) \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples

Elementary induced representation $0_4(C_4) \uparrow O$


Projection reduction of induced representation $0_4(C_4) \uparrow O$

Introduction to ortho-complete eigenvalue-parameter relations

Examples in SF_6 spectroscopy

Introduction to octahedral/ tetrahedral symmetry $O_h \supset O \sim T_d \supset T$

Octahedral-cubic O symmetry




Order $^{\circ}O = 6 \text{ hexahedron squares} \cdot 4 \text{ pts} = 24$
 $= 8 \text{ octahedron triangles} \cdot 3 \text{ pts} = 24$
 $= 12 \text{ lines} \cdot 2 \text{ pts} = 24 \text{ positions}$

Octahedral group O operations


Class of 1: **1**

$$\mathbf{r}_k = \mathbf{r}_k$$

Class of 8:
 $\pm 120^{\circ}$ rotations
 on $[111]$ axes



$$\tilde{\mathbf{r}}_k = \mathbf{r}_k^2 = \mathbf{r}_k^{-1}$$




Class of 3:
 180° rotations
 on $[100]$ axes

$$\rho_{x,y,z} = \mathbf{R}_{1,2,3}^2$$


$$\mathbf{R}_{x,y,z} = \mathbf{R}_{1,2,3}$$

Class of 6:
 $\pm 90^{\circ}$ rotations
 on $[100]$ axes




Class of 6:
 180° rotations
 on $[110]$ diagonals

$$\mathbf{i}_k = \mathbf{i}_k$$



$$\tilde{\mathbf{R}}_{x,y,z} = \mathbf{R}_{1,2,3}^3 = \mathbf{R}_{1,2,3}^{-1}$$




Tetrahedral symmetry becomes Icosahedral

T symmetry

T_h symmetry

I_h symmetry

(If rectangles have
 Golden Ratio $\frac{1 \pm \sqrt{5}}{2}$)



Introduction to octahedral tetrahedral symmetry $O_h \supset O \sim T_d \supset T$

Octahedral groups $O_h \supset O \sim T_d$ and $O_h \supset T_h \supset T$




Figure 4.1.5 The full octahedral group (O_h) and four non-Abelian subgroups T , T_h , T_d , and O . The Abelian D_2 subgroup of T is indicated also.

Fig. 4.1.5 from *Principles of Symmetry, Dynamics and Spectroscopy*

Review Octahedral $O_h \supset O$ group operator structure

 *Review Octahedral $O_h \supset O \supset D_4 \supset C_4$ subgroup chain correlations* 

Comparison of $O \supset D_4 \supset C_4$ and $O \supset D_4 \supset D_2$ correlations and level/projector splitting

$O \supset D_4 \supset C_4$ subgroup chain splitting

$O \supset D_4 \supset D_2$ subgroup chain splitting (normal D_2 vs. unnormal D_2)

$O_h \supset O \supset D_4 \supset C_{4v}$ and $O_h \supset O \supset D_4 \supset C_{4v} \supset C_{2v}$ subgroup splitting

Splitting O class projectors P^μ into irreducible projectors $P^\mu_{m+m_4}$ for $O \supset C_4$

Development of irreducible projectors $P^\mu_{m+m_4}$ and representations $D^\mu_{m+m_4}$

Calculating $P^{E_{0404}}$, $P^{E_{2424}}$, $P^{T_{10404}}$, $P^{T_{11414}}$, $P^{T_{22424}}$, $P^{T_{21414}}$,

$O \supset C_4$ induced representation $0_4(C_4) \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples

Elementary induced representation $0_4(C_4) \uparrow O$

Projection reduction of induced representation $0_4(C_4) \uparrow O$

Introduction to ortho-complete eigenvalue-parameter relations

Examples in SF_6 spectroscopy

Octahedral $O_h \supset O \supset D_4 \supset C_4$ subgroup correlations


$O \downarrow D_4$ subduction

$\chi_g^\mu(O)$	$g=1$	$r_{1..4}$	180° ρ_{xyz}	90° R_{xyz}	180° $i_{1..6}$
A_1	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

$D_4: \mathbf{1}, \rho_{z180^\circ}, R_{z\pm 90^\circ}, \rho_{x,y180^\circ}, i_{3,4}$

$A_1(O) \downarrow D_4 = 1, 1, 1, 1, 1.$
 $A_2(O) \downarrow D_4 = 1, 1, -1, 1, -1.$
 $E(O) \downarrow D_4 = 2, 2, 0, 2, 0.$
 $T_2(O) \downarrow D_4 = 3, -1, 1, -1, -1.$
 $T_2(O) \downarrow D_4 = 3, -1, -1, -1, 1.$

Fig. 3.1.1 PSDS



$\chi_g^\mu(D_4)$	$g=1$	ρ_{z180°	$R_{z\pm 90^\circ}$	$\rho_{x,y180^\circ}$	$i_{3,4}$
A_1	1	1	1	1	1
B_1	1	1	-1	1	-1
A_2	1	1	1	-1	-1
B_2	1	1	-1	-1	1
E	2	-2	0	0	0

$D_4 \downarrow C_4$ subduction

$C_4: \mathbf{1}, R_{z+90^\circ}, \rho_{z180^\circ}, R_{z-90^\circ}$
 $A_1(D_4) \downarrow C_4 = 1, 1, 1, 1. = (0)_4$
 $B_1(D_4) \downarrow C_4 = 1, -1, 1, -1. = (2)_4$
 $A_2(D_4) \downarrow C_4 = 1, 1, 1, 1. = (0)_4$
 $B_2(D_4) \downarrow C_4 = 1, -1, 1, -1. = (2)_4$
 $E(D_4) \downarrow C_4 = 2, 0, -2, 0. = (1)_4 \oplus (3)_4$


$O \downarrow D_4$	A_1	B_1	A_2	B_2	E
A_1	1
A_2	.	1	.	.	.
E	1	1	.	.	.
T_1	.	.	1	.	1
T_2	.	.	.	1	1

$\chi_g^\mu(C_4)$	$g=1$	R_{z+90°	R_{z+180°	R_{z-90°
$(0)_4$	1	1	1	1
$(1)_4$	1	i	-1	$-i$
$(2)_4$	1	-1	1	-1
$(3)_4$	1	$-i$	-1	i

$O \downarrow C_4$ subduction

$O \downarrow C_4$	0_4	1_4	2_4	$3_4 = \bar{1}_4$
A_1	1	.	.	.
A_2	.	.	1	.
E	1	.	1	.
T_1	1	1	.	1
T_2	.	$\rightarrow 1$	$\rightarrow 1$	$\rightarrow 1$

$D_4 \downarrow C_4$	0_4	1_4	2_4	$3_4 = \bar{1}_4$
A_1	1	.	.	.
B_1	.	.	1	.
A_2	1	.	.	.
B_2	.	.	$\rightarrow 1$.
E	.	$\rightarrow 1$.	$\rightarrow 1$




Octahedral $O_h \supset O \supset D_4 \supset C_4$ subgroup correlations

$O \downarrow D_4$ subduction

D_4 : $\mathbf{1}, \rho_{z180^\circ}, \mathbf{R}_{z\pm 90^\circ}, \rho_{x,y180^\circ}, \mathbf{i}_{3,4}$


$$\begin{aligned}
 A_1(O) \downarrow D_4 &= 1, 1, 1, 1, 1. \\
 A_2(O) \downarrow D_4 &= 1, 1, -1, 1, -1. \\
 E(O) \downarrow D_4 &= 2, 2, 0, 2, 0. \\
 T_2(O) \downarrow D_4 &= 3, -1, 1, -1, -1. \\
 T_2(O) \downarrow D_4 &= 3, -1, -1, -1, 1.
 \end{aligned}$$

Fig. 3.1.1 PSDS



$O \supset D_4 \supset C_4$ subgroup and level-splitting/relabeling correlations

O levels \downarrow D_4 levels \downarrow C_4 levels



$D_4 \downarrow C_4$ subduction

C_4 : $\mathbf{1}, \mathbf{R}_{z+90^\circ}, \rho_{z180^\circ}, \mathbf{R}_{z-90^\circ}$


$$\begin{aligned}
 A_1(D_4) \downarrow C_4 &= 1, 1, 1, 1. = (0)_4 \\
 B_1(D_4) \downarrow C_4 &= 1, -1, 1, -1. = (2)_4 \\
 A_2(D_4) \downarrow C_4 &= 1, 1, 1, 1. = (0)_4 \\
 B_2(D_4) \downarrow C_4 &= 1, -1, 1, -1. = (2)_4 \\
 E(D_4) \downarrow C_4 &= 2, 0, -2, 0. = (1)_4 \oplus (3)_4
 \end{aligned}$$

$O \downarrow D_4$	A_1	B_1	A_2	B_2	E
A_1	1
A_2	.	1	.	.	.
E	1	1	.	.	.
T_1	.	.	1	.	1
T_2	.	.	.	1	1

$O \downarrow C_4$ subduction

$O \downarrow C_4$	0_4	1_4	2_4	$3_4 = \bar{1}_4$
A_1	1	.	.	.
A_2	.	.	1	.
E	1	.	1	.
T_1	1	1	.	1
T_2	.	$\rightarrow 1$	$\rightarrow 1$	$\rightarrow 1$

$D_4 \downarrow C_4$	0_4	1_4	2_4	$3_4 = \bar{1}_4$
A_1	1	.	.	.
B_1	.	.	1	.
A_2	1	.	.	.
B_2	.	.	$\rightarrow 1$.
E	.	$\rightarrow 1$.	$\rightarrow 1$



Review Octahedral $O_h \supset O$ group operator structure

Review Octahedral $O_h \supset O \supset D_4 \supset C_4$ subgroup chain correlations

→ *Comparison of $O \supset D_4 \supset C_4$ and $O \supset D_4 \supset D_2$ correlations and level/projector splitting* **←**

$O \supset D_4 \supset C_4$ subgroup chain splitting

$O \supset D_4 \supset D_2$ subgroup chain splitting (normal D_2 vs. unnormal D_2)

$O_h \supset O \supset D_4 \supset C_{4v}$ and $O_h \supset O \supset D_4 \supset C_{4v} \supset C_{2v}$ subgroup splitting

Splitting O class projectors P^μ into irreducible projectors $P^\mu_{m_4 m_4}$ for $O \supset C_4$

Development of irreducible projectors $P^\mu_{m_4 m_4}$ and representations $D^\mu_{m_4 m_4}$

Calculating $P^{E_{0_4 0_4}}$, $P^{E_{2_4 2_4}}$, $P^{T_{1_4 0_4}}$, $P^{T_{1_4 1_4}}$, $P^{T_{2_4 2_4}}$, $P^{T_{2_4 1_4}}$,

$O \supset C_4$ induced representation $0_4(C_4) \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples

Elementary induced representation $0_4(C_4) \uparrow O$

Projection reduction of induced representation $0_4(C_4) \uparrow O$


Introduction to ortho-complete eigenvalue-parameter relations

Examples in SF_6 spectroscopy

D_4	$\mathbf{1}$	ρ_z	\mathbf{R}_z	$\rho_{x,y}$	$\mathbf{i}_{3,4}$
A_1	1	1	1	1	1
B_1	1	1	-1	1	-1
A_2	1	1	1	-1	-1
B_2	1	1	-1	-1	1
E	2	-2	0	0	0

$O \supset D_4 \supset C_4$ level splitting

Tetragonal Moving Wave Chain



$C_4 \{ \mathbf{1}, \mathbf{R}_z^1, \mathbf{R}_z^2, \mathbf{R}_z^3 \}$
 $\{ \mathbf{1}, \mathbf{R}_3^2, \mathbf{R}_3^3, \mathbf{R}_3^3 \}$

C_4	$\mathbf{1}$	\mathbf{R}_z^1	\mathbf{R}_z^2	\mathbf{R}_z^3
0_4	1	1	1	1
1_4	1	i	-1	$-i$
2_4	1	-1	1	-1
3_4	1	$-i$	-1	i

$-1_4 =$


$D_4 \downarrow C_4$	0_4	1_4	2_4	3_4
A_1	1	.	.	.
B_1	.	.	1	.
A_2	1	.	.	.
B_2	.	.	1	.
E	.	1	.	1

O	$\mathbf{1}$	$\mathbf{r}, \tilde{\mathbf{r}}_i$	ρ_{xyz}	$\mathbf{R}, \tilde{\mathbf{R}}_{xyz}$	\mathbf{R}^2	\mathbf{R}^3	\mathbf{i}_k
A_1	1	1	1	1	1	1	1
A_2	1	1	1	-1	-1	-1	-1
E	2	-1	2	0	0	0	0
T_1	3	0	-1	1	-1	-1	-1
T_2	3	0	-1	-1	-1	1	1

$O \downarrow D_4$	A_1	B_1	A_2	B_2	E
A_1	1
A_2	.	1	.	.	.
E	1	1	.	.	.
T_1	.	.	1	.	1
T_2	.	.	.	1	1

$O \downarrow C_4$	0_4	1_4	2_4	$3_4 = \bar{1}_4$
A_1	1	.	.	.
A_2	.	.	1	.
E	1	.	1	.
T_1	1	1	.	1
T_2	.	1	1	1

$O \supset D_4 \supset D_2$ level splitting




D_4	1	ρ_z	R_z	$\rho_{x,y}$	$i_{3,4}$
A_1	1	1	1	1	1
B_1	1	1	-1	1	-1
A_2	1	1	1	-1	-1
B_2	1	1	-1	-1	1
E	2	-2	0	0	0

Normal $D_2 = \{1, R_3^2, R_1^2, R_2^2\}$

$D_4 \downarrow D_2$	A_1	B_1	A_2	B_2
A_1	1	.	.	.
B_1	1	.	.	.
A_2	.	.	1	.
B_2	.	.	1	.
E	.	1	.	1

$O \supset D_4 \supset C_4$ level splitting



$D_2^{Nm} \{1, R_z^2, R_x^2, R_y^2\}$

A_1	1	1	1	1
B_1	1	-1	1	-1
A_2	1	1	-1	-1
B_2	1	-1	-1	1

$-1_4 =$

$D_4 \downarrow C_4$	0_4	1_4	2_4	3_4
A_1	1	.	.	.
B_1	.	.	1	.
A_2	1	.	.	.
B_2	.	.	1	.
E	.	1	.	1

	$r, \tilde{r}_i \quad \rho_{xyz} \quad R, \tilde{R}_{xyz}$				
O	1	r	R^2	R^3	i_k
A_1	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

Normal $D_2 = \{1, R_3^2, R_1^2, R_2^2\}$

$O \downarrow D_2$	A_1	B_1	A_2	B_2
A_1	1	.	.	.
A_2	1	.	.	.
E	2	.	.	.
T_1	.	1	1	1
T_2	.	1	1	1

$O \downarrow D_4$	A_1	B_1	A_2	B_2	E
A_1	1
A_2	.	1	.	.	.
E	1	1	.	.	.
T_1	.	.	1	.	1
T_2	.	.	.	1	1

$O \downarrow C_4$	0_4	1_4	2_4	$3_4 = \bar{1}_4$
A_1	1	.	.	.
A_2	.	.	1	.
E	1	.	1	.
T_1	1	1	.	1
T_2	.	1	1	1

Review Octahedral $O_h \supset O$ group operator structure

Review Octahedral $O_h \supset O \supset D_4 \supset C_4$ subgroup chain correlations

Comparison of $O \supset D_4 \supset C_4$ and $O \supset D_4 \supset D_2$ correlations and level/projector splitting

$O \supset D_4 \supset C_4$ subgroup chain splitting

$O \supset D_4 \supset D_2$ subgroup chain splitting (normal D_2 vs. unnormal D_2)

$O_h \supset O \supset D_4 \supset C_{4v}$ and $O_h \supset O \supset D_4 \supset C_{4v} \supset C_{2v}$ subgroup splitting

Splitting O class projectors P^μ into irreducible projectors $P^\mu_{m_4m_4}$ for $O \supset C_4$

Development of irreducible projectors $P^\mu_{m_4m_4}$ and representations $D^\mu_{m_4m_4}$

Calculating $P^{E_{0404}}$, $P^{E_{2424}}$, $P^{T_{10404}}$, $P^{T_{11414}}$, $P^{T_{22424}}$, $P^{T_{21414}}$,



$O \supset C_4$ induced representation $0_4(C_4) \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples

Elementary induced representation $0_4(C_4) \uparrow O$

Projection reduction of induced representation $0_4(C_4) \uparrow O$

Introduction to ortho-complete eigenvalue-parameter relations

Examples in SF_6 spectroscopy



$O_h \supset O \supset D_4 \supset D_2$ subgroup splitting

D_4	1	ρ_z	R_z	$\rho_{x,y}$	$i_{3,4}$
A_1	1	1	1	1	1
B_1	1	1	-1	1	-1
A_2	1	1	1	-1	-1
B_2	1	1	-1	-1	1
E	2	-2	0	0	0






Normal $D_2 = \{1, R_3^2, R_1^2, R_2^2\}$

$D_4 \downarrow D_2$	A_1	B_1	A_2	B_2
A_1	1	.	.	.
B_1	1	.	.	.
A_2	.	.	1	.
B_2	.	.	1	.
E	.	1	.	1

UnNormal $D_2 = \{1, R_3^2, i_3, i_4\}$

$D_4 \downarrow D_2$	A_1	B_1	A_2	B_2
A_1	1	.	.	.
B_1	.	.	1	.
A_2	.	.	1	.
B_2	1	.	.	.
E	.	1	.	1

Tetragonal Moving Wave Chain



$D_2^{Nm} \{1, R_z^2, R_x^2, R_y^2\}$






$D_2^{Un} \{1, R_z^2, i_3, i_4\}$

A_1	1	1	1	1
B_1	1	-1	1	-1
A_2	1	1	-1	-1
B_2	1	-1	-1	1

$-1_4 =$

$D_4 \downarrow C_4$	0_4	1_4	2_4	3_4
A_1	1	.	.	.
B_1	.	.	1	.
A_2	1	.	.	.
B_2	.	.	1	.
E	.	1	.	1

	$r, \tilde{r}_i \quad \rho_{xyz} \quad R, \tilde{R}_{xyz}$				
O	1	r	R^2	R^3	i_k
A_1	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1



two kinds of D_2 subgroup

Normal $D_2 = \{1, R_3^2, R_1^2, R_2^2\}$



$O \downarrow D_2$	A_1	B_1	A_2	B_2
A_1	1	.	.	.
A_2	1	.	.	.
E	2	.	.	.
T_1	.	1	1	1
T_2	.	1	1	1

$O \downarrow D_4$	A_1	B_1	A_2	B_2	E
A_1	1
A_2	.	1	.	.	.
E	1	1	.	.	.
T_1	.	.	1	.	1
T_2	.	.	.	1	1

$O \downarrow C_4$	0_4	1_4	2_4	$3_4 = \bar{1}_4$
A_1	1	.	.	.
A_2	.	.	1	.
E	1	.	1	.
T_1	1	1	.	1
T_2	.	1	1	1

$O_h \supset O \supset D_4 \supset D_2$ subgroup splitting

D_4	1	ρ_z	R_z	$\rho_{x,y}$	$i_{3,4}$
A_1	1	1	1	1	1
B_1	1	1	-1	1	-1
A_2	1	1	1	-1	-1
B_2	1	1	-1	-1	1
E	2	-2	0	0	0




$D_2^{Nm} \{ 1, R_z^2, R_x^2, R_y^2 \}$

$D_2^{Un} \{ 1, R_z^2, i_3, i_4 \}$

A_1	1	1	1	1
B_1	1	-1	1	-1
A_2	1	1	-1	-1
B_2	1	-1	-1	1

$-1_4 =$




Normal $D_2 = \{ 1, R_3^2, R_1^2, R_2^2 \}$

$D_4 \downarrow D_2$	A_1	B_1	A_2	B_2
A_1	1	.	.	.
B_1	1	.	.	.
A_2	.	.	1	.
B_2	.	.	1	.
E	.	1	.	1

UnNormal $D_2 = \{ 1, R_3^2, i_3, i_4 \}$

$D_4 \downarrow D_2$	A_1	B_1	A_2	B_2
A_1	1	.	.	.
B_1	.	.	1	.
A_2	.	.	1	.
B_2	1	.	.	.
E	.	1	.	1



$D_4 \downarrow C_4$	0_4	1_4	2_4	3_4
A_1	1	.	.	.
B_1	.	.	1	.
A_2	1	.	.	.
B_2	.	.	1	.
E	.	1	.	1

	1	r, \tilde{r}_i	ρ_{xyz}	R, \tilde{R}_{xyz}	R^2	R^3	i_k
O	1	r	ρ_{xyz}	R, \tilde{R}_{xyz}	R^2	R^3	i_k
A_1	1	1	1	1	1	1	1
A_2	1	1	1	-1	-1	-1	-1
E	2	-1	2	0	0	0	0
T_1	3	0	-1	1	-1	-1	-1
T_2	3	0	-1	-1	1	1	1

two kinds of D_2 subgroup splitting

Normal $D_2 = \{ 1, R_3^2, R_1^2, R_2^2 \}$ *UnNormal* $D_2 = \{ 1, R_3^2, i_3, i_4 \}$

$O \downarrow D_2$	A_1	B_1	A_2	B_2
A_1	1	.	.	.
A_2	1	.	.	.
E	2	<i>degeneracy ambiguity</i>		.
T_1	.	1	1	1
T_2	.	1	1	1

$O \downarrow D_2$	A_1	B_1	A_2	B_2
A_1	1	.	.	.
A_2	.	.	1	.
E	1	.	1	.
T_1	.	1	1	1
T_2	1	1	.	1

$O \downarrow D_4$	A_1	B_1	A_2	B_2	E
A_1	1
A_2	.	1	.	.	.
E	1	1	.	.	.
T_1	.	.	1	.	1
T_2	.	.	.	1	1

$O \downarrow C_4$	0_4	1_4	2_4	$3_4 = \bar{1}_4$
A_1	1	.	.	.
A_2	.	.	1	.
E	1	.	1	.
T_1	1	1	.	1
T_2	.	1	1	1

Review Octahedral $O_h \supset O$ group operator structure

Review Octahedral $O_h \supset O \supset D_4 \supset C_4$ subgroup chain correlations

Comparison of $O \supset D_4 \supset C_4$ and $O \supset D_4 \supset D_2$ correlations and level/projector splitting

$O \supset D_4 \supset C_4$ subgroup chain splitting

$O \supset D_4 \supset D_2$ subgroup chain splitting (normal D_2 vs. unnormal D_2)

$O_h \supset O \supset D_4 \supset C_{4v}$ and $O_h \supset O \supset D_4 \supset C_{4v} \supset C_{2v}$ subgroup splitting

Splitting O class projectors P^μ into irreducible projectors $P^\mu_{m_4m_4}$ for $O \supset C_4$

Development of irreducible projectors $P^\mu_{m_4m_4}$ and representations $D^\mu_{m_4m_4}$

Calculating $P^{E_{0404}}$, $P^{E_{2424}}$, $P^{T_{10404}}$, $P^{T_{11414}}$, $P^{T_{22424}}$, $P^{T_{21414}}$,

$O \supset C_4$ induced representation $0_4(C_4) \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples

Elementary induced representation $0_4(C_4) \uparrow O$

Projection reduction of induced representation $0_4(C_4) \uparrow O$


Introduction to ortho-complete eigenvalue-parameter relations

Examples in SF_6 spectroscopy

$O_h \supset D_{4h} \supset C_{4v} \supset C_4$ subgroup splitting

$O_h \supset C_{4v}$	A'	B'	A''	B''	E
$A_{1g} \downarrow C_{4v}$	1	·	·	·	·
$A_{2g} \downarrow C_{4v}$	·	1	·	·	·
$E_g \downarrow C_{4v}$	1	1	·	·	·
$T_{1g} \downarrow C_{4v}$	·	·	1	·	1
$T_{2g} \downarrow C_{4v}$	·	·	·	1	1
$A_{1g} \downarrow C_{4v}$	·	·	1	·	·
$A_{2u} \downarrow C_{4v}$	·	·	·	1	·
$E_u \downarrow C_{4v}$	·	·	1	1	·
$T_{1u} \downarrow C_{4v}$	1	·	·	·	1
$T_{2u} \downarrow C_{4v}$	·	1	·	·	1

Fig. 3.1.1 PSDS



$O_h \supset D_{4h} \supset C_{4v} \supset C_{2v}$ subgroup splitting

$O_h \supset C_{4v}$	A'	B'	A''	B''	E
$A_{1g} \downarrow C_{4v}$	1	·	·	·	·
$A_{2g} \downarrow C_{4v}$	·	1	·	·	·
$E_g \downarrow C_{4v}$	1	1	·	·	·
$T_{1g} \downarrow C_{4v}$	·	·	1	·	1
$T_{2g} \downarrow C_{4v}$	·	·	·	1	1
$A_{1g} \downarrow C_{4v}$	·	·	1	·	·
$A_{2u} \downarrow C_{4v}$	·	·	·	1	·
$E_u \downarrow C_{4v}$	·	·	1	1	·
$T_{1u} \downarrow C_{4v}$	1	·	·	·	1
$T_{2u} \downarrow C_{4v}$	·	1	·	·	1


has degeneracy ambiguity

$O_h \supset C_{2v}^z$	A'	B'	A''	B''
$A_{1g} \downarrow C_{2v}^z$	1	·	·	·
$A_{2g} \downarrow C_{2v}^z$	1	·	·	·
$E_g \downarrow C_{2v}^z$	2	·	·	·
$T_{1g} \downarrow C_{2v}^z$	·	1	1	1
$T_{2g} \downarrow C_{2v}^z$	·	1	1	1
$A_{1g} \downarrow C_{2v}^z$	·	·	1	·
$A_{2u} \downarrow C_{2v}^z$	·	·	1	·
$E_u \downarrow C_{2v}^z$	·	·	2	·
$T_{1u} \downarrow C_{2v}^z$	1	1	·	1
$T_{2u} \downarrow C_{2v}^z$	1	1	·	1

has no degeneracy ambiguity

$O_h \supset C_{2v}^i$	A'	B'	A''	B''
$A_{1g} \downarrow C_{2v}^i$	1	·	·	·
$A_{2g} \downarrow C_{2v}^i$	·	1	·	·
$E_g \downarrow C_{2v}^i$	1	1	·	·
$T_{1g} \downarrow C_{2v}^i$	·	1	1	1
$T_{2g} \downarrow C_{2v}^i$	1	·	1	1
$A_{1g} \downarrow C_{2v}^i$	·	·	1	·
$A_{2u} \downarrow C_{2v}^i$	·	·	·	1
$E_u \downarrow C_{2v}^i$	·	·	1	1
$T_{1u} \downarrow C_{2v}^i$	1	1	·	1
$T_{2u} \downarrow C_{2v}^i$	1	1	1	·

Fig. 3.1.1 PSDS



Review Octahedral $O_h \supset O$ group operator structure


Review Octahedral $O_h \supset O \supset D_4 \supset C_4$ subgroup chain correlations

Comparison of $O \supset D_4 \supset C_4$ and $O \supset D_4 \supset D_2$ correlations and level/projector splitting

$O \supset D_4 \supset C_4$ subgroup chain splitting

$O \supset D_4 \supset D_2$ subgroup chain splitting (normal D_2 vs. unnormal D_2)


$O_h \supset O \supset D_4 \supset C_{4v}$ and $O_h \supset O \supset D_4 \supset C_{4v} \supset C_{2v}$ subgroup splitting



Splitting O class projectors P^μ into irreducible projectors $P^\mu_{m_4 m_4}$ for $O \supset C_4$

Development of irreducible projectors $P^\mu_{m_4 m_4}$ and representations $D^\mu_{m_4 m_4}$

Calculating $P^{E_{0404}}$, $P^{E_{2424}}$, $P^{T_{10404}}$, $P^{T_{11414}}$, $P^{T_{22424}}$, $P^{T_{21414}}$,



$O \supset C_4$ induced representation $0_4(C_4) \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples

Elementary induced representation $0_4(C_4) \uparrow O$

Projection reduction of induced representation $0_4(C_4) \uparrow O$

Introduction to ortho-complete eigenvalue-parameter relations

Examples in SF_6 spectroscopy

Splitting O class projectors \mathbf{P}^μ into irreducible projectors $\mathbf{P}^\mu_{m_4 m_4}$ for $O \supset C_4$

$O \supset C_4$ Correlation table shows which \mathbf{P}^μ splittings are allowed:

$O \supset C_4$	0_4	1_4	2_4	3_4
$A_1 \downarrow C_4$	1	.	.	.
$A_2 \downarrow C_4$.	.	1	.
$E \downarrow C_4$	1	.	1	.
$T_1 \downarrow C_4$	1	1	.	1
$T_2 \downarrow C_4$.	1	1	1

$\mathbf{P}^{A_1} = \mathbf{P}^{A_1}_{0_4 0_4}$
 and
 $\mathbf{P}^{A_2} = \mathbf{P}^{A_2}_{2_4 2_4}$
 cannot split

$O: \chi_g^\mu$	O characters				
	$g=1$	\mathbf{r}_{1-4} $\tilde{\mathbf{r}}_{1-4}$	ρ_{xyz}	\mathbf{R}_{xyz} $\tilde{\mathbf{R}}_{xyz}$	\mathbf{i}_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

$C_4: d_{\mathbf{R}^p}^{m_4}$	C_4 characters			
	$g=1$	\mathbf{R}_z^1	$\rho_z = \mathbf{R}_z^2$	$\tilde{\mathbf{R}}_z = \mathbf{R}_z^3$
$m_4=0_4$	1	1	1	1
1_4	1	-i	-1	i
2_4	1	-1	1	-1
3_4	1	-i	-1	-i

O operators (Two notations: Older Princ. of Symm. Dynamics and Spectra. and Newer Int. J. Mol. Sci)

PSDS:	1	\mathbf{r}_1	\mathbf{r}_2	\mathbf{r}_3	\mathbf{r}_4	\mathbf{r}_1^2	\mathbf{r}_2^2	\mathbf{r}_3^2	\mathbf{r}_4^2	\mathbf{R}_1^2	\mathbf{R}_2^2	\mathbf{R}_3^2	\mathbf{R}_1	\mathbf{R}_2	\mathbf{R}_3	\mathbf{R}_1^3	\mathbf{R}_2^3	\mathbf{R}_3^3	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6
IJMS:	1	\mathbf{r}_1	\mathbf{r}_2	\mathbf{r}_3	\mathbf{r}_4	$\tilde{\mathbf{r}}_1$	$\tilde{\mathbf{r}}_2$	$\tilde{\mathbf{r}}_3$	$\tilde{\mathbf{r}}_4$	ρ_x	ρ_y	ρ_z	\mathbf{R}_x	\mathbf{R}_y	\mathbf{R}_z	$\tilde{\mathbf{R}}_x$	$\tilde{\mathbf{R}}_y$	$\tilde{\mathbf{R}}_z$	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6

Splitting O class projectors \mathbf{P}^μ into irreducible projectors $\mathbf{P}^\mu_{m_4 m_4}$ for $O \supset C_4$

$O \supset C_4$ Correlation table shows which \mathbf{P}^μ splittings are allowed:

$O \supset C_4$	0_4	1_4	2_4	3_4
$A_1 \downarrow C_4$	1	.	.	.
$A_2 \downarrow C_4$.	.	1	.
$E \downarrow C_4$	1	.	1	.
$T_1 \downarrow C_4$	1	1	.	1
$T_2 \downarrow C_4$.	1	1	1

$\mathbf{P}^{A_1} = \mathbf{P}_{0_4 0_4}^{A_1}$
and
 $\mathbf{P}^{A_2} = \mathbf{P}_{2_4 2_4}^{A_2}$
cannot
split

$$\mathbf{1} \cdot \mathbf{P}^\mu = (\mathbf{p}_{0_4} + \mathbf{p}_{1_4} + \mathbf{p}_{2_4} + \mathbf{p}_{3_4}) \cdot \mathbf{P}^\mu$$

$$\mathbf{1} \cdot \mathbf{P}^{A_1} = \mathbf{P}_{0_4 0_4}^{A_1} + 0 + 0 + 0$$

$$\mathbf{1} \cdot \mathbf{P}^{A_2} = 0 + 0 + \mathbf{P}_{2_4 2_4}^{A_2} + 0$$

$$\mathbf{1} \cdot \mathbf{P}^E = \mathbf{P}_{0_4 0_4}^E + 0 + \mathbf{P}_{2_4 2_4}^E + 0$$

$$\mathbf{1} \cdot \mathbf{P}^{T_1} = \mathbf{P}_{0_4 0_4}^{T_1} + \mathbf{P}_{1_4 1_4}^{T_1} + 0 + \mathbf{P}_{3_4 3_4}^{T_1}$$

$$\mathbf{1} \cdot \mathbf{P}^{T_2} = 0 + \mathbf{P}_{1_4 1_4}^{T_2} + \mathbf{P}_{2_4 2_4}^{T_2} + \mathbf{P}_{3_4 3_4}^{T_2}$$

$O: \chi_g^\mu$	O characters				
	$g=1$	\mathbf{r}_{1-4} $\tilde{\mathbf{r}}_{1-4}$	ρ_{xyz} $\tilde{\mathbf{R}}_{xyz}$	\mathbf{R}_{xyz} $\tilde{\mathbf{R}}_{xyz}$	\mathbf{i}_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

$C_4: d_{R^p}^{m_4}$	C_4 characters			
	$g=1$	\mathbf{R}_z^1	$\rho_z = \mathbf{R}_z^2$	$\tilde{\mathbf{R}}_z = \mathbf{R}_z^3$
$m_4=0_4$	1	1	1	1
1_4	1	-i	-1	i
2_4	1	-1	1	-1
3_4	1	-i	-1	-i

O operators (Two notations: Older Princ. of Symm. Dynamics and Spectra. and Newer Int. J. Mol. Sci)

PSDS:	1	\mathbf{r}_1	\mathbf{r}_2	\mathbf{r}_3	\mathbf{r}_4	\mathbf{r}_1^2	\mathbf{r}_2^2	\mathbf{r}_3^2	\mathbf{r}_4^2	\mathbf{R}_1^2	\mathbf{R}_2^2	\mathbf{R}_3^2	\mathbf{R}_1	\mathbf{R}_2	\mathbf{R}_3	\mathbf{R}_1^3	\mathbf{R}_2^3	\mathbf{R}_3^3	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6
IJMS:	1	\mathbf{r}_1	\mathbf{r}_2	\mathbf{r}_3	\mathbf{r}_4	$\tilde{\mathbf{r}}_1$	$\tilde{\mathbf{r}}_2$	$\tilde{\mathbf{r}}_3$	$\tilde{\mathbf{r}}_4$	ρ_x	ρ_y	ρ_z	\mathbf{R}_x	\mathbf{R}_y	\mathbf{R}_z	$\tilde{\mathbf{R}}_x$	$\tilde{\mathbf{R}}_y$	$\tilde{\mathbf{R}}_z$	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6

Splitting O class projectors \mathbf{P}^μ into irreducible projectors $\mathbf{P}^\mu_{m_4 m_4}$ for $O \supset C_4$

$O \supset C_4$ Correlation table shows which \mathbf{P}^μ splittings are allowed:

$O \supset C_4$	0_4	1_4	2_4	3_4
$A_1 \downarrow C_4$	1	.	.	.
$A_2 \downarrow C_4$.	.	1	.
$E \downarrow C_4$	1	.	1	.
$T_1 \downarrow C_4$	1	1	.	1
$T_2 \downarrow C_4$.	1	1	1

$\mathbf{P}^{A_1} = \mathbf{P}^{A_1}_{0_4 0_4}$
and
 $\mathbf{P}^{A_2} = \mathbf{P}^{A_2}_{2_4 2_4}$
cannot split

$$\mathbf{1} \cdot \mathbf{P}^\mu = (\mathbf{p}_{0_4} + \mathbf{p}_{1_4} + \mathbf{p}_{2_4} + \mathbf{p}_{3_4}) \cdot \mathbf{P}^\mu$$

$$\mathbf{1} \cdot \mathbf{P}^{A_1} = \mathbf{P}^{A_1}_{0_4 0_4} + 0 + 0 + 0$$

$$\mathbf{1} \cdot \mathbf{P}^{A_2} = 0 + 0 + \mathbf{P}^{A_2}_{2_4 2_4} + 0$$

$$\mathbf{1} \cdot \mathbf{P}^E = \mathbf{P}^E_{0_4 0_4} + 0 + \mathbf{P}^E_{2_4 2_4} + 0$$

$$\mathbf{1} \cdot \mathbf{P}^{T_1} = \mathbf{P}^{T_1}_{0_4 0_4} + \mathbf{P}^{T_1}_{1_4 1_4} + 0 + \mathbf{P}^{T_1}_{3_4 3_4}$$

$$\mathbf{1} \cdot \mathbf{P}^{T_2} = 0 + \mathbf{P}^{T_2}_{1_4 1_4} + \mathbf{P}^{T_2}_{2_4 2_4} + \mathbf{P}^{T_2}_{3_4 3_4}$$

$O: \chi_g^\mu$	O characters				
	$g=1$	\mathbf{r}_{1-4} $\tilde{\mathbf{r}}_{1-4}$	ρ_{xyz} $\tilde{\mathbf{R}}_{xyz}$	\mathbf{R}_{xyz} $\tilde{\mathbf{R}}_{xyz}$	\mathbf{i}_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

$C_4: d_{R^p}^{m_4}$	C_4 characters			
	$g=1$	\mathbf{R}_z^1	$\rho_z = \mathbf{R}_z^2$	$\tilde{\mathbf{R}}_z = \mathbf{R}_z^3$
$m_4=0_4$	1	1	1	1
1_4	1	-i	-1	i
2_4	1	-1	1	-1
3_4	1	-i	-1	-i

$O \supset C_4$ splitting done by C_4 projectors applied to O class projectors

$$\mathbf{P}^E = \frac{2}{8} \mathbf{1} - \frac{1}{8} \mathbf{c}_r + \frac{2}{8} \mathbf{c}_\rho + \frac{0}{8} \mathbf{c}_R - \frac{0}{8} \mathbf{c}_i$$

$$\mathbf{P}^{T_1} = \frac{3}{8} \mathbf{1} + \frac{0}{8} \mathbf{c}_r - \frac{1}{8} \mathbf{c}_\rho + \frac{1}{8} \mathbf{c}_R - \frac{1}{8} \mathbf{c}_i$$

$$\mathbf{P}^{T_2} = \frac{3}{8} \mathbf{1} + \frac{0}{8} \mathbf{c}_r - \frac{1}{8} \mathbf{c}_\rho - \frac{1}{8} \mathbf{c}_R + \frac{1}{8} \mathbf{c}_i$$

O operators (Two notations: Older Princ. of Symm. Dynamics and Spectra. and Newer Int. J. Mol. Sci)

PSDS:	1	\mathbf{r}_1	\mathbf{r}_2	\mathbf{r}_3	\mathbf{r}_4	\mathbf{r}_1^2	\mathbf{r}_2^2	\mathbf{r}_3^2	\mathbf{r}_4^2	\mathbf{R}_1^2	\mathbf{R}_2^2	\mathbf{R}_3^2	\mathbf{R}_1	\mathbf{R}_2	\mathbf{R}_3	\mathbf{R}_1^3	\mathbf{R}_2^3	\mathbf{R}_3^3	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6
IJMS:	1	\mathbf{r}_1	\mathbf{r}_2	\mathbf{r}_3	\mathbf{r}_4	$\tilde{\mathbf{r}}_1$	$\tilde{\mathbf{r}}_2$	$\tilde{\mathbf{r}}_3$	$\tilde{\mathbf{r}}_4$	ρ_x	ρ_y	ρ_z	\mathbf{R}_x	\mathbf{R}_y	\mathbf{R}_z	$\tilde{\mathbf{R}}_x$	$\tilde{\mathbf{R}}_y$	$\tilde{\mathbf{R}}_z$	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6

Splitting O class projectors \mathbf{P}^μ into irreducible projectors $\mathbf{P}^\mu_{m_4 m_4}$ for $O \supset C_4$

$O \supset C_4$ Correlation table shows which \mathbf{P}^μ splittings are allowed:

$O \supset C_4$	0_4	1_4	2_4	3_4
$A_1 \downarrow C_4$	1	.	.	.
$A_2 \downarrow C_4$.	.	1	.
$E \downarrow C_4$	1	.	1	.
$T_1 \downarrow C_4$	1	1	.	1
$T_2 \downarrow C_4$.	1	1	1

$\mathbf{P}^{A_1} = \mathbf{P}^{A_1}_{0_4 0_4}$
and
 $\mathbf{P}^{A_2} = \mathbf{P}^{A_2}_{2_4 2_4}$
cannot split

$$\mathbf{1} \cdot \mathbf{P}^\mu = (\mathbf{p}_{0_4} + \mathbf{p}_{1_4} + \mathbf{p}_{2_4} + \mathbf{p}_{3_4}) \cdot \mathbf{P}^\mu$$

$$\mathbf{1} \cdot \mathbf{P}^{A_1} = \mathbf{P}^{A_1}_{0_4 0_4} + 0 + 0 + 0$$

$$\mathbf{1} \cdot \mathbf{P}^{A_2} = 0 + 0 + \mathbf{P}^{A_2}_{2_4 2_4} + 0$$

$$\mathbf{1} \cdot \mathbf{P}^E = \mathbf{P}^E_{0_4 0_4} + 0 + \mathbf{P}^E_{2_4 2_4} + 0$$

$$\mathbf{1} \cdot \mathbf{P}^{T_1} = \mathbf{P}^{T_1}_{0_4 0_4} + \mathbf{P}^{T_1}_{1_4 1_4} + 0 + \mathbf{P}^{T_1}_{3_4 3_4}$$

$$\mathbf{1} \cdot \mathbf{P}^{T_2} = 0 + \mathbf{P}^{T_2}_{1_4 1_4} + \mathbf{P}^{T_2}_{2_4 2_4} + \mathbf{P}^{T_2}_{3_4 3_4}$$

$O: \chi_g^\mu$	O characters				
	$g=1$	\mathbf{r}_{1-4} $\tilde{\mathbf{r}}_{1-4}$	ρ_{xyz} $\tilde{\mathbf{R}}_{xyz}$	\mathbf{R}_{xyz} $\tilde{\mathbf{R}}_{xyz}$	\mathbf{i}_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

$C_4: d_{R^p}^{m_4}$	C_4 characters			
	$g=1$	\mathbf{R}_z^1	$\rho_z = \mathbf{R}_z^2$	$\tilde{\mathbf{R}}_z = \mathbf{R}_z^3$
$m_4=0_4$	1	1	1	1
1_4	1	-i	-1	i
2_4	1	-1	1	-1
3_4	1	-i	-1	-i

$O \supset C_4$ splitting done by C_4 projectors applied to O class projectors

$$\mathbf{P}^E = \frac{2}{8} \mathbf{1} - \frac{1}{8} \mathbf{c}_r + \frac{2}{8} \mathbf{c}_\rho + \frac{0}{8} \mathbf{c}_R - \frac{0}{8} \mathbf{c}_i$$

$$\mathbf{P}^{T_1} = \frac{3}{8} \mathbf{1} + \frac{0}{8} \mathbf{c}_r - \frac{1}{8} \mathbf{c}_\rho + \frac{1}{8} \mathbf{c}_R - \frac{1}{8} \mathbf{c}_i$$

$$\mathbf{P}^{T_2} = \frac{3}{8} \mathbf{1} + \frac{0}{8} \mathbf{c}_r - \frac{1}{8} \mathbf{c}_\rho - \frac{1}{8} \mathbf{c}_R + \frac{1}{8} \mathbf{c}_i$$

$$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 e^{\frac{2\pi i m_4 \cdot p}{4}} \mathbf{R}_z^p = \underbrace{\left\{ \begin{array}{l} \mathbf{p}_{0_4} = (1 + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{1_4} = (1 + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{2_4} = (1 - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{3_4} = (1 - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4 \end{array} \right.}_{C_4 \text{ characters}}$$

$$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$$

O operators (Two notations: Older Princ. of Symm. Dynamics and Spectra. and Newer Int. J. Mol. Sci)

PSDS:	1	\mathbf{r}_1	\mathbf{r}_2	\mathbf{r}_3	\mathbf{r}_4	\mathbf{r}_1^2	\mathbf{r}_2^2	\mathbf{r}_3^2	\mathbf{r}_4^2	\mathbf{R}_1^2	\mathbf{R}_2^2	\mathbf{R}_3^2	\mathbf{R}_1	\mathbf{R}_2	\mathbf{R}_3	\mathbf{R}_1^3	\mathbf{R}_2^3	\mathbf{R}_3^3	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6
IJMS:	1	\mathbf{r}_1	\mathbf{r}_2	\mathbf{r}_3	\mathbf{r}_4	$\tilde{\mathbf{r}}_1$	$\tilde{\mathbf{r}}_2$	$\tilde{\mathbf{r}}_3$	$\tilde{\mathbf{r}}_4$	ρ_x	ρ_y	ρ_z	\mathbf{R}_x	\mathbf{R}_y	\mathbf{R}_z	$\tilde{\mathbf{R}}_x$	$\tilde{\mathbf{R}}_y$	$\tilde{\mathbf{R}}_z$	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6

Splitting O class projectors \mathbf{P}^μ into irreducible projectors $\mathbf{P}^\mu_{m_4 m_4}$ for $O \supset C_4$

$O \supset C_4$ Correlation table shows which \mathbf{P}^μ splittings are allowed:

$O \supset C_4$	0_4	1_4	2_4	3_4
$A_1 \downarrow C_4$	1	.	.	.
$A_2 \downarrow C_4$.	.	1	.
$E \downarrow C_4$	1	.	1	.
$T_1 \downarrow C_4$	1	1	.	1
$T_2 \downarrow C_4$.	1	1	1

$\mathbf{P}^{A_1} = \mathbf{P}^{A_1}_{0_4 0_4}$
and
 $\mathbf{P}^{A_2} = \mathbf{P}^{A_2}_{2_4 2_4}$
cannot split

$$\mathbf{1} \cdot \mathbf{P}^\mu = (\mathbf{p}_{0_4} + \mathbf{p}_{1_4} + \mathbf{p}_{2_4} + \mathbf{p}_{3_4}) \cdot \mathbf{P}^\mu$$

$$\mathbf{1} \cdot \mathbf{P}^{A_1} = \mathbf{P}^{A_1}_{0_4 0_4} + 0 + 0 + 0$$

$$\mathbf{1} \cdot \mathbf{P}^{A_2} = 0 + 0 + \mathbf{P}^{A_2}_{2_4 2_4} + 0$$

$$\mathbf{1} \cdot \mathbf{P}^E = \mathbf{P}^E_{0_4 0_4} + 0 + \mathbf{P}^E_{2_4 2_4} + 0$$

$$\mathbf{1} \cdot \mathbf{P}^{T_1} = \mathbf{P}^{T_1}_{0_4 0_4} + \mathbf{P}^{T_1}_{1_4 1_4} + 0 + \mathbf{P}^{T_1}_{3_4 3_4}$$

$$\mathbf{1} \cdot \mathbf{P}^{T_2} = 0 + \mathbf{P}^{T_2}_{1_4 1_4} + \mathbf{P}^{T_2}_{2_4 2_4} + \mathbf{P}^{T_2}_{3_4 3_4}$$

$O: \chi_g^\mu$	<i>O characters</i>				
	$g=1$	\mathbf{r}_{1-4} $\tilde{\mathbf{r}}_{1-4}$	ρ_{xyz}	\mathbf{R}_{xyz} $\tilde{\mathbf{R}}_{xyz}$	\mathbf{i}_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

$C_4: d_{R^p}^{m_4}$	<i>C₄ characters</i>			
	$g=1$	\mathbf{R}_z^1	$\rho_z = \mathbf{R}_z^2$	$\tilde{\mathbf{R}}_z = \mathbf{R}_z^3$
$m_4=0_4$	1	1	1	1
1_4	1	-i	-1	i
2_4	1	-1	1	-1
3_4	1	-i	-1	-i

$O \supset C_4$ splitting done by C_4 projectors applied to O class projectors

$$\mathbf{P}^E = \frac{2}{8} \mathbf{1} - \frac{1}{8} \mathbf{c}_r + \frac{2}{8} \mathbf{c}_\rho + \frac{0}{8} \mathbf{c}_R - \frac{0}{8} \mathbf{c}_i$$

$$\mathbf{P}^{T_1} = \frac{3}{8} \mathbf{1} + \frac{0}{8} \mathbf{c}_r - \frac{1}{8} \mathbf{c}_\rho + \frac{1}{8} \mathbf{c}_R - \frac{1}{8} \mathbf{c}_i$$

$$\mathbf{P}^{T_2} = \frac{3}{8} \mathbf{1} + \frac{0}{8} \mathbf{c}_r - \frac{1}{8} \mathbf{c}_\rho - \frac{1}{8} \mathbf{c}_R + \frac{1}{8} \mathbf{c}_i$$

$$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 e^{\frac{2\pi i m_4 \cdot p}{4}} \mathbf{R}_z^p = \underbrace{\left\{ \begin{array}{l} \mathbf{p}_{0_4} = (1 + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{1_4} = (1 + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{2_4} = (1 - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{3_4} = (1 - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4 \end{array} \right.}_{C_4 \text{ characters}}$$

$$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$$

Following development of irreducible projectors:

$$\mathbf{P}^\mu_{m_4 m_4} \equiv \mathbf{p}_{m_4} \mathbf{P}^\mu = \mathbf{P}^\mu \mathbf{p}_{m_4}$$

O operators (Two notations: Older Princ. of Symm. Dynamics and Spectra. and Newer Int. J. Mol. Sci)

PSDS:	1	\mathbf{r}_1	\mathbf{r}_2	\mathbf{r}_3	\mathbf{r}_4	\mathbf{r}_1^2	\mathbf{r}_2^2	\mathbf{r}_3^2	\mathbf{r}_4^2	\mathbf{R}_1^2	\mathbf{R}_2^2	\mathbf{R}_3^2	\mathbf{R}_1	\mathbf{R}_2	\mathbf{R}_3	\mathbf{R}_1^3	\mathbf{R}_2^3	\mathbf{R}_3^3	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6
IJMS:	1	\mathbf{r}_1	\mathbf{r}_2	\mathbf{r}_3	\mathbf{r}_4	$\tilde{\mathbf{r}}_1$	$\tilde{\mathbf{r}}_2$	$\tilde{\mathbf{r}}_3$	$\tilde{\mathbf{r}}_4$	ρ_x	ρ_y	ρ_z	\mathbf{R}_x	\mathbf{R}_y	\mathbf{R}_z	$\tilde{\mathbf{R}}_x$	$\tilde{\mathbf{R}}_y$	$\tilde{\mathbf{R}}_z$	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6

Splitting O class projectors \mathbf{P}^μ into irreducible projectors $\mathbf{P}^\mu_{m_4 m_4}$ for $O \supset C_4$

$O \supset C_4$ Correlation table shows which \mathbf{P}^μ splittings are allowed:

$O \supset C_4$	0_4	1_4	2_4	3_4
$A_1 \downarrow C_4$	1	.	.	.
$A_2 \downarrow C_4$.	.	1	.
$E \downarrow C_4$	1	.	1	.
$T_1 \downarrow C_4$	1	1	.	1
$T_2 \downarrow C_4$.	1	1	1

$\mathbf{P}^{A_1} = \mathbf{P}^{A_1}_{0_4 0_4}$
and
 $\mathbf{P}^{A_2} = \mathbf{P}^{A_2}_{2_4 2_4}$
cannot split

$$\mathbf{1} \cdot \mathbf{P}^\mu = (\mathbf{p}_{0_4} + \mathbf{p}_{1_4} + \mathbf{p}_{2_4} + \mathbf{p}_{3_4}) \cdot \mathbf{P}^\mu$$

$$\mathbf{1} \cdot \mathbf{P}^{A_1} = \mathbf{P}^{A_1}_{0_4 0_4} + 0 + 0 + 0$$

$$\mathbf{1} \cdot \mathbf{P}^{A_2} = 0 + 0 + \mathbf{P}^{A_2}_{2_4 2_4} + 0$$

$$\mathbf{1} \cdot \mathbf{P}^E = \mathbf{P}^E_{0_4 0_4} + 0 + \mathbf{P}^E_{2_4 2_4} + 0$$

$$\mathbf{1} \cdot \mathbf{P}^{T_1} = \mathbf{P}^{T_1}_{0_4 0_4} + \mathbf{P}^{T_1}_{1_4 1_4} + 0 + \mathbf{P}^{T_1}_{3_4 3_4}$$

$$\mathbf{1} \cdot \mathbf{P}^{T_2} = 0 + \mathbf{P}^{T_2}_{1_4 1_4} + \mathbf{P}^{T_2}_{2_4 2_4} + \mathbf{P}^{T_2}_{3_4 3_4}$$

$O: \chi_g^\mu$	O characters				
	$g=1$	\mathbf{r}_{1-4} $\tilde{\mathbf{r}}_{1-4}$	ρ_{xyz} $\tilde{\mathbf{R}}_{xyz}$	\mathbf{R}_{xyz} $\tilde{\mathbf{R}}_{xyz}$	\mathbf{i}_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

$C_4: d_{R^p}^{m_4}$	C_4 characters			
	$g=1$	\mathbf{R}_z^1	$\rho_z = \mathbf{R}_z^2$	$\tilde{\mathbf{R}}_z = \mathbf{R}_z^3$
$m_4=0_4$	1	1	1	1
1_4	1	-i	-1	i
2_4	1	-1	1	-1
3_4	1	-i	-1	-i

$O \supset C_4$ splitting done by C_4 projectors applied to O class projectors

$$\mathbf{P}^E = \frac{2}{8} \mathbf{1} - \frac{1}{8} \mathbf{c}_r + \frac{2}{8} \mathbf{c}_\rho + \frac{0}{8} \mathbf{c}_R - \frac{0}{8} \mathbf{c}_i$$

$$\mathbf{P}^{T_1} = \frac{3}{8} \mathbf{1} + \frac{0}{8} \mathbf{c}_r - \frac{1}{8} \mathbf{c}_\rho + \frac{1}{8} \mathbf{c}_R - \frac{1}{8} \mathbf{c}_i$$

$$\mathbf{P}^{T_2} = \frac{3}{8} \mathbf{1} + \frac{0}{8} \mathbf{c}_r - \frac{1}{8} \mathbf{c}_\rho - \frac{1}{8} \mathbf{c}_R + \frac{1}{8} \mathbf{c}_i$$

$$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 \underbrace{e^{\frac{2\pi i m_4 \cdot p}{4}}}_{C_4 \text{ characters}} \mathbf{R}_z^p = \left\{ \begin{array}{l} \mathbf{p}_{0_4} = (\mathbf{1} + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z) / 4 \\ \mathbf{p}_{1_4} = (\mathbf{1} + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z) / 4 \\ \mathbf{p}_{2_4} = (\mathbf{1} - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z) / 4 \\ \mathbf{p}_{3_4} = (\mathbf{1} - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z) / 4 \end{array} \right.$$

$$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$$

Following development of irreducible projectors:

$$\mathbf{P}^\mu_{m_4 m_4} \equiv \mathbf{p}_{m_4} \mathbf{P}^\mu = \mathbf{P}^\mu \mathbf{p}_{m_4}$$

...with examples:

$$\mathbf{P}^{T_1}_{0_4 0_4} \equiv \mathbf{p}_{0_4} \mathbf{P}^{T_1} = \mathbf{P}^{T_1} \mathbf{p}_{0_4}$$

$$\mathbf{P}^{T_1}_{1_4 1_4} \equiv \mathbf{p}_{1_4} \mathbf{P}^{T_1} = \mathbf{P}^{T_1} \mathbf{p}_{1_4}$$

etc.

O operators (Two notations: Older Princ. of Symm. Dynamics and Spectra. and Newer Int. J. Mol. Sci)

PSDS:	1	\mathbf{r}_1	\mathbf{r}_2	\mathbf{r}_3	\mathbf{r}_4	\mathbf{r}_1^2	\mathbf{r}_2^2	\mathbf{r}_3^2	\mathbf{r}_4^2	\mathbf{R}_1^2	\mathbf{R}_2^2	\mathbf{R}_3^2	\mathbf{R}_1	\mathbf{R}_2	\mathbf{R}_3	\mathbf{R}_1^3	\mathbf{R}_2^3	\mathbf{R}_3^3	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6
IJMS:	1	\mathbf{r}_1	\mathbf{r}_2	\mathbf{r}_3	\mathbf{r}_4	$\tilde{\mathbf{r}}_1$	$\tilde{\mathbf{r}}_2$	$\tilde{\mathbf{r}}_3$	$\tilde{\mathbf{r}}_4$	ρ_x	ρ_y	ρ_z	\mathbf{R}_x	\mathbf{R}_y	\mathbf{R}_z	$\tilde{\mathbf{R}}_x$	$\tilde{\mathbf{R}}_y$	$\tilde{\mathbf{R}}_z$	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6

Splitting O class projectors \mathbf{P}^μ into irreducible projectors $\mathbf{P}^\mu_{m_4m_4}$ for $O \supset C_4$

$O \supset C_4$ Correlation table shows which \mathbf{P}^μ splittings are allowed:

$O \supset C_4$	0_4	1_4	2_4	3_4
$A_1 \downarrow C_4$	1	.	.	.
$A_2 \downarrow C_4$.	.	1	.
$E \downarrow C_4$	1	.	1	.
$T_1 \downarrow C_4$	1	1	.	1
$T_2 \downarrow C_4$.	1	1	1

$\mathbf{P}^{A_1} = \mathbf{P}^{A_1}_{0_4 0_4}$
and
 $\mathbf{P}^{A_2} = \mathbf{P}^{A_2}_{2_4 2_4}$
cannot split

$$\mathbf{1} \cdot \mathbf{P}^\mu = (\mathbf{p}_{0_4} + \mathbf{p}_{1_4} + \mathbf{p}_{2_4} + \mathbf{p}_{3_4}) \cdot \mathbf{P}^\mu$$

$$\mathbf{1} \cdot \mathbf{P}^{A_1} = \mathbf{P}^{A_1}_{0_4 0_4} + 0 + 0 + 0$$

$$\mathbf{1} \cdot \mathbf{P}^{A_2} = 0 + 0 + \mathbf{P}^{A_2}_{2_4 2_4} + 0$$

$$\mathbf{1} \cdot \mathbf{P}^E = \mathbf{P}^E_{0_4 0_4} + 0 + \mathbf{P}^E_{2_4 2_4} + 0$$

$$\mathbf{1} \cdot \mathbf{P}^{T_1} = \mathbf{P}^{T_1}_{0_4 0_4} + \mathbf{P}^{T_1}_{1_4 1_4} + 0 + \mathbf{P}^{T_1}_{3_4 3_4}$$

$$\mathbf{1} \cdot \mathbf{P}^{T_2} = 0 + \mathbf{P}^{T_2}_{1_4 1_4} + \mathbf{P}^{T_2}_{2_4 2_4} + \mathbf{P}^{T_2}_{3_4 3_4}$$

$O: \chi_g^\mu$	<i>O characters</i>				
	$g=1$	\mathbf{r}_{1-4} $\tilde{\mathbf{r}}_{1-4}$	ρ_{xyz} $\tilde{\mathbf{R}}_{xyz}$	\mathbf{R}_{xyz} $\tilde{\mathbf{R}}_{xyz}$	\mathbf{i}_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1
$C_4: d_{R^p}^{m_4}$	<i>C₄ characters</i>				
	$g=1$	\mathbf{R}_z^1	$\rho_z = \mathbf{R}_z^2$	$\tilde{\mathbf{R}}_z = \mathbf{R}_z^3$	
$m_4=0_4$	1	1	1	1	
1_4	1	-i	-1	i	
2_4	1	-1	1	-1	
3_4	1	-i	-1	-i	

$O \supset C_4$ splitting done by C_4 projectors applied to O class projectors

$$\mathbf{P}^E = \frac{2}{8}\mathbf{1} - \frac{1}{8}\mathbf{c}_r + \frac{2}{8}\mathbf{c}_\rho + \frac{0}{8}\mathbf{c}_R - \frac{0}{8}\mathbf{c}_i$$

$$\mathbf{P}^{T_1} = \frac{3}{8}\mathbf{1} + \frac{0}{8}\mathbf{c}_r - \frac{1}{8}\mathbf{c}_\rho + \frac{1}{8}\mathbf{c}_R - \frac{1}{8}\mathbf{c}_i$$

$$\mathbf{P}^{T_2} = \frac{3}{8}\mathbf{1} + \frac{0}{8}\mathbf{c}_r - \frac{1}{8}\mathbf{c}_\rho - \frac{1}{8}\mathbf{c}_R + \frac{1}{8}\mathbf{c}_i$$

$$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 e^{\frac{2\pi i m_4 \cdot p}{4}} \mathbf{R}_z^p = \underbrace{\dots}_{C_4 \text{ characters}}$$

$$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$$

$$\mathbf{p}_{0_4} = (\mathbf{1} + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z) / 4$$

$$\mathbf{p}_{1_4} = (\mathbf{1} + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z) / 4$$

$$\mathbf{p}_{2_4} = (\mathbf{1} - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z) / 4$$

$$\mathbf{p}_{3_4} = (\mathbf{1} - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z) / 4$$

Following development of irreducible projectors:

$$\mathbf{P}^\mu_{m_4 m_4} \equiv \mathbf{p}_{m_4} \mathbf{P}^\mu = \mathbf{P}^\mu \mathbf{p}_{m_4}$$

...with examples:

$$\mathbf{P}^{T_1}_{0_4 0_4} \equiv \mathbf{p}_{0_4} \mathbf{P}^{T_1} = \mathbf{P}^{T_1} \mathbf{p}_{0_4}$$

$$\mathbf{P}^{T_1}_{1_4 1_4} \equiv \mathbf{p}_{1_4} \mathbf{P}^{T_1} = \mathbf{P}^{T_1} \mathbf{p}_{1_4}$$

etc.

...uses left-coset combinations...

...and projector "factoring"...

$$\mathbf{1}C_4 = \mathbf{1}\{\mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z\}, \rho_x C_4 = \{\rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3\}, \mathbf{r}_1 C_4 = \{\mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y\}, \mathbf{r}_2 C_4 = \{\mathbf{r}_2, \mathbf{r}_3, \mathbf{i}_2, \tilde{\mathbf{R}}_y\}, \tilde{\mathbf{r}}_1 C_4 = \{\tilde{\mathbf{r}}_1, \tilde{\mathbf{r}}_3, \tilde{\mathbf{R}}_x, \mathbf{i}_6\}, \tilde{\mathbf{r}}_2 C_4 = \{\tilde{\mathbf{r}}_2, \tilde{\mathbf{r}}_4, \mathbf{R}_x, \mathbf{i}_5\}.$$

O operators (Two notations: Older Princ. of Symm. Dynamics and Spectra. and Newer Int. J. Mol. Sci)

PSDS:	1	\mathbf{r}_1	\mathbf{r}_2	\mathbf{r}_3	\mathbf{r}_4	\mathbf{r}_1^2	\mathbf{r}_2^2	\mathbf{r}_3^2	\mathbf{r}_4^2	\mathbf{R}_1^2	\mathbf{R}_2^2	\mathbf{R}_3^2	\mathbf{R}_1	\mathbf{R}_2	\mathbf{R}_3	\mathbf{R}_1^3	\mathbf{R}_2^3	\mathbf{R}_3^3	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6
IJMS:	1	\mathbf{r}_1	\mathbf{r}_2	\mathbf{r}_3	\mathbf{r}_4	$\tilde{\mathbf{r}}_1$	$\tilde{\mathbf{r}}_2$	$\tilde{\mathbf{r}}_3$	$\tilde{\mathbf{r}}_4$	ρ_x	ρ_y	ρ_z	\mathbf{R}_x	\mathbf{R}_y	\mathbf{R}_z	$\tilde{\mathbf{R}}_x$	$\tilde{\mathbf{R}}_y$	$\tilde{\mathbf{R}}_z$	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6


Following development of irreducible projectors:

$$\mathbf{P}_{m_4 m_4}^\mu \equiv \mathbf{p}_{m_4} \mathbf{P}^\mu = \mathbf{P}^\mu \mathbf{p}_{m_4}$$

...uses left-coset combinations...

...and projector "factoring"...

$$1C_4 = 1\{\mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z\}, \rho_x C_4 = \{\rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3\}, \mathbf{r}_1 C_4 = \{\mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y\}, \mathbf{r}_2 C_4 = \{\mathbf{r}_2, \mathbf{r}_3, \mathbf{i}_2, \tilde{\mathbf{R}}_y\}, \tilde{\mathbf{r}}_1 C_4 = \{\tilde{\mathbf{r}}_1, \tilde{\mathbf{r}}_3, \tilde{\mathbf{R}}_x, \mathbf{i}_6\}, \tilde{\mathbf{r}}_2 C_4 = \{\tilde{\mathbf{r}}_2, \tilde{\mathbf{r}}_4, \mathbf{R}_x, \mathbf{i}_5\}$$



Review Octahedral $O_h \supset O$ group operator structure

Review Octahedral $O_h \supset O \supset D_4 \supset C_4$ subgroup chain correlations

Comparison of $O \supset D_4 \supset C_4$ and $O \supset D_4 \supset D_2$ correlations and level/projector splitting

$O \supset D_4 \supset C_4$ subgroup chain splitting

$O \supset D_4 \supset D_2$ subgroup chain splitting (normal D_2 vs. unnormal D_2)

$O_h \supset O \supset D_4 \supset C_{4v}$ and $O_h \supset O \supset D_4 \supset C_{4v} \supset C_{2v}$ subgroup splitting

Splitting O class projectors \mathbf{P}^μ into irreducible projectors $\mathbf{P}^\mu_{m_4m_4}$ for $O \supset C_4$

Development of irreducible projectors $\mathbf{P}^\mu_{m_4m_4}$ and representations $D^\mu_{m_4m_4}$

Calculating $\mathbf{P}^E_{0_40_4}$, $\mathbf{P}^E_{2_42_4}$, $\mathbf{P}^{T_1}_{0_40_4}$, $\mathbf{P}^{T_1}_{1_41_4}$, $\mathbf{P}^{T_2}_{2_42_4}$, $\mathbf{P}^{T_2}_{1_41_4}$,


$O \supset C_4$ induced representation $0_4(C_4) \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples

Elementary induced representation $0_4(C_4) \uparrow O$

Projection reduction of induced representation $0_4(C_4) \uparrow O$

Introduction to ortho-complete eigenvalue-parameter relations

Examples in SF_6 spectroscopy



General development of $O \supset C_4$ irreducible projectors


$$\mathbf{P}_{m_4 m_4}^\mu = \sum_g \frac{\ell^\mu}{\circ O} D_{m_4 m_4}^{\mu*}(g) \mathbf{g}$$

$$\mathbf{P}_{m_4 m_4}^\mu = \mathbf{p}^{m_4} \mathbf{P}^\mu = \mathbf{P}^\mu \mathbf{p}^{m_4}$$

Deriving diagonal irreducible O -representation components $D_{m_4 m_4}^{\mu*}(g)$

$$= \sum_g \frac{\ell^\mu}{\circ O} (\chi_g^{\mu*}) \cdot \mathbf{g}(\mathbf{p}^{m_4})$$

$$= \left(\frac{\ell^\mu}{24} \right) (\chi_1^{\mu*}) \cdot \mathbf{1} \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4}$$



$$\rho_x C_4 = \rho_x \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3 \} \text{Coset}$$

$O: \chi_g^\mu$	O characters				
	$\mathbf{g}=\mathbf{1}$	\mathbf{r}_{1-4} $\tilde{\mathbf{r}}_{1-4}$	ρ_{xyz}	\mathbf{R}_{xyz} $\tilde{\mathbf{R}}_{xyz}$	\mathbf{i}_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

$$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 \underbrace{e^{\frac{2\pi i m_4 \cdot p}{4}}}_{C_4 \text{ characters}} \mathbf{R}_z^p =$$

$$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$$

}

$$\begin{cases} \mathbf{p}_{0_4} = (\mathbf{1} + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{1_4} = (\mathbf{1} + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{2_4} = (\mathbf{1} - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{3_4} = (\mathbf{1} - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4 \end{cases}$$

$$\mathbf{r}_1 C_4 = \mathbf{r}_1 \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y \} \text{Coset}$$

General development of irep projectors $\mathbf{P}_{m_4 m_4}^\mu = \sum_g \frac{\ell^\mu}{\circ O} D_{m_4 m_4}^{\mu*}(g) \mathbf{g}$ for subgroup chain $O \supset D_4 \supset C_4$

$$\mathbf{P}_{m_4 m_4}^\mu = \mathbf{p}^{m_4} \mathbf{P}^\mu = \mathbf{P}^\mu \mathbf{p}^{m_4}$$

$$= \sum_g \frac{\ell^\mu}{\circ O} (\chi_g^{\mu*}) \cdot \mathbf{g}(\mathbf{p}^{m_4})$$

$$= \left(\frac{\ell^\mu}{24} \right) (\chi_1^{\mu*}) \cdot \mathbf{1} \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

(Deriving diagonal irreducible O -representation (“irep”) components $D_{m_4 m_4}^{\mu*}(g)$)

+
+
+

$$\rho_x C_4 = \rho_x \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3 \} \text{Coset}$$

+
+
+
+

$O: \chi_g^\mu$	O characters				
	$\mathbf{g}=\mathbf{1}$	\mathbf{r}_{1-4} $\tilde{\mathbf{r}}_{1-4}$	ρ_{xyz}	\mathbf{R}_{xyz} $\tilde{\mathbf{R}}_{xyz}$	\mathbf{i}_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

$$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 \underbrace{e^{\frac{2\pi i m_4 \cdot p}{4}}}_{C_4 \text{ characters}} \mathbf{R}_z^p = \begin{cases} \mathbf{p}_{0_4} = (\mathbf{1} + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{1_4} = (\mathbf{1} + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{2_4} = (\mathbf{1} - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{3_4} = (\mathbf{1} - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4 \end{cases}$$

$$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$$

$$\mathbf{r}_1 C_4 = \mathbf{r}_1 \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y \} \text{Coset}$$

General development of irep projectors $\mathbf{P}_{m_4 m_4}^\mu = \sum_g \frac{\ell^\mu}{\circ O} D_{m_4 m_4}^{\mu*}(g) \mathbf{g}$ for subgroup chain $O \supset D_4 \supset C_4$

$$\mathbf{P}_{m_4 m_4}^\mu = \mathbf{p}^{m_4} \mathbf{P}^\mu = \mathbf{P}^\mu \mathbf{p}^{m_4}$$

$$= \sum_g \frac{\ell^\mu}{\circ O} (\chi_g^{\mu*}) \cdot \mathbf{g}(\mathbf{p}^{m_4})$$

(Deriving diagonal irreducible O -representation (“irep”) components $D_{m_4 m_4}^{\mu*}(g)$)

$$= \left(\frac{\ell^\mu}{24}\right) (\chi_1^{\mu*}) \cdot \mathbf{1} \left(\mathbf{1} + d_{\rho_z}^{m_4} \boldsymbol{\rho}_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96}\right) \left(\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \boldsymbol{\rho}_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24}\right) (\chi_{\rho_z}^{\mu*}) \cdot \boldsymbol{\rho}_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \boldsymbol{\rho}_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4}$$

+

+

$$\boldsymbol{\rho}_x C_4 = \boldsymbol{\rho}_x \{ \mathbf{1}, \boldsymbol{\rho}_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \boldsymbol{\rho}_x, \boldsymbol{\rho}_y, \mathbf{i}_4, \mathbf{i}_3 \} \text{Coset}$$

+

+

+

+

$O: \chi_g^\mu$	O characters				
	$\mathbf{g}=\mathbf{1}$	\mathbf{r}_{1-4} $\tilde{\mathbf{r}}_{1-4}$	$\boldsymbol{\rho}_{xyz}$	\mathbf{R}_{xyz} $\tilde{\mathbf{R}}_{xyz}$	\mathbf{i}_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

$$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 \underbrace{e^{\frac{2\pi i m_4 \cdot p}{4}}}_{C_4 \text{ characters}} \mathbf{R}_z^p = \left\{ \begin{array}{l} \mathbf{p}_{0_4} = (\mathbf{1} + \mathbf{R}_z + \boldsymbol{\rho}_z + \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{1_4} = (\mathbf{1} + i\mathbf{R}_z - \boldsymbol{\rho}_z - i\tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{2_4} = (\mathbf{1} - \mathbf{R}_z + \boldsymbol{\rho}_z - \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{3_4} = (\mathbf{1} - i\mathbf{R}_z - \boldsymbol{\rho}_z + i\tilde{\mathbf{R}}_z)/4 \end{array} \right.$$

$$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$$

$$\mathbf{r}_1 C_4 = \mathbf{r}_1 \{ \mathbf{1}, \boldsymbol{\rho}_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y \} \text{Coset}$$

General development of irep projectors $\mathbf{P}_{m_4 m_4}^\mu = \sum_g \frac{\ell^\mu}{\circ O} D_{m_4 m_4}^{\mu*}(g) \mathbf{g}$ for subgroup chain $O \supset D_4 \supset C_4$

$$\mathbf{P}_{m_4 m_4}^\mu = \mathbf{p}^{m_4} \mathbf{P}^\mu = \mathbf{P}^\mu \mathbf{p}^{m_4}$$

$$= \sum_g \frac{\ell^\mu}{\circ O} (\chi_g^{\mu*}) \cdot \mathbf{g}(\mathbf{p}^{m_4})$$

(Deriving diagonal irreducible O -representation (“irep”) components $D_{m_4 m_4}^{\mu*}(g)$)

$$= \left(\frac{\ell^\mu}{24} \right) (\chi_1^{\mu*}) \cdot \mathbf{1} \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_z}^{\mu*}) \cdot \rho_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_z + d_{\rho_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z \right)$$

+
+

$$\rho_x C_4 = \rho_x \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3 \} \text{Coset}$$

$O: \chi_g^\mu$	$\mathbf{g}=\mathbf{1}$	O characters			
		\mathbf{r}_{1-4} $\tilde{\mathbf{r}}_{1-4}$	ρ_{xyz}	\mathbf{R}_{xyz} $\tilde{\mathbf{R}}_{xyz}$	\mathbf{i}_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 \underbrace{e^{\frac{2\pi i m_4 \cdot p}{4}}}_{C_4 \text{ characters}} \mathbf{R}_z^p = \begin{cases} \mathbf{p}_{0_4} = (\mathbf{1} + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{1_4} = (\mathbf{1} + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{2_4} = (\mathbf{1} - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{3_4} = (\mathbf{1} - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4 \end{cases}$

$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$

$$\mathbf{r}_1 C_4 = \mathbf{r}_1 \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y \} \text{Coset}$$

General development of irep projectors $\mathbf{P}_{m_4 m_4}^\mu = \sum_g \frac{\ell^\mu}{\circ O} D_{m_4 m_4}^{\mu*}(g) \mathbf{g}$ for subgroup chain $O \supset D_4 \supset C_4$

$$\mathbf{P}_{m_4 m_4}^\mu = \mathbf{p}^{m_4} \mathbf{P}^\mu = \mathbf{P}^\mu \mathbf{p}^{m_4}$$

$$= \sum_g \frac{\ell^\mu}{\circ O} (\chi_g^{\mu*}) \cdot \mathbf{g}(\mathbf{p}^{m_4})$$

(Deriving diagonal irreducible O -representation (“irep”) components $D_{m_4 m_4}^{\mu*}(g)$)

$$= \left(\frac{\ell^\mu}{24} \right) (\chi_1^{\mu*}) \cdot \mathbf{1} \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_z}^{\mu*}) \cdot \rho_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_z + d_{\rho_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z \right) = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) \left(d_{\rho_z}^{m_4} \mathbf{1} + \mathbf{1} \cdot \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

+
+

$$\rho_x C_4 = \rho_x \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3 \} \text{Coset}$$

$O: \chi_g^\mu$	$\mathbf{g}=\mathbf{1}$	O characters			
		\mathbf{r}_{1-4} $\tilde{\mathbf{r}}_{1-4}$	ρ_{xyz}	\mathbf{R}_{xyz} $\tilde{\mathbf{R}}_{xyz}$	\mathbf{i}_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

$$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 \underbrace{e^{\frac{2\pi i m_4 \cdot p}{4}}}_{C_4 \text{ characters}} \mathbf{R}_z^p =$$

$$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$$

}

$$\begin{cases} \mathbf{p}_{0_4} = (\mathbf{1} + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{1_4} = (\mathbf{1} + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{2_4} = (\mathbf{1} - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{3_4} = (\mathbf{1} - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4 \end{cases}$$

$$\mathbf{r}_1 C_4 = \mathbf{r}_1 \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y \} \text{Coset}$$

General development of irep projectors $\mathbf{P}_{m_4 m_4}^\mu = \sum_g \frac{\ell^\mu}{\circ O} D_{m_4 m_4}^{\mu*}(g) \mathbf{g}$ for subgroup chain $O \supset D_4 \supset C_4$

$$\mathbf{P}_{m_4 m_4}^\mu = \mathbf{p}^{m_4} \mathbf{P}^\mu = \mathbf{P}^\mu \mathbf{p}^{m_4}$$

(Deriving diagonal irreducible O -representation (“irep”) components $D_{m_4 m_4}^{\mu*}(g)$)

$$= \sum_g \frac{\ell^\mu}{\circ O} (\chi_g^{\mu*}) \cdot \mathbf{g}(\mathbf{p}^{m_4})$$

$$= \left(\frac{\ell^\mu}{24} \right) (\chi_1^{\mu*}) \cdot \mathbf{1} (\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z) \frac{1}{4} = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) (\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z) = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) (\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_z}^{\mu*}) \cdot \rho_z (\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) (\mathbf{1} \cdot \rho_z + d_{\rho_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z) = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) (d_{\rho_z}^{m_4} \mathbf{1} + \mathbf{1} \cdot \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{R_z}^{\mu*}) \cdot \mathbf{R}_z (\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z) \frac{1}{4}$$

+

$$\rho_x C_4 = \rho_x \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3 \} \text{Coset}$$

+

+

+

+

$O: \chi_g^\mu$	$\mathbf{g}=\mathbf{1}$	O characters			
		\mathbf{r}_{1-4} $\tilde{\mathbf{r}}_{1-4}$	ρ_{xyz}	\mathbf{R}_{xyz} $\tilde{\mathbf{R}}_{xyz}$	\mathbf{i}_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

$$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 \underbrace{e^{\frac{2\pi i m_4 \cdot p}{4}}}_{C_4 \text{ characters}} \mathbf{R}_z^p = \begin{cases} \mathbf{p}_{0_4} = (\mathbf{1} + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{1_4} = (\mathbf{1} + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{2_4} = (\mathbf{1} - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{3_4} = (\mathbf{1} - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4 \end{cases}$$

$$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$$

$$\mathbf{r}_1 C_4 = \mathbf{r}_1 \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y \} \text{Coset}$$

General development of irep projectors $\mathbf{P}_{m_4 m_4}^\mu = \sum_g \frac{\ell^\mu}{\circ O} D_{m_4 m_4}^{\mu*}(g) \mathbf{g}$ for subgroup chain $O \supset D_4 \supset C_4$

$$\mathbf{P}_{m_4 m_4}^\mu = \mathbf{p}^{m_4} \mathbf{P}^\mu = \mathbf{P}^\mu \mathbf{p}^{m_4}$$

$$= \sum_g \frac{\ell^\mu}{\circ O} (\chi_g^{\mu*}) \cdot \mathbf{g}(\mathbf{p}^{m_4})$$

(Deriving diagonal irreducible O -representation (“irep”) components $D_{m_4 m_4}^{\mu*}(g)$)

$$= \left(\frac{\ell^\mu}{24} \right) (\chi_1^{\mu*}) \cdot \mathbf{1} \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_z}^{\mu*}) \cdot \rho_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_z + d_{\rho_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z \right) = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) \left(d_{\rho_z}^{m_4} \mathbf{1} + \mathbf{1} \cdot \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{R_z}^{\mu*}) \cdot \mathbf{R}_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{R_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{R}_z + d_{\rho_z}^{m_4} \tilde{\mathbf{R}}_z + d_{R_z}^{m_4} \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{1} \right) =$$

$$+$$

$$\rho_x C_4 = \rho_x \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3 \} \text{Coset}$$

$O: \chi_g^\mu$	$g=1$	O characters			
		\mathbf{r}_{1-4} $\tilde{\mathbf{r}}_{1-4}$	ρ_{xyz}	\mathbf{R}_{xyz} $\tilde{\mathbf{R}}_{xyz}$	\mathbf{i}_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

$$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 \underbrace{e^{\frac{2\pi i m_4 \cdot p}{4}}}_{C_4 \text{ characters}} \mathbf{R}_z^p =$$

$$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$$

}

$$\begin{cases} \mathbf{p}_{0_4} = (\mathbf{1} + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{1_4} = (\mathbf{1} + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{2_4} = (\mathbf{1} - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{3_4} = (\mathbf{1} - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4 \end{cases}$$

$$\mathbf{r}_1 C_4 = \mathbf{r}_1 \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y \} \text{Coset}$$

General development of irep projectors $\mathbf{P}_{m_4 m_4}^\mu = \sum_g \frac{\ell^\mu}{\circ O} D_{m_4 m_4}^{\mu*}(g) \mathbf{g}$ for subgroup chain $O \supset D_4 \supset C_4$

$$\mathbf{P}_{m_4 m_4}^\mu = \mathbf{p}^{m_4} \mathbf{P}^\mu = \mathbf{P}^\mu \mathbf{p}^{m_4}$$

$$= \sum_g \frac{\ell^\mu}{\circ O} (\chi_g^{\mu*}) \cdot \mathbf{g}(\mathbf{p}^{m_4})$$

(Deriving diagonal irreducible O -representation (“irep”) components $D_{m_4 m_4}^{\mu*}(g)$)

$$= \left(\frac{\ell^\mu}{24} \right) (\chi_1^{\mu*}) \cdot \mathbf{1} \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_z}^{\mu*}) \cdot \rho_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_z + d_{\rho_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z \right) = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) \left(d_{\rho_z}^{m_4} \mathbf{1} + \mathbf{1} \cdot \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{R_z}^{\mu*}) \cdot \mathbf{R}_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{R_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{R}_z + d_{\rho_z}^{m_4} \tilde{\mathbf{R}}_z + d_{R_z}^{m_4} \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{1} \right) = \left(\frac{\ell^\mu \chi_{R_z}^{\mu*}}{96} \right) \left(d_{\tilde{R}_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \rho_z + \mathbf{1} \cdot \mathbf{R}_z + d_{\rho_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+$$

$$\rho_x C_4 = \rho_x \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3 \} \text{Coset}$$

$O: \chi_g^\mu$	$\mathbf{g}=\mathbf{1}$	O characters				\mathbf{i}_{1-6}
		\mathbf{r}_{1-4} $\tilde{\mathbf{r}}_{1-4}$	ρ_{xyz}	\mathbf{R}_{xyz} $\tilde{\mathbf{R}}_{xyz}$		
$\mu=A_1$	1	1	1	1	1	$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 \underbrace{e^{\frac{2\pi i m_4 \cdot p}{4}}}_{C_4 \text{ characters}} \mathbf{R}_z^p = \begin{cases} \mathbf{p}_{0_4} = (\mathbf{1} + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{1_4} = (\mathbf{1} + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{2_4} = (\mathbf{1} - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{3_4} = (\mathbf{1} - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4 \end{cases}$ $d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$
A_2	1	1	1	-1	-1	
E	2	-1	2	0	0	
T_1	3	0	-1	1	-1	
T_2	3	0	-1	-1	1	

$$\mathbf{r}_1 C_4 = \mathbf{r}_1 \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y \} \text{Coset}$$

General development of irep projectors $\mathbf{P}_{m_4 m_4}^\mu = \sum_g \frac{\ell^\mu}{\circ O} D_{m_4 m_4}^{\mu*}(g) \mathbf{g}$ for subgroup chain $O \supset D_4 \supset C_4$

$$\mathbf{P}_{m_4 m_4}^\mu = \mathbf{p}^{m_4} \mathbf{P}^\mu = \mathbf{P}^\mu \mathbf{p}^{m_4}$$

(Deriving diagonal irreducible O -representation (“irep”) components $D_{m_4 m_4}^{\mu*}(g)$)

$$= \sum_g \frac{\ell^\mu}{\circ O} (\chi_g^{\mu*}) \cdot \mathbf{g}(\mathbf{p}^{m_4})$$

$$= \left(\frac{\ell^\mu}{24} \right) (\chi_1^{\mu*}) \cdot \mathbf{1} (\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z) \frac{1}{4} = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) (\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z) = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) (\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_z}^{\mu*}) \cdot \rho_z (\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) (\mathbf{1} \cdot \rho_z + d_{\rho_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z) = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) (d_{\rho_z}^{m_4} \mathbf{1} + \mathbf{1} \cdot \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{R_z}^{\mu*}) \cdot \mathbf{R}_z (\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{R_z}^{\mu*}}{96} \right) (\mathbf{1} \cdot \mathbf{R}_z + d_{\rho_z}^{m_4} \tilde{\mathbf{R}}_z + d_{R_z}^{m_4} \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{1}) = \left(\frac{\ell^\mu \chi_{R_z}^{\mu*}}{96} \right) (d_{\tilde{R}_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \rho_z + \mathbf{1} \cdot \mathbf{R}_z + d_{\rho_z}^{m_4} \tilde{\mathbf{R}}_z)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\tilde{R}_z}^{\mu*}) \cdot \tilde{\mathbf{R}}_z (\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\tilde{R}_z}^{\mu*}}{96} \right) (\mathbf{1} \cdot \tilde{\mathbf{R}}_z + d_{\rho_z}^{m_4} \mathbf{R}_z + d_{R_z}^{m_4} \mathbf{1} + d_{\tilde{R}_z}^{m_4} \rho_z) = \left(\frac{\ell^\mu \chi_{\tilde{R}_z}^{\mu*}}{96} \right) (d_{R_z}^{m_4} \mathbf{1} + d_{\tilde{R}_z}^{m_4} \rho_z + d_{\rho_z}^{m_4} \mathbf{R}_z + \mathbf{1} \cdot \tilde{\mathbf{R}}_z)$$

$$\rho_x C_4 = \rho_x \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3 \} \text{Coset}$$

$O: \chi_g^\mu$	$g=1$	O characters			
		\mathbf{r}_{1-4} $\tilde{\mathbf{r}}_{1-4}$	ρ_{xyz}	\mathbf{R}_{xyz} $\tilde{\mathbf{R}}_{xyz}$	\mathbf{i}_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

$$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 \underbrace{e^{\frac{2\pi i m_4 \cdot p}{4}}}_{C_4 \text{ characters}} \mathbf{R}_z^p =$$

$$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$$

}

$$\begin{cases} \mathbf{p}_{0_4} = (\mathbf{1} + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{1_4} = (\mathbf{1} + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{2_4} = (\mathbf{1} - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{3_4} = (\mathbf{1} - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4 \end{cases}$$

$$\mathbf{r}_1 C_4 = \mathbf{r}_1 \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y \} \text{Coset}$$

General development of irep projectors $\mathbf{P}_{m_4 m_4}^\mu = \sum_g \frac{\ell^\mu}{\circ O} D_{m_4 m_4}^{\mu*}(g) \mathbf{g}$ for subgroup chain $O \supset D_4 \supset C_4$

$$\mathbf{P}_{m_4 m_4}^\mu = \mathbf{p}^{m_4} \mathbf{P}^\mu = \mathbf{P}^\mu \mathbf{p}^{m_4}$$

(Deriving diagonal irreducible O -representation (“irep”) components $D_{m_4 m_4}^{\mu*}(g)$)

$$= \sum_g \frac{\ell^\mu}{\circ O} (\chi_g^{\mu*}) \cdot \mathbf{g}(\mathbf{p}^{m_4})$$

$$= \left(\frac{\ell^\mu}{24} \right) (\chi_1^{\mu*}) \cdot \mathbf{1} \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_z}^{\mu*}) \cdot \rho_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_z + d_{\rho_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z \right) = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) \left(d_{\rho_z}^{m_4} \mathbf{1} + \mathbf{1} \cdot \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{R_z}^{\mu*}) \cdot \mathbf{R}_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{R_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{R}_z + d_{\rho_z}^{m_4} \tilde{\mathbf{R}}_z + d_{R_z}^{m_4} \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{1} \right) = \left(\frac{\ell^\mu \chi_{R_z}^{\mu*}}{96} \right) \left(d_{\tilde{R}_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \rho_z + \mathbf{1} \cdot \mathbf{R}_z + d_{\rho_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\tilde{R}_z}^{\mu*}) \cdot \tilde{\mathbf{R}}_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\tilde{R}_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \tilde{\mathbf{R}}_z + d_{\rho_z}^{m_4} \mathbf{R}_z + d_{R_z}^{m_4} \mathbf{1} + d_{\tilde{R}_z}^{m_4} \rho_z \right) = \left(\frac{\ell^\mu \chi_{\tilde{R}_z}^{\mu*}}{96} \right) \left(d_{R_z}^{m_4} \mathbf{1} + d_{\tilde{R}_z}^{m_4} \rho_z + d_{\rho_z}^{m_4} \mathbf{R}_z + \mathbf{1} \cdot \tilde{\mathbf{R}}_z \right)$$

$$\rho_x C_4 = \rho_x \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3 \} \text{Coset}$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_x}^{\mu*}) \cdot \rho_x \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4}$$

		O characters				
		$\mathbf{g}=\mathbf{1}$	\mathbf{r}_{1-4}	ρ_{xyz}	\mathbf{R}_{xyz}	\mathbf{i}_{1-6}
		$O: \chi_g^\mu$	$\tilde{\mathbf{r}}_{1-4}$	$\tilde{\mathbf{R}}_{xyz}$		
+	$\mu=A_1$	1	1	1	1	1
+	A_2	1	1	1	-1	-1
	E	2	-1	2	0	0
	T_1	3	0	-1	1	-1
	T_2	3	0	-1	-1	1

$$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 \underbrace{e^{\frac{2\pi i m_4 \cdot p}{4}}}_{C_4 \text{ characters}} \mathbf{R}_z^p =$$

$$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$$

}

$$\begin{cases} \mathbf{p}_{0_4} = (\mathbf{1} + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{1_4} = (\mathbf{1} + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{2_4} = (\mathbf{1} - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{3_4} = (\mathbf{1} - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4 \end{cases}$$

$$\mathbf{r}_1 C_4 = \mathbf{r}_1 \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y \} \text{Coset}$$

General development of irep projectors $\mathbf{P}_{m_4 m_4}^\mu = \sum_g \frac{\ell^\mu}{\circ O} D_{m_4 m_4}^{\mu*}(g) \mathbf{g}$ for subgroup chain $O \supset D_4 \supset C_4$

$$\mathbf{P}_{m_4 m_4}^\mu = \mathbf{p}^{m_4} \mathbf{P}^\mu = \mathbf{P}^\mu \mathbf{p}^{m_4}$$

(Deriving diagonal irreducible O -representation (“irep”) components $D_{m_4 m_4}^{\mu*}(g)$)

$$= \sum_g \frac{\ell^\mu}{\circ O} (\chi_g^{\mu*}) \cdot \mathbf{g}(\mathbf{p}^{m_4})$$

$$= \left(\frac{\ell^\mu}{24} \right) (\chi_1^{\mu*}) \cdot \mathbf{1} (\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z) \frac{1}{4} = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) (\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z) = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) (\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_z}^{\mu*}) \cdot \rho_z (\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) (\mathbf{1} \cdot \rho_z + d_{\rho_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z) = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) (d_{\rho_z}^{m_4} \mathbf{1} + \mathbf{1} \cdot \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{R_z}^{\mu*}) \cdot \mathbf{R}_z (\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{R_z}^{\mu*}}{96} \right) (\mathbf{1} \cdot \mathbf{R}_z + d_{\rho_z}^{m_4} \tilde{\mathbf{R}}_z + d_{R_z}^{m_4} \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{1}) = \left(\frac{\ell^\mu \chi_{R_z}^{\mu*}}{96} \right) (d_{\tilde{R}_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \rho_z + \mathbf{1} \cdot \mathbf{R}_z + d_{\rho_z}^{m_4} \tilde{\mathbf{R}}_z)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\tilde{R}_z}^{\mu*}) \cdot \tilde{\mathbf{R}}_z (\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\tilde{R}_z}^{\mu*}}{96} \right) (\mathbf{1} \cdot \tilde{\mathbf{R}}_z + d_{\rho_z}^{m_4} \mathbf{R}_z + d_{R_z}^{m_4} \mathbf{1} + d_{\tilde{R}_z}^{m_4} \rho_z) = \left(\frac{\ell^\mu \chi_{\tilde{R}_z}^{\mu*}}{96} \right) (d_{R_z}^{m_4} \mathbf{1} + d_{\tilde{R}_z}^{m_4} \rho_z + d_{\rho_z}^{m_4} \mathbf{R}_z + \mathbf{1} \cdot \tilde{\mathbf{R}}_z)$$

$$\rho_x C_4 = \rho_x \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3 \} \text{Coset}$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_x}^{\mu*}) \cdot \rho_x (\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_x}^{\mu*}}{96} \right) (\mathbf{1} \cdot \rho_x + d_{\rho_z}^{m_4} \rho_y + d_{R_z}^{m_4} \mathbf{i}_4 + d_{\tilde{R}_z}^{m_4} \mathbf{i}_3) =$$

	$O: \chi_g^\mu$	O characters				
		$\mathbf{g}=\mathbf{1}$	\mathbf{r}_{1-4} $\tilde{\mathbf{r}}_{1-4}$	ρ_{xyz}	\mathbf{R}_{xyz} $\tilde{\mathbf{R}}_{xyz}$	\mathbf{i}_{1-6}
+	$\mu=A_1$	1	1	1	1	1
+	A_2	1	1	1	-1	-1
+	E	2	-1	2	0	0
+	T_1	3	0	-1	1	-1
+	T_2	3	0	-1	-1	1

$$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 \underbrace{e^{\frac{2\pi i m_4 \cdot p}{4}}}_{C_4 \text{ characters}} \mathbf{R}_z^p =$$

$$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$$

}

$$\begin{cases} \mathbf{p}_{0_4} = (\mathbf{1} + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{1_4} = (\mathbf{1} + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{2_4} = (\mathbf{1} - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{3_4} = (\mathbf{1} - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4 \end{cases}$$

$$\mathbf{r}_1 C_4 = \mathbf{r}_1 \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y \} \text{Coset}$$

General development of irep projectors $\mathbf{P}_{m_4 m_4}^\mu = \sum_g \frac{\ell^\mu}{\circ O} D_{m_4 m_4}^{\mu*}(g) \mathbf{g}$ for subgroup chain $O \supset D_4 \supset C_4$

$$\mathbf{P}_{m_4 m_4}^\mu = \mathbf{p}^{m_4} \mathbf{P}^\mu = \mathbf{P}^\mu \mathbf{p}^{m_4}$$

(Deriving diagonal irreducible O -representation (“irep”) components $D_{m_4 m_4}^{\mu*}(g)$)

$$= \sum_g \frac{\ell^\mu}{\circ O} (\chi_g^{\mu*}) \cdot \mathbf{g}(\mathbf{p}^{m_4})$$

$$= \left(\frac{\ell^\mu}{24} \right) (\chi_1^{\mu*}) \cdot \mathbf{1} \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_z}^{\mu*}) \cdot \rho_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_z + d_{\rho_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z \right) = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) \left(d_{\rho_z}^{m_4} \mathbf{1} + \mathbf{1} \cdot \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{R_z}^{\mu*}) \cdot \mathbf{R}_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{R_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{R}_z + d_{\rho_z}^{m_4} \tilde{\mathbf{R}}_z + d_{R_z}^{m_4} \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{1} \right) = \left(\frac{\ell^\mu \chi_{R_z}^{\mu*}}{96} \right) \left(d_{\tilde{R}_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \rho_z + \mathbf{1} \cdot \mathbf{R}_z + d_{\rho_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\tilde{R}_z}^{\mu*}) \cdot \tilde{\mathbf{R}}_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\tilde{R}_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \tilde{\mathbf{R}}_z + d_{\rho_z}^{m_4} \mathbf{R}_z + d_{R_z}^{m_4} \mathbf{1} + d_{\tilde{R}_z}^{m_4} \rho_z \right) = \left(\frac{\ell^\mu \chi_{\tilde{R}_z}^{\mu*}}{96} \right) \left(d_{R_z}^{m_4} \mathbf{1} + d_{\tilde{R}_z}^{m_4} \rho_z + d_{\rho_z}^{m_4} \mathbf{R}_z + \mathbf{1} \cdot \tilde{\mathbf{R}}_z \right)$$

$$\rho_x C_4 = \rho_x \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3 \} \text{Coset}$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_x}^{\mu*}) \cdot \rho_x \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_x}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_x + d_{\rho_z}^{m_4} \rho_y + d_{R_z}^{m_4} \mathbf{i}_4 + d_{\tilde{R}_z}^{m_4} \mathbf{i}_3 \right) = \left(\frac{\ell^\mu \chi_{\rho_x}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_x + d_{\rho_z}^{m_4} \rho_y + d_{R_z}^{m_4} \mathbf{i}_4 + d_{\tilde{R}_z}^{m_4} \mathbf{i}_3 \right)$$

O characters

$O: \chi_g^\mu$	$g=1$	\mathbf{r}_{1-4}	ρ_{xyz}	\mathbf{R}_{xyz}	\mathbf{i}_{1-6}
		$\tilde{\mathbf{r}}_{1-4}$		$\tilde{\mathbf{R}}_{xyz}$	
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

$$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 \underbrace{e^{\frac{2\pi i m_4 \cdot p}{4}}}_{C_4 \text{ characters}} \mathbf{R}_z^p =$$

$$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$$

}

$$\begin{cases} \mathbf{p}_{0_4} = (\mathbf{1} + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{1_4} = (\mathbf{1} + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{2_4} = (\mathbf{1} - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{3_4} = (\mathbf{1} - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4 \end{cases}$$

$$\mathbf{r}_1 C_4 = \mathbf{r}_1 \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y \} \text{Coset}$$

General development of irep projectors $\mathbf{P}_{m_4 m_4}^\mu = \sum_g \frac{\ell^\mu}{\circ O} D_{m_4 m_4}^{\mu*}(g) \mathbf{g}$ for subgroup chain $O \supset D_4 \supset C_4$

$$\mathbf{P}_{m_4 m_4}^\mu = \mathbf{p}^{m_4} \mathbf{P}^\mu = \mathbf{P}^\mu \mathbf{p}^{m_4}$$

(Deriving diagonal irreducible O -representation (“irep”) components $D_{m_4 m_4}^{\mu*}(g)$)

$$= \sum_g \frac{\ell^\mu}{\circ O} (\chi_g^{\mu*}) \cdot \mathbf{g}(\mathbf{p}^{m_4})$$

$$= \left(\frac{\ell^\mu}{24} \right) (\chi_1^{\mu*}) \cdot \mathbf{1} \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_z}^{\mu*}) \cdot \rho_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_z + d_{\rho_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z \right) = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) \left(d_{\rho_z}^{m_4} \mathbf{1} + \mathbf{1} \cdot \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{R_z}^{\mu*}) \cdot \mathbf{R}_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{R_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{R}_z + d_{\rho_z}^{m_4} \tilde{\mathbf{R}}_z + d_{R_z}^{m_4} \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{1} \right) = \left(\frac{\ell^\mu \chi_{R_z}^{\mu*}}{96} \right) \left(d_{\tilde{R}_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \rho_z + \mathbf{1} \cdot \mathbf{R}_z + d_{\rho_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\tilde{R}_z}^{\mu*}) \cdot \tilde{\mathbf{R}}_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\tilde{R}_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \tilde{\mathbf{R}}_z + d_{\rho_z}^{m_4} \mathbf{R}_z + d_{R_z}^{m_4} \mathbf{1} + d_{\tilde{R}_z}^{m_4} \rho_z \right) = \left(\frac{\ell^\mu \chi_{\tilde{R}_z}^{\mu*}}{96} \right) \left(d_{R_z}^{m_4} \mathbf{1} + d_{\tilde{R}_z}^{m_4} \rho_z + d_{\rho_z}^{m_4} \mathbf{R}_z + \mathbf{1} \cdot \tilde{\mathbf{R}}_z \right)$$

$$\rho_x C_4 = \rho_x \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3 \} \text{Coset}$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_x}^{\mu*}) \cdot \rho_x \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_x}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_x + d_{\rho_z}^{m_4} \rho_y + d_{R_z}^{m_4} \mathbf{i}_4 + d_{\tilde{R}_z}^{m_4} \mathbf{i}_3 \right) = \left(\frac{\ell^\mu \chi_{\rho_x}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_x + d_{\rho_z}^{m_4} \rho_y + d_{R_z}^{m_4} \mathbf{i}_4 + d_{\tilde{R}_z}^{m_4} \mathbf{i}_3 \right)$$

O characters

$O: \chi_g^\mu$	$g=1$	\mathbf{r}_{1-4}	ρ_{xyz}	\mathbf{R}_{xyz}	\mathbf{i}_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 e^{\frac{2\pi i m_4 \cdot p}{4}} \mathbf{R}_z^p =$

$\underbrace{\hspace{10em}}_{C_4 \text{ characters}}$

$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$

$\left\{ \begin{array}{l} \mathbf{p}_{0_4} = (\mathbf{1} + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{1_4} = (\mathbf{1} + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{2_4} = (\mathbf{1} - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{3_4} = (\mathbf{1} - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4 \end{array} \right.$

$$\mathbf{r}_1 C_4 = \mathbf{r}_1 \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y \} \text{Coset}$$

General development of irep projectors $\mathbf{P}_{m_4 m_4}^\mu = \sum_g \frac{\ell^\mu}{\circ O} D_{m_4 m_4}^{\mu*}(g) \mathbf{g}$ for subgroup chain $O \supset D_4 \supset C_4$

$$\mathbf{P}_{m_4 m_4}^\mu = \mathbf{p}^{m_4} \mathbf{P}^\mu = \mathbf{P}^\mu \mathbf{p}^{m_4}$$

(Deriving diagonal irreducible O -representation (“irep”) components $D_{m_4 m_4}^{\mu*}(g)$)

$$= \sum_g \frac{\ell^\mu}{\circ O} (\chi_g^{\mu*}) \cdot \mathbf{g}(\mathbf{p}^{m_4})$$

$$= \left(\frac{\ell^\mu}{24} \right) (\chi_1^{\mu*}) \cdot \mathbf{1} \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_z}^{\mu*}) \cdot \rho_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_z + d_{\rho_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z \right) = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) \left(d_{\rho_z}^{m_4} \mathbf{1} + \mathbf{1} \cdot \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{R_z}^{\mu*}) \cdot \mathbf{R}_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{R_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{R}_z + d_{\rho_z}^{m_4} \tilde{\mathbf{R}}_z + d_{R_z}^{m_4} \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{1} \right) = \left(\frac{\ell^\mu \chi_{R_z}^{\mu*}}{96} \right) \left(d_{\tilde{R}_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \rho_z + \mathbf{1} \cdot \mathbf{R}_z + d_{\rho_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\tilde{R}_z}^{\mu*}) \cdot \tilde{\mathbf{R}}_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\tilde{R}_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \tilde{\mathbf{R}}_z + d_{\rho_z}^{m_4} \mathbf{R}_z + d_{R_z}^{m_4} \mathbf{1} + d_{\tilde{R}_z}^{m_4} \rho_z \right) = \left(\frac{\ell^\mu \chi_{\tilde{R}_z}^{\mu*}}{96} \right) \left(d_{R_z}^{m_4} \mathbf{1} + d_{\tilde{R}_z}^{m_4} \rho_z + d_{\rho_z}^{m_4} \mathbf{R}_z + \mathbf{1} \cdot \tilde{\mathbf{R}}_z \right)$$

$$\rho_x C_4 = \rho_x \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3 \} \text{Coset}$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_x}^{\mu*}) \cdot \rho_x \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_x}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_x + d_{\rho_z}^{m_4} \rho_y + d_{R_z}^{m_4} \mathbf{i}_4 + d_{\tilde{R}_z}^{m_4} \mathbf{i}_3 \right) = \left(\frac{\ell^\mu \chi_{\rho_x}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_x + d_{\rho_z}^{m_4} \rho_y + d_{R_z}^{m_4} \mathbf{i}_4 + d_{\tilde{R}_z}^{m_4} \mathbf{i}_3 \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_y}^{\mu*}) \cdot \rho_y \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_y}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_y + d_{\rho_z}^{m_4} \rho_x + d_{R_z}^{m_4} \mathbf{i}_3 + d_{\tilde{R}_z}^{m_4} \mathbf{i}_4 \right) =$$

$$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 e^{\frac{2\pi i m_4 \cdot p}{4}} \mathbf{R}_z^p = \begin{cases} \mathbf{p}_{0_4} = (\mathbf{1} + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{1_4} = (\mathbf{1} + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{2_4} = (\mathbf{1} - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{3_4} = (\mathbf{1} - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4 \end{cases}$$

C_4 characters

$$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$$

$$\mathbf{r}_1 C_4 = \mathbf{r}_1 \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y \} \text{Coset}$$

General development of irep projectors $\mathbf{P}_{m_4 m_4}^\mu = \sum_g \frac{\ell^\mu}{\circ O} D_{m_4 m_4}^{\mu*}(g) \mathbf{g}$ for subgroup chain $O \supset D_4 \supset C_4$

$$\mathbf{P}_{m_4 m_4}^\mu = \mathbf{p}^{m_4} \mathbf{P}^\mu = \mathbf{P}^\mu \mathbf{p}^{m_4}$$

(Deriving diagonal irreducible O -representation (“irep”) components $D_{m_4 m_4}^{\mu*}(g)$)

$$= \sum_g \frac{\ell^\mu}{\circ O} (\chi_g^{\mu*}) \cdot \mathbf{g}(\mathbf{p}^{m_4})$$

$$= \left(\frac{\ell^\mu}{24} \right) (\chi_1^{\mu*}) \cdot \mathbf{1} \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_z}^{\mu*}) \cdot \rho_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_z + d_{\rho_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z \right) = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) \left(d_{\rho_z}^{m_4} \mathbf{1} + \mathbf{1} \cdot \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$


$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{R_z}^{\mu*}) \cdot \mathbf{R}_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{R_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{R}_z + d_{\rho_z}^{m_4} \tilde{\mathbf{R}}_z + d_{R_z}^{m_4} \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{1} \right) = \left(\frac{\ell^\mu \chi_{R_z}^{\mu*}}{96} \right) \left(d_{\tilde{R}_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \rho_z + \mathbf{1} \cdot \mathbf{R}_z + d_{\rho_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\tilde{R}_z}^{\mu*}) \cdot \tilde{\mathbf{R}}_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\tilde{R}_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \tilde{\mathbf{R}}_z + d_{\rho_z}^{m_4} \mathbf{R}_z + d_{R_z}^{m_4} \mathbf{1} + d_{\tilde{R}_z}^{m_4} \rho_z \right) = \left(\frac{\ell^\mu \chi_{\tilde{R}_z}^{\mu*}}{96} \right) \left(d_{R_z}^{m_4} \mathbf{1} + d_{\tilde{R}_z}^{m_4} \rho_z + d_{\rho_z}^{m_4} \mathbf{R}_z + \mathbf{1} \cdot \tilde{\mathbf{R}}_z \right)$$

$$\rho_x C_4 = \rho_x \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3 \} \text{Coset}$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_x}^{\mu*}) \cdot \rho_x \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_x}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_x + d_{\rho_z}^{m_4} \rho_y + d_{R_z}^{m_4} \mathbf{i}_4 + d_{\tilde{R}_z}^{m_4} \mathbf{i}_3 \right) = \left(\frac{\ell^\mu \chi_{\rho_x}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_x + d_{\rho_z}^{m_4} \rho_y + d_{R_z}^{m_4} \mathbf{i}_4 + d_{\tilde{R}_z}^{m_4} \mathbf{i}_3 \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_y}^{\mu*}) \cdot \rho_y \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_y}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_y + d_{\rho_z}^{m_4} \rho_x + d_{R_z}^{m_4} \mathbf{i}_3 + d_{\tilde{R}_z}^{m_4} \mathbf{i}_4 \right) = \left(\frac{\ell^\mu \chi_{\rho_y}^{\mu*}}{96} \right) \left(d_{\rho_z}^{m_4} \rho_x + \mathbf{1} \cdot \rho_y + d_{\tilde{R}_z}^{m_4} \mathbf{i}_4 + d_{R_z}^{m_4} \mathbf{i}_3 \right)$$



$$\mathbf{r}_1 C_4 = \mathbf{r}_1 \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y \} \text{Coset}$$

General development of irep projectors $\mathbf{P}_{m_4 m_4}^\mu = \sum_g \frac{\ell^\mu}{\circ O} D_{m_4 m_4}^{\mu*}(g) \mathbf{g}$ for subgroup chain $O \supset D_4 \supset C_4$

$$\mathbf{P}_{m_4 m_4}^\mu = \mathbf{p}^{m_4} \mathbf{P}^\mu = \mathbf{P}^\mu \mathbf{p}^{m_4}$$

(Deriving diagonal irreducible O -representation (“irep”) components $D_{m_4 m_4}^{\mu*}(g)$)

$$= \sum_g \frac{\ell^\mu}{\circ O} (\chi_g^{\mu*}) \cdot \mathbf{g}(\mathbf{p}^{m_4})$$

$$= \left(\frac{\ell^\mu}{24} \right) (\chi_1^{\mu*}) \cdot \mathbf{1} \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_z}^{\mu*}) \cdot \rho_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_z + d_{\rho_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z \right) = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) \left(d_{\rho_z}^{m_4} \mathbf{1} + \mathbf{1} \cdot \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{R_z}^{\mu*}) \cdot \mathbf{R}_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{R_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{R}_z + d_{\rho_z}^{m_4} \tilde{\mathbf{R}}_z + d_{R_z}^{m_4} \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{1} \right) = \left(\frac{\ell^\mu \chi_{R_z}^{\mu*}}{96} \right) \left(d_{\tilde{R}_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \rho_z + \mathbf{1} \cdot \mathbf{R}_z + d_{\rho_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\tilde{R}_z}^{\mu*}) \cdot \tilde{\mathbf{R}}_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\tilde{R}_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \tilde{\mathbf{R}}_z + d_{\rho_z}^{m_4} \mathbf{R}_z + d_{R_z}^{m_4} \mathbf{1} + d_{\tilde{R}_z}^{m_4} \rho_z \right) = \left(\frac{\ell^\mu \chi_{\tilde{R}_z}^{\mu*}}{96} \right) \left(d_{R_z}^{m_4} \mathbf{1} + d_{\tilde{R}_z}^{m_4} \rho_z + d_{\rho_z}^{m_4} \mathbf{R}_z + \mathbf{1} \cdot \tilde{\mathbf{R}}_z \right)$$

$$\rho_x C_4 = \rho_x \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3 \} \text{Coset}$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_x}^{\mu*}) \cdot \rho_x \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_x}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_x + d_{\rho_z}^{m_4} \rho_y + d_{R_z}^{m_4} \mathbf{i}_4 + d_{\tilde{R}_z}^{m_4} \mathbf{i}_3 \right) = \left(\frac{\ell^\mu \chi_{\rho_x}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_x + d_{\rho_z}^{m_4} \rho_y + d_{R_z}^{m_4} \mathbf{i}_4 + d_{\tilde{R}_z}^{m_4} \mathbf{i}_3 \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_y}^{\mu*}) \cdot \rho_y \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_y}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_y + d_{\rho_z}^{m_4} \rho_x + d_{R_z}^{m_4} \mathbf{i}_3 + d_{\tilde{R}_z}^{m_4} \mathbf{i}_4 \right) = \left(\frac{\ell^\mu \chi_{\rho_y}^{\mu*}}{96} \right) \left(d_{\rho_z}^{m_4} \rho_x + \mathbf{1} \cdot \rho_y + d_{\tilde{R}_z}^{m_4} \mathbf{i}_4 + d_{R_z}^{m_4} \mathbf{i}_3 \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\mathbf{i}_4}^{\mu*}) \cdot \mathbf{i}_4 \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\mathbf{i}_4}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{i}_4 + d_{\rho_z}^{m_4} \mathbf{i}_3 + d_{R_z}^{m_4} \rho_y + d_{\tilde{R}_z}^{m_4} \rho_x \right) = \left(\frac{\ell^\mu \chi_{\mathbf{i}_4}^{\mu*}}{96} \right) \left(d_{\tilde{R}_z}^{m_4} \rho_x + d_{R_z}^{m_4} \rho_y + \mathbf{1} \cdot \mathbf{i}_4 + d_{\rho_z}^{m_4} \mathbf{i}_3 \right)$$

$$+$$

$$\mathbf{r}_1 C_4 = \mathbf{r}_1 \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y \} \text{Coset}$$

General development of irep projectors $\mathbf{P}_{m_4 m_4}^\mu = \sum_g \frac{\ell^\mu}{\circ O} D_{m_4 m_4}^{\mu*}(g) \mathbf{g}$ for subgroup chain $O \supset D_4 \supset C_4$

$$\mathbf{P}_{m_4 m_4}^\mu = \mathbf{p}^{m_4} \mathbf{P}^\mu = \mathbf{P}^\mu \mathbf{p}^{m_4}$$

(Deriving diagonal irreducible O -representation (“irep”) components $D_{m_4 m_4}^{\mu*}(g)$)

$$= \sum_g \frac{\ell^\mu}{\circ O} (\chi_g^{\mu*}) \cdot \mathbf{g}(\mathbf{p}^{m_4})$$

$$= \left(\frac{\ell^\mu}{24} \right) (\chi_1^{\mu*}) \cdot \mathbf{1} \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) = \left(\frac{\ell^\mu \chi_1^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_z}^{\mu*}) \cdot \rho_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_z + d_{\rho_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z \right) = \left(\frac{\ell^\mu \chi_{\rho_z}^{\mu*}}{96} \right) \left(d_{\rho_z}^{m_4} \mathbf{1} + \mathbf{1} \cdot \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{R}_z + d_{R_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{R_z}^{\mu*}) \cdot \mathbf{R}_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{R_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{R}_z + d_{\rho_z}^{m_4} \tilde{\mathbf{R}}_z + d_{R_z}^{m_4} \rho_z + d_{\tilde{R}_z}^{m_4} \mathbf{1} \right) = \left(\frac{\ell^\mu \chi_{R_z}^{\mu*}}{96} \right) \left(d_{\tilde{R}_z}^{m_4} \mathbf{1} + d_{R_z}^{m_4} \rho_z + \mathbf{1} \cdot \mathbf{R}_z + d_{\rho_z}^{m_4} \tilde{\mathbf{R}}_z \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\tilde{R}_z}^{\mu*}) \cdot \tilde{\mathbf{R}}_z \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\tilde{R}_z}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \tilde{\mathbf{R}}_z + d_{\rho_z}^{m_4} \mathbf{R}_z + d_{R_z}^{m_4} \mathbf{1} + d_{\tilde{R}_z}^{m_4} \rho_z \right) = \left(\frac{\ell^\mu \chi_{\tilde{R}_z}^{\mu*}}{96} \right) \left(d_{R_z}^{m_4} \mathbf{1} + d_{\tilde{R}_z}^{m_4} \rho_z + d_{\rho_z}^{m_4} \mathbf{R}_z + \mathbf{1} \cdot \tilde{\mathbf{R}}_z \right)$$

$$\rho_x C_4 = \rho_x \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3 \} \text{Coset}$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_x}^{\mu*}) \cdot \rho_x \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_x}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_x + d_{\rho_z}^{m_4} \rho_y + d_{R_z}^{m_4} \mathbf{i}_4 + d_{\tilde{R}_z}^{m_4} \mathbf{i}_3 \right) = \left(\frac{\ell^\mu \chi_{\rho_x}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_x + d_{\rho_z}^{m_4} \rho_y + d_{R_z}^{m_4} \mathbf{i}_4 + d_{\tilde{R}_z}^{m_4} \mathbf{i}_3 \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\rho_y}^{\mu*}) \cdot \rho_y \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\rho_y}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \rho_y + d_{\rho_z}^{m_4} \rho_x + d_{R_z}^{m_4} \mathbf{i}_3 + d_{\tilde{R}_z}^{m_4} \mathbf{i}_4 \right) = \left(\frac{\ell^\mu \chi_{\rho_y}^{\mu*}}{96} \right) \left(d_{\rho_z}^{m_4} \rho_x + \mathbf{1} \cdot \rho_y + d_{\tilde{R}_z}^{m_4} \mathbf{i}_4 + d_{R_z}^{m_4} \mathbf{i}_3 \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\mathbf{i}_4}^{\mu*}) \cdot \mathbf{i}_4 \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\mathbf{i}_4}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{i}_4 + d_{\rho_z}^{m_4} \mathbf{i}_3 + d_{R_z}^{m_4} \rho_y + d_{\tilde{R}_z}^{m_4} \rho_x \right) = \left(\frac{\ell^\mu \chi_{\mathbf{i}_4}^{\mu*}}{96} \right) \left(d_{\tilde{R}_z}^{m_4} \rho_x + d_{R_z}^{m_4} \rho_y + \mathbf{1} \cdot \mathbf{i}_4 + d_{\rho_z}^{m_4} \mathbf{i}_3 \right)$$

$$+ \left(\frac{\ell^\mu}{24} \right) (\chi_{\mathbf{i}_3}^{\mu*}) \cdot \mathbf{i}_3 \left(\mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{R_z}^{m_4} \mathbf{R}_z + d_{\tilde{R}_z}^{m_4} \tilde{\mathbf{R}}_z \right) \frac{1}{4} = \left(\frac{\ell^\mu \chi_{\mathbf{i}_3}^{\mu*}}{96} \right) \left(\mathbf{1} \cdot \mathbf{i}_3 + d_{\rho_z}^{m_4} \mathbf{i}_4 + d_{R_z}^{m_4} \mathbf{1} + d_{\tilde{R}_z}^{m_4} \rho_y \right) = \left(\frac{\ell^\mu \chi_{\mathbf{i}_3}^{\mu*}}{96} \right) \left(d_{R_z}^{m_4} \rho_x + d_{\tilde{R}_z}^{m_4} \rho_y + d_{\rho_z}^{m_4} \mathbf{i}_4 + \mathbf{1} \cdot \mathbf{i}_3 \right)$$

$$\mathbf{r}_1 C_4 = \mathbf{r}_1 \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} = \{ \mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y \} \text{Coset}$$

etc. etc.

General development of irep projectors $\mathbf{P}_{m_4 m_4}^\mu = \sum_g \frac{\ell^\mu}{\circ O} D_{m_4 m_4}^{\mu*}(g) \mathbf{g}$ for subgroup chain $O \supset D_4 \supset C_4$

$\mathbf{P}_{m_4 m_4}^\mu = \mathbf{p}_{m_4} \mathbf{P}^\mu = \mathbf{P}^\mu \mathbf{p}_{m_4}$ (Deriving diagonal irreducible O -representation (“irep”) components $D_{m_4 m_4}^{\mu*}(g)$)

$$= \sum_g \frac{\ell^\mu}{\circ O} (\chi_g^{\mu*}) \cdot \mathbf{g} \cdot (\mathbf{p}_{m_4}) = \sum_g \frac{\ell^\mu}{4 \circ O} (\chi_g^{\mu*}) \cdot \mathbf{g} \cdot (d_1^{m_4} \mathbf{1} + d_{\rho_z}^{m_4} \rho_z + d_{\mathbf{R}_z}^{m_4} \mathbf{R}_z + d_{\tilde{\mathbf{R}}_z}^{m_4} \tilde{\mathbf{R}}_z)$$

O characters χ_g^μ

$O: \chi_g^\mu$	$\mathbf{g}=\mathbf{1}$	\mathbf{r}_{1-4}^p	ρ_{xyz}	\mathbf{R}_{xyz}^p	\mathbf{i}_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

$$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 e^{\frac{2\pi i m_4 \cdot p}{4}} \mathbf{R}_z^p = \begin{cases} \mathbf{p}_{0_4} = (\mathbf{1} + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{1_4} = (\mathbf{1} + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{2_4} = (\mathbf{1} - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{3_4} = (\mathbf{1} - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4 \end{cases}$$

C_4 characters $d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$

$$\begin{aligned} \mathbf{1}C_4 &= \mathbf{1}\{\mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z\} & \rho_x C_4 &= \{\rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3\} & \mathbf{r}_1 C_4 &= \{\mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y\} & \mathbf{r}_2 C_4 &= \{\mathbf{r}_2, \mathbf{r}_3, \mathbf{i}_2, \tilde{\mathbf{R}}_y\} & \tilde{\mathbf{r}}_1 C_4 &= \{\tilde{\mathbf{r}}_1, \tilde{\mathbf{r}}_3, \tilde{\mathbf{R}}_x, \mathbf{i}_6\} & \tilde{\mathbf{r}}_2 C_4 &= \{\tilde{\mathbf{r}}_2, \tilde{\mathbf{r}}_4, \mathbf{R}_x, \mathbf{i}_5\} \\ &= \frac{\ell^\mu}{96} \chi_{\mathbf{1}}^{\mu*}(1, d_{\rho_z}^{m_4}, d_{\mathbf{R}_z}^{m_4}, d_{\tilde{\mathbf{R}}_z}^{m_4}) + \frac{\ell^\mu}{96} \chi_{\rho_x}^{\mu*}(1, d_{\rho_z}^{m_4}, d_{\mathbf{R}_z}^{m_4}, d_{\tilde{\mathbf{R}}_z}^{m_4}) + \frac{\ell^\mu}{96} \chi_{\mathbf{r}_1}^{\mu*}(1, d_{\rho_z}^{m_4}, d_{\mathbf{R}_z}^{m_4}, d_{\tilde{\mathbf{R}}_z}^{m_4}) + \frac{\ell^\mu}{96} \chi_{\mathbf{r}_2}^{\mu*}(1, d_{\rho_z}^{m_4}, d_{\mathbf{R}_z}^{m_4}, d_{\tilde{\mathbf{R}}_z}^{m_4}) + \frac{\ell^\mu}{96} \chi_{\tilde{\mathbf{r}}_1}^{\mu*}(1, d_{\rho_z}^{m_4}, d_{\mathbf{R}_z}^{m_4}, d_{\tilde{\mathbf{R}}_z}^{m_4}) + \frac{\ell^\mu}{96} \chi_{\tilde{\mathbf{r}}_2}^{\mu*}(1, d_{\rho_z}^{m_4}, d_{\mathbf{R}_z}^{m_4}, d_{\tilde{\mathbf{R}}_z}^{m_4}) \\ &+ \frac{\ell^\mu}{96} \chi_{\rho_z}^{\mu*}(d_{\rho_z}^{m_4}, 1, d_{\tilde{\mathbf{R}}_z}^{m_4}, d_{\mathbf{R}_z}^{m_4}) + \frac{\ell^\mu}{96} \chi_{\rho_y}^{\mu*}(d_{\rho_z}^{m_4}, 1, d_{\tilde{\mathbf{R}}_z}^{m_4}, d_{\mathbf{R}_z}^{m_4}) + \frac{\ell^\mu}{96} \chi_{\mathbf{r}_4}^{\mu*}(d_{\rho_z}^{m_4}, 1, d_{\tilde{\mathbf{R}}_z}^{m_4}, d_{\mathbf{R}_z}^{m_4}) + \frac{\ell^\mu}{96} \chi_{\mathbf{r}_3}^{\mu*}(d_{\rho_z}^{m_4}, 1, d_{\tilde{\mathbf{R}}_z}^{m_4}, d_{\mathbf{R}_z}^{m_4}) + \frac{\ell^\mu}{96} \chi_{\tilde{\mathbf{r}}_3}^{\mu*}(d_{\rho_z}^{m_4}, 1, d_{\tilde{\mathbf{R}}_z}^{m_4}, d_{\mathbf{R}_z}^{m_4}) + \frac{\ell^\mu}{96} \chi_{\tilde{\mathbf{r}}_4}^{\mu*}(d_{\rho_z}^{m_4}, 1, d_{\tilde{\mathbf{R}}_z}^{m_4}, d_{\mathbf{R}_z}^{m_4}) \\ &+ \frac{\ell^\mu}{96} \chi_{\mathbf{R}_z}^{\mu*}(d_{\tilde{\mathbf{R}}_z}^{m_4}, d_{\mathbf{R}_z}^{m_4}, 1, d_{\rho_z}^{m_4}) + \frac{\ell^\mu}{96} \chi_{\mathbf{i}_4}^{\mu*}(d_{\tilde{\mathbf{R}}_z}^{m_4}, d_{\mathbf{R}_z}^{m_4}, 1, d_{\rho_z}^{m_4}) + \frac{\ell^\mu}{96} \chi_{\mathbf{i}_1}^{\mu*}(d_{\tilde{\mathbf{R}}_z}^{m_4}, d_{\mathbf{R}_z}^{m_4}, 1, d_{\rho_z}^{m_4}) + \frac{\ell^\mu}{96} \chi_{\mathbf{i}_2}^{\mu*}(d_{\tilde{\mathbf{R}}_z}^{m_4}, d_{\mathbf{R}_z}^{m_4}, 1, d_{\rho_z}^{m_4}) + \frac{\ell^\mu}{96} \chi_{\tilde{\mathbf{R}}_x}^{\mu*}(d_{\tilde{\mathbf{R}}_z}^{m_4}, d_{\mathbf{R}_z}^{m_4}, 1, d_{\rho_z}^{m_4}) + \frac{\ell^\mu}{96} \chi_{\mathbf{R}_x}^{\mu*}(d_{\tilde{\mathbf{R}}_z}^{m_4}, d_{\mathbf{R}_z}^{m_4}, 1, d_{\rho_z}^{m_4}) \\ &+ \frac{\ell^\mu}{96} \chi_{\tilde{\mathbf{R}}_z}^{\mu*}(d_{\mathbf{R}_z}^{m_4}, d_{\tilde{\mathbf{R}}_z}^{m_4}, d_{\rho_z}^{m_4}, 1) + \frac{\ell^\mu}{96} \chi_{\mathbf{i}_3}^{\mu*}(d_{\mathbf{R}_z}^{m_4}, d_{\tilde{\mathbf{R}}_z}^{m_4}, d_{\rho_z}^{m_4}, 1) + \frac{\ell^\mu}{96} \chi_{\mathbf{R}_y}^{\mu*}(d_{\mathbf{R}_z}^{m_4}, d_{\tilde{\mathbf{R}}_z}^{m_4}, d_{\rho_z}^{m_4}, 1) + \frac{\ell^\mu}{96} \chi_{\tilde{\mathbf{R}}_y}^{\mu*}(d_{\mathbf{R}_z}^{m_4}, d_{\tilde{\mathbf{R}}_z}^{m_4}, d_{\rho_z}^{m_4}, 1) + \frac{\ell^\mu}{96} \chi_{\mathbf{i}_6}^{\mu*}(d_{\mathbf{R}_z}^{m_4}, d_{\tilde{\mathbf{R}}_z}^{m_4}, d_{\rho_z}^{m_4}, 1) + \frac{\ell^\mu}{96} \chi_{\mathbf{i}_5}^{\mu*}(d_{\mathbf{R}_z}^{m_4}, d_{\tilde{\mathbf{R}}_z}^{m_4}, d_{\rho_z}^{m_4}, 1) \end{aligned}$$

Each of 24 columns is a sum of 4 products $\frac{\ell^\mu}{96} \chi_g^{\mu*} d_{\rho_p}^{m_4}$ that gives coefficient $= \frac{\ell^\mu}{\circ O} D_{m_4 m_4}^{\mu*}(g)$ of $\mathbf{P}_{m_4 m_4}^\mu$

$$\frac{1}{96} (\mathbf{1} + \rho_z + \mathbf{R}_z + \tilde{\mathbf{R}}_z + \rho_x + \rho_y + \mathbf{i}_4 + \mathbf{i}_3 + \mathbf{r}_1 + \mathbf{r}_4 + \mathbf{i}_1 + \mathbf{R}_y + \mathbf{r}_2 + \mathbf{r}_3 + \mathbf{i}_2 + \tilde{\mathbf{R}}_y + \tilde{\mathbf{r}}_1 + \tilde{\mathbf{r}}_3 + \tilde{\mathbf{R}}_x + \mathbf{i}_6 + \tilde{\mathbf{r}}_2 + \tilde{\mathbf{r}}_4 + \mathbf{R}_x + \mathbf{i}_5)$$

This $\mathbf{P}_{m_4 m_4}^\mu$ -sum is in order of left cosets $\mathbf{g} \cdot C_4$ of C_4 in O . (Examples follow.)

$$\{\mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z\} \quad \{\rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3\} \quad \{\mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y\} \quad \{\mathbf{r}_2, \mathbf{r}_3, \mathbf{i}_2, \tilde{\mathbf{R}}_y\} \quad \{\tilde{\mathbf{r}}_1, \tilde{\mathbf{r}}_3, \tilde{\mathbf{R}}_x, \mathbf{i}_6\} \quad \{\tilde{\mathbf{r}}_2, \tilde{\mathbf{r}}_4, \mathbf{R}_x, \mathbf{i}_5\}$$

Review Octahedral $O_h \supset O$ group operator structure

Review Octahedral $O_h \supset O \supset D_4 \supset C_4$ subgroup chain correlations

Comparison of $O \supset D_4 \supset C_4$ and $O \supset D_4 \supset D_2$ correlations and level/projector splitting

$O \supset D_4 \supset C_4$ subgroup chain splitting

$O \supset D_4 \supset D_2$ subgroup chain splitting (normal D_2 vs. unnormal D_2)

$O_h \supset O \supset D_4 \supset C_{4v}$ and $O_h \supset O \supset D_4 \supset C_{4v} \supset C_{2v}$ subgroup splitting

Splitting O class projectors P^μ into irreducible projectors $P^\mu_{m_4m_4}$ for $O \supset C_4$

Development of irreducible projectors $P^\mu_{m_4m_4}$ and representations $D^\mu_{m_4m_4}$

Calculating $P^{E_{0404}}$, $P^{E_{2424}}$, $P^{T_{10404}}$, $P^{T_{11414}}$, $P^{T_{22424}}$, $P^{T_{21414}}$,



$O \supset C_4$ induced representation $0_4(C_4) \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples

Elementary induced representation $0_4(C_4) \uparrow O$

Projection reduction of induced representation $0_4(C_4) \uparrow O$

Introduction to ortho-complete eigenvalue-parameter relations

Examples in SF_6 spectroscopy



Calculating $\mathbf{P}^E_{0_40_4}$

$O \supset C_4$	0_4	1_4	2_4	3_4
$A_1 \downarrow C_4$	1	.	.	.
$A_2 \downarrow C_4$.	.	1	.
$E \downarrow C_4$	1	.	1	.
$T_1 \downarrow C_4$	1	1	.	1
$T_2 \downarrow C_4$.	1	1	1

$O: \chi_g^\mu$	$g=1$	r_{1-4}^p	ρ_{xyz}	R_{xyz}^p	i_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

O characters

C_4 characters

$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$

$$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 e^{\frac{2\pi i m_4 \cdot p}{4}} \mathbf{R}_z^p = \begin{cases} \mathbf{p}_{0_4} = (1 + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{1_4} = (1 + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{2_4} = (1 - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{3_4} = (1 - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4 \end{cases}$$

$$\mathbf{P}^E_{0_40_4} = \mathbf{p}_{0_4} \mathbf{P}^E = \mathbf{P}^E \mathbf{p}_{0_4}$$

$$= \sum_g \frac{\ell^E}{\circ O} (\chi_g^E) \cdot \mathbf{g} \cdot (\mathbf{p}_{0_4}) = \sum_g \frac{2}{96} (\chi_g^E) \cdot \mathbf{g} \cdot (1 \cdot \mathbf{1} + 1 \cdot \rho_z + 1 \cdot \mathbf{R}_z + 1 \cdot \tilde{\mathbf{R}}_z)$$

Calculating \mathbf{P}^E_{0404}

$O \supset C_4$	0_4	1_4	2_4	3_4
$A_1 \downarrow C_4$	1	.	.	.
$A_2 \downarrow C_4$.	.	1	.
$E \downarrow C_4$	1	.	1	.
$T_1 \downarrow C_4$	1	1	.	1
$T_2 \downarrow C_4$.	1	1	1

$O: \chi_g^\mu$	$g=1$	r_{1-4}^p	ρ_{xyz}	R_{xyz}^p	i_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

O characters

C_4 characters

$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$

$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 e^{\frac{2\pi i m_4 \cdot p}{4}} \mathbf{R}_z^p =$

$\mathbf{p}_{0_4} = (1 + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4$

$\mathbf{p}_{1_4} = (1 + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4$

$\mathbf{p}_{2_4} = (1 - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4$

$\mathbf{p}_{3_4} = (1 - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4$

$$\mathbf{P}^E_{0404} = \mathbf{p}_{0_4} \mathbf{P}^E = \mathbf{P}^E \mathbf{p}_{0_4}$$

$$= \sum_g \frac{\ell^E}{\circ O} (\chi_g^E) \cdot \mathbf{g} \cdot (\mathbf{p}_{0_4}) = \sum_g \frac{2}{96} (\chi_g^E) \cdot \mathbf{g} \cdot (1 \cdot \mathbf{1} + 1 \cdot \rho_z + 1 \cdot \mathbf{R}_z + 1 \cdot \tilde{\mathbf{R}}_z)$$

$$1C_4 = \mathbf{1} \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} \quad \rho_x C_4 = \{ \rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3 \} \quad \mathbf{r}_1 C_4 = \{ \mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y \} \quad \mathbf{r}_2 C_4 = \{ \mathbf{r}_2, \mathbf{r}_3, \mathbf{i}_2, \tilde{\mathbf{R}}_y \} \quad \tilde{\mathbf{r}}_1 C_4 = \{ \tilde{\mathbf{r}}_1, \tilde{\mathbf{r}}_3, \tilde{\mathbf{R}}_x, \mathbf{i}_6 \} \quad \tilde{\mathbf{r}}_2 C_4 = \{ \tilde{\mathbf{r}}_2, \tilde{\mathbf{r}}_4, \mathbf{R}_x, \mathbf{i}_5 \}$$

$$= \frac{1}{48} \chi_1^E(1, d_{\rho_z}^{04}, d_{R_z}^{04}, d_{\tilde{R}_z}^{04}) + \frac{1}{48} \chi_{\rho_x}^E(1, d_{\rho_z}^{04}, d_{R_z}^{04}, d_{\tilde{R}_z}^{04}) + \frac{1}{48} \chi_{\mathbf{r}_1}^E(1, d_{\rho_z}^{04}, d_{R_z}^{04}, d_{\tilde{R}_z}^{04}) + \frac{1}{48} \chi_{\mathbf{r}_2}^E(1, d_{\rho_z}^{04}, d_{R_z}^{04}, d_{\tilde{R}_z}^{04}) + \frac{1}{48} \chi_{\tilde{\mathbf{r}}_1}^E(1, d_{\rho_z}^{04}, d_{R_z}^{04}, d_{\tilde{R}_z}^{04}) + \frac{1}{48} \chi_{\tilde{\mathbf{r}}_2}^E(1, d_{\rho_z}^{04}, d_{R_z}^{04}, d_{\tilde{R}_z}^{04})$$

$$+ \frac{1}{48} \chi_{\rho_z}^E(d_{\rho_z}^{04}, 1, d_{\tilde{R}_z}^{04}, d_{R_z}^{04}) + \frac{1}{48} \chi_{\rho_y}^E(d_{\rho_z}^{04}, 1, d_{\tilde{R}_z}^{04}, d_{R_z}^{04}) + \frac{1}{48} \chi_{\mathbf{r}_4}^E(d_{\rho_z}^{04}, 1, d_{\tilde{R}_z}^{04}, d_{R_z}^{04}) + \frac{1}{48} \chi_{\mathbf{r}_3}^E(d_{\rho_z}^{04}, 1, d_{\tilde{R}_z}^{04}, d_{R_z}^{04}) + \frac{1}{48} \chi_{\tilde{\mathbf{r}}_3}^E(d_{\rho_z}^{04}, 1, d_{\tilde{R}_z}^{04}, d_{R_z}^{04}) + \frac{1}{48} \chi_{\tilde{\mathbf{r}}_4}^E(d_{\rho_z}^{04}, 1, d_{\tilde{R}_z}^{04}, d_{R_z}^{04})$$

$$+ \frac{1}{48} \chi_{\mathbf{R}_z}^E(d_{\tilde{R}_z}^{04}, d_{R_z}^{04}, 1, d_{\rho_z}^{04}) + \frac{1}{48} \chi_{\mathbf{i}_4}^E(d_{\tilde{R}_z}^{04}, d_{R_z}^{04}, 1, d_{\rho_z}^{04}) + \frac{1}{48} \chi_{\mathbf{i}_1}^E(d_{\tilde{R}_z}^{04}, d_{R_z}^{04}, 1, d_{\rho_z}^{04}) + \frac{1}{48} \chi_{\mathbf{i}_2}^E(d_{\tilde{R}_z}^{04}, d_{R_z}^{04}, 1, d_{\rho_z}^{04}) + \frac{1}{48} \chi_{\tilde{\mathbf{R}}_x}^E(d_{\tilde{R}_z}^{04}, d_{R_z}^{04}, 1, d_{\rho_z}^{04}) + \frac{1}{48} \chi_{\mathbf{R}_x}^E(d_{\tilde{R}_z}^{04}, d_{R_z}^{04}, 1, d_{\rho_z}^{04})$$

$$+ \frac{1}{48} \chi_{\tilde{\mathbf{R}}_z}^E(d_{R_z}^{04}, d_{\tilde{R}_z}^{04}, d_{\rho_z}^{04}, 1) + \frac{1}{48} \chi_{\mathbf{i}_3}^E(d_{R_z}^{04}, d_{\tilde{R}_z}^{04}, d_{\rho_z}^{04}, 1) + \frac{1}{48} \chi_{\mathbf{R}_y}^E(d_{R_z}^{04}, d_{\tilde{R}_z}^{04}, d_{\rho_z}^{04}, 1) + \frac{1}{48} \chi_{\tilde{\mathbf{R}}_y}^E(d_{R_z}^{04}, d_{\tilde{R}_z}^{04}, d_{\rho_z}^{04}, 1) + \frac{1}{48} \chi_{\mathbf{i}_6}^E(d_{R_z}^{04}, d_{\tilde{R}_z}^{04}, d_{\rho_z}^{04}, 1) + \frac{1}{48} \chi_{\mathbf{i}_5}^E(d_{R_z}^{04}, d_{\tilde{R}_z}^{04}, d_{\rho_z}^{04}, 1)$$

Calculating $\mathbf{P}^E_{0_40_4}$

$O \supset C_4$	0_4	1_4	2_4	3_4
$A_1 \downarrow C_4$	1	.	.	.
$A_2 \downarrow C_4$.	.	1	.
$E \downarrow C_4$	1	.	1	.
$T_1 \downarrow C_4$	1	1	.	1
$T_2 \downarrow C_4$.	1	1	1

$O: \chi_g^\mu$	$g=1$	r_{1-4}^p	ρ_{xyz}	R_{xyz}^p	i_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

O characters

C_4 characters

$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$

$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 e^{\frac{2\pi i m_4 \cdot p}{4}} \mathbf{R}_z^p =$

$\mathbf{p}_{0_4} = (1 + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4$

$\mathbf{p}_{1_4} = (1 + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4$

$\mathbf{p}_{2_4} = (1 - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4$

$\mathbf{p}_{3_4} = (1 - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4$

$\mathbf{P}^E_{0_40_4} = \mathbf{p}_{0_4} \mathbf{P}^E = \mathbf{P}^E \mathbf{p}_{0_4}$

$= \sum_g \frac{\ell^E}{\circ O} (\chi_g^E) \cdot \mathbf{g} \cdot (\mathbf{p}_{0_4}) = \sum_g \frac{2}{96} (\chi_g^E) \cdot \mathbf{g} \cdot (1 \cdot \mathbf{1} + 1 \cdot \rho_z + 1 \cdot \mathbf{R}_z + 1 \cdot \tilde{\mathbf{R}}_z)$

$1C_4 = \mathbf{1} \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \}$ $\rho_x C_4 = \{ \rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3 \}$ $\mathbf{r}_1 C_4 = \{ \mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y \}$ $\mathbf{r}_2 C_4 = \{ \mathbf{r}_2, \mathbf{r}_3, \mathbf{i}_2, \tilde{\mathbf{R}}_y \}$ $\tilde{\mathbf{r}}_1 C_4 = \{ \tilde{\mathbf{r}}_1, \tilde{\mathbf{r}}_3, \tilde{\mathbf{R}}_x, \mathbf{i}_6 \}$ $\tilde{\mathbf{r}}_2 C_4 = \{ \tilde{\mathbf{r}}_2, \tilde{\mathbf{r}}_4, \mathbf{R}_x, \mathbf{i}_5 \}$

$= \frac{1}{48} \chi_1^E(1, d_{\rho_z}^{0_4}, d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4}) + \frac{1}{48} \chi_{\rho_x}^E(1, d_{\rho_z}^{0_4}, d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4}) + \frac{1}{48} \chi_{\mathbf{r}_1}^E(1, d_{\rho_z}^{0_4}, d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4}) + \frac{1}{48} \chi_{\mathbf{r}_2}^E(1, d_{\rho_z}^{0_4}, d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4}) + \frac{1}{48} \chi_{\tilde{\mathbf{r}}_1}^E(1, d_{\rho_z}^{0_4}, d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4}) + \frac{1}{48} \chi_{\tilde{\mathbf{r}}_2}^E(1, d_{\rho_z}^{0_4}, d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4})$

$+ \frac{1}{48} \chi_{\rho_z}^E(d_{\rho_z}^{0_4}, 1, d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4}) + \frac{1}{48} \chi_{\rho_y}^E(d_{\rho_z}^{0_4}, 1, d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4}) + \frac{1}{48} \chi_{\mathbf{r}_4}^E(d_{\rho_z}^{0_4}, 1, d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4}) + \frac{1}{48} \chi_{\mathbf{r}_3}^E(d_{\rho_z}^{0_4}, 1, d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4}) + \frac{1}{48} \chi_{\tilde{\mathbf{r}}_3}^E(d_{\rho_z}^{0_4}, 1, d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4}) + \frac{1}{48} \chi_{\tilde{\mathbf{r}}_4}^E(d_{\rho_z}^{0_4}, 1, d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4})$

$+ \frac{1}{48} \chi_{\mathbf{R}_z}^E(d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4}, 1, d_{\rho_z}^{0_4}) + \frac{1}{48} \chi_{\mathbf{i}_4}^E(d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4}, 1, d_{\rho_z}^{0_4}) + \frac{1}{48} \chi_{\mathbf{i}_1}^E(d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4}, 1, d_{\rho_z}^{0_4}) + \frac{1}{48} \chi_{\mathbf{i}_2}^E(d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4}, 1, d_{\rho_z}^{0_4}) + \frac{1}{48} \chi_{\tilde{\mathbf{R}}_x}^E(d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4}, 1, d_{\rho_z}^{0_4}) + \frac{1}{48} \chi_{\mathbf{R}_x}^E(d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4}, 1, d_{\rho_z}^{0_4})$

$+ \frac{1}{48} \chi_{\tilde{\mathbf{R}}_z}^E(d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4}, d_{\rho_z}^{0_4}, 1) + \frac{1}{48} \chi_{\mathbf{i}_3}^E(d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4}, d_{\rho_z}^{0_4}, 1) + \frac{1}{48} \chi_{\mathbf{R}_y}^E(d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4}, d_{\rho_z}^{0_4}, 1) + \frac{1}{48} \chi_{\tilde{\mathbf{R}}_y}^E(d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4}, d_{\rho_z}^{0_4}, 1) + \frac{1}{48} \chi_{\mathbf{i}_6}^E(d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4}, d_{\rho_z}^{0_4}, 1) + \frac{1}{48} \chi_{\mathbf{i}_5}^E(d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4}, d_{\rho_z}^{0_4}, 1)$

$= \frac{1}{48} (+2)(1, +1, +1, +1) + \frac{1}{48} (+2)(1, +1, +1, +1) + \frac{1}{48} (-1)(1, +1, +1, +1) + \frac{1}{48} (-1)(1, +1, +1, +1) + \frac{1}{48} (-1)(1, +1, +1, +1) + \frac{1}{48} (-1)(1, +1, +1, +1)$

$+ \frac{1}{48} (+2)(+1, 1, +1, +1) + \frac{1}{48} (+2)(+1, 1, +1, +1) + \frac{1}{48} (-1)(+1, 1, +1, +1) + \frac{1}{48} (-1)(+1, 1, +1, +1) + \frac{1}{48} (-1)(+1, 1, +1, +1) + \frac{1}{48} (-1)(+1, 1, +1, +1)$

$+ \frac{1}{48} (0)(+1, +1, 1, +1) + \frac{1}{48} (0)(+1, +1, 1, +1) + \frac{1}{48} (0)(+1, +1, 1, +1) + \frac{1}{48} (0)(+1, +1, 1, +1) + \frac{1}{48} (0)(+1, +1, 1, +1) + \frac{1}{48} (0)(+1, +1, 1, +1)$

$+ \frac{1}{48} (0)(+1, +1, +1, 1) + \frac{1}{48} (0)(+1, +1, +1, 1) + \frac{1}{48} (0)(+1, +1, +1, 1) + \frac{1}{48} (0)(+1, +1, +1, 1) + \frac{1}{48} (0)(+1, +1, +1, 1) + \frac{1}{48} (0)(+1, +1, +1, 1)$

$\underline{\underline{4, 4, 4, 4, \quad 4, 4, 4, 4, \quad -2, -2, -2, -2, \quad -2, -2, -2, -2, \quad -2, -2, -2, -2, \quad -2, -2, -2, -2}}$

Calculating $\mathbf{P}^E_{0_4 0_4}$

$O \supset C_4$	0_4	1_4	2_4	3_4
$A_1 \downarrow C_4$	1	.	.	.
$A_2 \downarrow C_4$.	.	1	.
$E \downarrow C_4$	1	.	1	.
$T_1 \downarrow C_4$	1	1	.	1
$T_2 \downarrow C_4$.	1	1	1

$O: \chi_g^\mu$	$g=1$	r_{1-4}^p	ρ_{xyz}	R_{xyz}^p	i_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

O characters

C_4 characters

$$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$$

$$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 e^{\frac{2\pi i m_4 \cdot p}{4}} \mathbf{R}_z^p =$$

$$\left\{ \begin{aligned} \mathbf{p}_{0_4} &= (1 + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{1_4} &= (1 + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{2_4} &= (1 - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{3_4} &= (1 - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4 \end{aligned} \right.$$

$$\mathbf{P}_{0_4 0_4}^E = \mathbf{p}_{0_4} \mathbf{P}^E = \mathbf{P}^E \mathbf{p}_{0_4}$$

$$= \sum_g \frac{\ell^E}{\circ O} (\chi_g^E) \cdot \mathbf{g} \cdot (\mathbf{p}_{0_4}) = \sum_g \frac{2}{96} (\chi_g^E) \cdot \mathbf{g} \cdot (1 \cdot \mathbf{1} + 1 \cdot \rho_z + 1 \cdot \mathbf{R}_z + 1 \cdot \tilde{\mathbf{R}}_z)$$

$$\mathbf{1}C_4 = \mathbf{1}\{\mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z\} \quad \rho_x C_4 = \{\rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3\} \quad \mathbf{r}_1 C_4 = \{\mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y\} \quad \mathbf{r}_2 C_4 = \{\mathbf{r}_2, \mathbf{r}_3, \mathbf{i}_2, \tilde{\mathbf{R}}_y\} \quad \tilde{\mathbf{r}}_1 C_4 = \{\tilde{\mathbf{r}}_1, \tilde{\mathbf{r}}_3, \tilde{\mathbf{R}}_x, \mathbf{i}_6\} \quad \tilde{\mathbf{r}}_2 C_4 = \{\tilde{\mathbf{r}}_2, \tilde{\mathbf{r}}_4, \mathbf{R}_x, \mathbf{i}_5\}$$

$$= \frac{1}{48} \chi_1^E(1, d_{\rho_z}^{0_4}, d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4}) + \frac{1}{48} \chi_{\rho_x}^E(1, d_{\rho_z}^{0_4}, d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4}) + \frac{1}{48} \chi_{\mathbf{r}_1}^E(1, d_{\rho_z}^{0_4}, d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4}) + \frac{1}{48} \chi_{\mathbf{r}_2}^E(1, d_{\rho_z}^{0_4}, d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4}) + \frac{1}{48} \chi_{\tilde{\mathbf{r}}_1}^E(1, d_{\rho_z}^{0_4}, d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4}) + \frac{1}{48} \chi_{\tilde{\mathbf{r}}_2}^E(1, d_{\rho_z}^{0_4}, d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4})$$

$$+ \frac{1}{48} \chi_{\rho_z}^E(d_{\rho_z}^{0_4}, 1, d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4}) + \frac{1}{48} \chi_{\rho_y}^E(d_{\rho_z}^{0_4}, 1, d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4}) + \frac{1}{48} \chi_{\mathbf{r}_4}^E(d_{\rho_z}^{0_4}, 1, d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4}) + \frac{1}{48} \chi_{\mathbf{r}_3}^E(d_{\rho_z}^{0_4}, 1, d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4}) + \frac{1}{48} \chi_{\tilde{\mathbf{r}}_3}^E(d_{\rho_z}^{0_4}, 1, d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4}) + \frac{1}{48} \chi_{\tilde{\mathbf{r}}_4}^E(d_{\rho_z}^{0_4}, 1, d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4})$$

$$+ \frac{1}{48} \chi_{\mathbf{R}_z}^E(d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4}, 1, d_{\rho_z}^{0_4}) + \frac{1}{48} \chi_{\mathbf{i}_4}^E(d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4}, 1, d_{\rho_z}^{0_4}) + \frac{1}{48} \chi_{\mathbf{i}_1}^E(d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4}, 1, d_{\rho_z}^{0_4}) + \frac{1}{48} \chi_{\mathbf{i}_2}^E(d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4}, 1, d_{\rho_z}^{0_4}) + \frac{1}{48} \chi_{\tilde{\mathbf{R}}_x}^E(d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4}, 1, d_{\rho_z}^{0_4}) + \frac{1}{48} \chi_{\mathbf{R}_x}^E(d_{\tilde{R}_z}^{0_4}, d_{R_z}^{0_4}, 1, d_{\rho_z}^{0_4})$$

$$+ \frac{1}{48} \chi_{\tilde{\mathbf{R}}_z}^E(d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4}, d_{\rho_z}^{0_4}, 1) + \frac{1}{48} \chi_{\mathbf{i}_3}^E(d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4}, d_{\rho_z}^{0_4}, 1) + \frac{1}{48} \chi_{\mathbf{R}_y}^E(d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4}, d_{\rho_z}^{0_4}, 1) + \frac{1}{48} \chi_{\tilde{\mathbf{R}}_y}^E(d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4}, d_{\rho_z}^{0_4}, 1) + \frac{1}{48} \chi_{\mathbf{i}_6}^E(d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4}, d_{\rho_z}^{0_4}, 1) + \frac{1}{48} \chi_{\mathbf{i}_5}^E(d_{R_z}^{0_4}, d_{\tilde{R}_z}^{0_4}, d_{\rho_z}^{0_4}, 1)$$

$$= \frac{1}{48} (+2)(1, +1, +1, +1) + \frac{1}{48} (+2)(1, +1, +1, +1) + \frac{1}{48} (-1)(1, +1, +1, +1) + \frac{1}{48} (-1)(1, +1, +1, +1) + \frac{1}{48} (-1)(1, +1, +1, +1) + \frac{1}{48} (-1)(1, +1, +1, +1)$$

$$+ \frac{1}{48} (+2)(+1, 1, +1, +1) + \frac{1}{48} (+2)(+1, 1, +1, +1) + \frac{1}{48} (-1)(+1, 1, +1, +1) + \frac{1}{48} (-1)(+1, 1, +1, +1) + \frac{1}{48} (-1)(+1, 1, +1, +1) + \frac{1}{48} (-1)(+1, 1, +1, +1)$$

$$+ \frac{1}{48} (0)(+1, +1, 1, +1) + \frac{1}{48} (0)(+1, +1, 1, +1) + \frac{1}{48} (0)(+1, +1, 1, +1) + \frac{1}{48} (0)(+1, +1, 1, +1) + \frac{1}{48} (0)(+1, +1, 1, +1) + \frac{1}{48} (0)(+1, +1, 1, +1)$$

$$+ \frac{1}{48} (0)(+1, +1, +1, 1) + \frac{1}{48} (0)(+1, +1, +1, 1) + \frac{1}{48} (0)(+1, +1, +1, 1) + \frac{1}{48} (0)(+1, +1, +1, 1) + \frac{1}{48} (0)(+1, +1, +1, 1) + \frac{1}{48} (0)(+1, +1, +1, 1)$$

$$\underline{\underline{4, 4, 4, 4, \quad 4, 4, 4, 4, \quad -2, -2, -2, -2, \quad -2, -2, -2, -2, \quad -2, -2, -2, -2, \quad -2, -2, -2, -2}}$$

$$\mathbf{P}_{0_4 0_4}^E = \frac{1}{12} (\underline{\underline{1}} \mathbf{1} + \underline{\underline{1}} \rho_z + \underline{\underline{1}} \mathbf{R}_z + \underline{\underline{1}} \tilde{\mathbf{R}}_z + \underline{\underline{1}} \rho_x + \underline{\underline{1}} \rho_y + \underline{\underline{1}} \mathbf{i}_4 + \underline{\underline{1}} \mathbf{i}_3 \quad \underline{\underline{-\frac{1}{2}}} \mathbf{r}_1 \underline{\underline{-\frac{1}{2}}} \mathbf{r}_4 \underline{\underline{-\frac{1}{2}}} \mathbf{i}_1 \underline{\underline{-\frac{1}{2}}} \mathbf{R}_y \quad \underline{\underline{-\frac{1}{2}}} \mathbf{r}_2 \underline{\underline{-\frac{1}{2}}} \mathbf{r}_3 \underline{\underline{-\frac{1}{2}}} \mathbf{i}_2 \underline{\underline{-\frac{1}{2}}} \tilde{\mathbf{R}}_y \quad \underline{\underline{-\frac{1}{2}}} \tilde{\mathbf{r}}_1 \underline{\underline{-\frac{1}{2}}} \tilde{\mathbf{r}}_3 \underline{\underline{-\frac{1}{2}}} \tilde{\mathbf{R}}_x \underline{\underline{-\frac{1}{2}}} \mathbf{i}_6 \quad \underline{\underline{-\frac{1}{2}}} \tilde{\mathbf{r}}_2 \underline{\underline{-\frac{1}{2}}} \tilde{\mathbf{r}}_4 \underline{\underline{-\frac{1}{2}}} \mathbf{R}_x \underline{\underline{-\frac{1}{2}}} \mathbf{i}_5)$$

Coset-factored sum:

$$\mathbf{P}_{0_4 0_4}^E = \frac{1}{12} [(1) \cdot \mathbf{1} \mathbf{p}_{0_4} + (1) \cdot \rho_x \mathbf{p}_{0_4} + (-\frac{1}{2}) \cdot \mathbf{r}_1 \mathbf{p}_{0_4} + (-\frac{1}{2}) \cdot \mathbf{r}_2 \mathbf{p}_{0_4} + (-\frac{1}{2}) \cdot \tilde{\mathbf{r}}_1 \mathbf{p}_{0_4} + (-\frac{1}{2}) \cdot \tilde{\mathbf{r}}_2 \mathbf{p}_{0_4}]$$

Broken-class-ordered sum:

$$\mathbf{P}_{0_4 0_4}^E = \frac{1}{12} (\mathbf{1} \cdot \mathbf{1} \quad \underline{\underline{-\frac{1}{2}}} \mathbf{r}_1 \underline{\underline{-\frac{1}{2}}} \mathbf{r}_2 \quad \underline{\underline{-\frac{1}{2}}} \mathbf{r}_3 \underline{\underline{-\frac{1}{2}}} \mathbf{r}_4 \quad \underline{\underline{-\frac{1}{2}}} \tilde{\mathbf{r}}_1 \underline{\underline{-\frac{1}{2}}} \tilde{\mathbf{r}}_2 \quad \underline{\underline{-\frac{1}{2}}} \tilde{\mathbf{r}}_3 \underline{\underline{-\frac{1}{2}}} \tilde{\mathbf{r}}_4 \quad + \mathbf{1} \rho_x + \mathbf{1} \rho_y + \mathbf{1} \rho_z \quad \underline{\underline{-\frac{1}{2}}} \mathbf{R}_x \underline{\underline{-\frac{1}{2}}} \mathbf{R}_y \quad + \mathbf{1} \mathbf{R}_z \quad \underline{\underline{-\frac{1}{2}}} \tilde{\mathbf{R}}_x \underline{\underline{-\frac{1}{2}}} \tilde{\mathbf{R}}_y \quad + \mathbf{1} \tilde{\mathbf{R}}_z \quad \underline{\underline{-\frac{1}{2}}} \mathbf{i}_1 \underline{\underline{-\frac{1}{2}}} \mathbf{i}_2 \quad + \mathbf{1} \mathbf{i}_3 + \mathbf{1} \mathbf{i}_4 \quad \underline{\underline{-\frac{1}{2}}} \mathbf{i}_5 \underline{\underline{-\frac{1}{2}}} \mathbf{i}_6)$$

Review Octahedral $O_h \supset O$ group operator structure

Review Octahedral $O_h \supset O \supset D_4 \supset C_4$ subgroup chain correlations

Comparison of $O \supset D_4 \supset C_4$ and $O \supset D_4 \supset D_2$ correlations and level/projector splitting

$O \supset D_4 \supset C_4$ subgroup chain splitting

$O \supset D_4 \supset D_2$ subgroup chain splitting (normal D_2 vs. unnormal D_2)

$O_h \supset O \supset D_4 \supset C_{4v}$ and $O_h \supset O \supset D_4 \supset C_{4v} \supset C_{2v}$ subgroup splitting

Splitting O class projectors P^μ into irreducible projectors $P^\mu_{m_4 m_4}$ for $O \supset C_4$

Development of irreducible projectors $P^\mu_{m_4 m_4}$ and representations $D^\mu_{m_4 m_4}$

Calculating $P^{E_{0404}}$, $P^{E_{2424}}$, $P^{T_{10404}}$, $P^{T_{11414}}$, $P^{T_{22424}}$, $P^{T_{21414}}$,



$O \supset C_4$ induced representation $0_4(C_4) \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples

Elementary induced representation $0_4(C_4) \uparrow O$

Projection reduction of induced representation $0_4(C_4) \uparrow O$

Introduction to ortho-complete eigenvalue-parameter relations

Examples in SF_6 spectroscopy



Calculating \mathbf{P}^E_{2424}

$O \supset C_4$	0_4	1_4	2_4	3_4
$A_1 \downarrow C_4$	1	.	.	.
$A_2 \downarrow C_4$.	.	1	.
$E \downarrow C_4$	1	.	1	.
$T_1 \downarrow C_4$	1	1	.	1
$T_2 \downarrow C_4$.	1	1	1

$O: \chi_g^\mu$	$g=1$	r_{1-4}^p	ρ_{xyz}	R_{xyz}^p	i_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

O characters

C_4 characters

$$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$$

$$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 e^{\frac{2\pi i m_4 \cdot p}{4}} \mathbf{R}_z^p =$$

$$\left\{ \begin{array}{l} \mathbf{p}_{0_4} = (1 + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{1_4} = (1 + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{2_4} = (1 - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{3_4} = (1 - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4 \end{array} \right.$$

$$\mathbf{P}^E_{2424} = \mathbf{p}_{2_4} \mathbf{P}^E = \mathbf{P}^E \mathbf{p}_{2_4}$$

$$= \sum_g \frac{\ell^E}{\circ O} (\chi_g^E) \cdot \mathbf{g} \cdot (\mathbf{p}_{2_4}) = \sum_g \frac{2}{96} (\chi_g^E) \cdot \mathbf{g} \cdot (1 \cdot \mathbf{1} + 1 \cdot \rho_z - 1 \cdot \mathbf{R}_z - 1 \cdot \tilde{\mathbf{R}}_z)$$

$$\mathbf{1}C_4 = \mathbf{1}\{\mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z\} \quad \rho_x C_4 = \{\rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3\} \quad \mathbf{r}_1 C_4 = \{\mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y\} \quad \mathbf{r}_2 C_4 = \{\mathbf{r}_2, \mathbf{r}_3, \mathbf{i}_2, \tilde{\mathbf{R}}_y\} \quad \tilde{\mathbf{r}}_1 C_4 = \{\tilde{\mathbf{r}}_1, \tilde{\mathbf{r}}_3, \tilde{\mathbf{R}}_x, \mathbf{i}_6\} \quad \tilde{\mathbf{r}}_2 C_4 = \{\tilde{\mathbf{r}}_2, \tilde{\mathbf{r}}_4, \mathbf{R}_x, \mathbf{i}_5\}$$

$$= \frac{1}{48} \chi_{\mathbf{1}}^E(1, d_{\rho_z}^{24}, d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24}) + \frac{1}{48} \chi_{\rho_x}^E(1, d_{\rho_z}^{24}, d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24}) + \frac{1}{48} \chi_{\mathbf{r}_1}^E(1, d_{\rho_z}^{24}, d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24}) + \frac{1}{48} \chi_{\mathbf{r}_2}^E(1, d_{\rho_z}^{24}, d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24}) + \frac{1}{48} \chi_{\tilde{\mathbf{r}}_1}^E(1, d_{\rho_z}^{24}, d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24}) + \frac{1}{48} \chi_{\tilde{\mathbf{r}}_2}^E(1, d_{\rho_z}^{24}, d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24})$$

$$+ \frac{1}{48} \chi_{\rho_z}^E(d_{\rho_z}^{24}, 1, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24}) + \frac{1}{48} \chi_{\rho_y}^E(d_{\rho_z}^{24}, 1, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24}) + \frac{1}{48} \chi_{\mathbf{r}_4}^E(d_{\rho_z}^{24}, 1, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24}) + \frac{1}{48} \chi_{\mathbf{r}_3}^E(d_{\rho_z}^{24}, 1, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24}) + \frac{1}{48} \chi_{\tilde{\mathbf{r}}_3}^E(d_{\rho_z}^{24}, 1, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24}) + \frac{1}{48} \chi_{\tilde{\mathbf{r}}_4}^E(d_{\rho_z}^{24}, 1, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24})$$

$$+ \frac{1}{48} \chi_{\mathbf{R}_z}^E(d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24}, 1, d_{\rho_z}^{24}) + \frac{1}{48} \chi_{\mathbf{i}_4}^E(d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24}, 1, d_{\rho_z}^{24}) + \frac{1}{48} \chi_{\mathbf{i}_1}^E(d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24}, 1, d_{\rho_z}^{24}) + \frac{1}{48} \chi_{\mathbf{i}_2}^E(d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24}, 1, d_{\rho_z}^{24}) + \frac{1}{48} \chi_{\tilde{\mathbf{R}}_x}^E(d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24}, 1, d_{\rho_z}^{24}) + \frac{1}{48} \chi_{\mathbf{R}_x}^E(d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24}, 1, d_{\rho_z}^{24})$$

$$+ \frac{1}{48} \chi_{\tilde{\mathbf{R}}_z}^E(d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\rho_z}^{24}, 1) + \frac{1}{48} \chi_{\mathbf{i}_3}^E(d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\rho_z}^{24}, 1) + \frac{1}{48} \chi_{\mathbf{R}_y}^E(d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\rho_z}^{24}, 1) + \frac{1}{48} \chi_{\tilde{\mathbf{R}}_y}^E(d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\rho_z}^{24}, 1) + \frac{1}{48} \chi_{\mathbf{i}_6}^E(d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\rho_z}^{24}, 1) + \frac{1}{48} \chi_{\mathbf{i}_5}^E(d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\rho_z}^{24}, 1)$$

$$= \frac{1}{48} (+2)(1, +1, -1, -1) = \frac{1}{48} (+2)(1, +1, -1, -1) + \frac{1}{48} (-1)(1, +1, -1, -1) + \frac{1}{48} (-1)(1, +1, -1, -1) + \frac{1}{48} (-1)(1, +1, -1, -1) + \frac{1}{48} (-1)(1, +1, -1, -1) + \frac{1}{48} (-1)(1, +1, -1, -1)$$

$$+ \frac{1}{48} (+2)(+1, 1, -1, -1) + \frac{1}{48} (+2)(+1, 1, -1, -1) + \frac{1}{48} (-1)(+1, 1, -1, -1) + \frac{1}{48} (-1)(+1, 1, -1, -1) + \frac{1}{48} (-1)(+1, 1, -1, -1) + \frac{1}{48} (-1)(+1, 1, -1, -1)$$

$$+ \frac{1}{48} (0)(-1, -1, 1, +1) + \frac{1}{48} (0)(-1, -1, 1, +1) + \frac{1}{48} (0)(-1, -1, 1, +1) + \frac{1}{48} (0)(-1, -1, 1, +1) + \frac{1}{48} (0)(-1, -1, 1, +1) + \frac{1}{48} (0)(-1, -1, 1, +1)$$

$$+ \frac{1}{48} (0)(-1, -1, +1, 1) + \frac{1}{48} (0)(-1, -1, +1, 1) + \frac{1}{48} (0)(-1, -1, +1, 1) + \frac{1}{48} (0)(-1, -1, +1, 1) + \frac{1}{48} (0)(-1, -1, +1, 1) + \frac{1}{48} (0)(-1, -1, +1, 1)$$

$$4, 4, -4, -4, \quad 4, 4, -4, -4, \quad -2, -2, 2, 2, \quad -2, -2, 2, 2, \quad -2, -2, 2, 2, \quad -2, -2, 2, 2,$$

$$\frac{1}{12} (\mathbf{1}\mathbf{1} + \mathbf{1}\rho_z - \mathbf{1}\mathbf{R}_z - \mathbf{1}\tilde{\mathbf{R}}_z + \mathbf{1}\rho_x + \mathbf{1}\rho_y - \mathbf{1}\mathbf{i}_4 - \mathbf{1}\mathbf{i}_3 \quad \frac{1}{2}\mathbf{r}_1 \frac{1}{2}\mathbf{r}_4 + \frac{1}{2}\mathbf{i}_1 + \frac{1}{2}\mathbf{R}_y \quad \frac{1}{2}\mathbf{r}_2 \frac{1}{2}\mathbf{r}_3 + \frac{1}{2}\mathbf{i}_2 + \frac{1}{2}\tilde{\mathbf{R}}_y \quad \frac{1}{2}\tilde{\mathbf{r}}_1 \frac{1}{2}\tilde{\mathbf{r}}_3 + \frac{1}{2}\tilde{\mathbf{R}}_x + \frac{1}{2}\mathbf{i}_6 \quad \frac{1}{2}\tilde{\mathbf{r}}_2 \frac{1}{2}\tilde{\mathbf{r}}_4 + \frac{1}{2}\mathbf{R}_x + \frac{1}{2}\mathbf{i}_5)$$

Coset-factored sum:

$$\mathbf{P}^E_{2424} = \frac{1}{12} [(1) \cdot \mathbf{1} \mathbf{p}_{2_4} + (1) \cdot \rho_x \mathbf{p}_{2_4} + (-\frac{1}{2}) \cdot \mathbf{r}_1 \mathbf{p}_{2_4} + (-\frac{1}{2}) \cdot \mathbf{r}_2 \mathbf{p}_{2_4} + (-\frac{1}{2}) \cdot \tilde{\mathbf{r}}_1 \mathbf{p}_{2_4} + (-\frac{1}{2}) \cdot \tilde{\mathbf{r}}_2 \mathbf{p}_{2_4}]$$

Broken-class-ordered sum:

$$\mathbf{P}^E_{2424} = \frac{1}{12} (\mathbf{1}\mathbf{1} - \frac{1}{2}\mathbf{r}_1 - \frac{1}{2}\mathbf{r}_2 - \frac{1}{2}\mathbf{r}_3 - \frac{1}{2}\mathbf{r}_4 - \frac{1}{2}\tilde{\mathbf{r}}_1 - \frac{1}{2}\tilde{\mathbf{r}}_2 - \frac{1}{2}\tilde{\mathbf{r}}_3 - \frac{1}{2}\tilde{\mathbf{r}}_4 + \mathbf{1}\rho_x + \mathbf{1}\rho_y + \mathbf{1}\rho_z + \frac{1}{2}\mathbf{R}_x + \frac{1}{2}\mathbf{R}_y - \mathbf{1}\mathbf{R}_z + \frac{1}{2}\tilde{\mathbf{R}}_x + \frac{1}{2}\tilde{\mathbf{R}}_y - \mathbf{1}\tilde{\mathbf{R}}_z + \frac{1}{2}\mathbf{i}_1 + \frac{1}{2}\mathbf{i}_2 - \mathbf{1}\mathbf{i}_3 - \mathbf{1}\mathbf{i}_4 + \frac{1}{2}\mathbf{i}_5 + \frac{1}{2}\mathbf{i}_6)$$

Review Octahedral $O_h \supset O$ group operator structure

Review Octahedral $O_h \supset O \supset D_4 \supset C_4$ subgroup chain correlations

Comparison of $O \supset D_4 \supset C_4$ and $O \supset D_4 \supset D_2$ correlations and level/projector splitting

$O \supset D_4 \supset C_4$ subgroup chain splitting

$O \supset D_4 \supset D_2$ subgroup chain splitting (normal D_2 vs. unnormal D_2)

$O_h \supset O \supset D_4 \supset C_{4v}$ and $O_h \supset O \supset D_4 \supset C_{4v} \supset C_{2v}$ subgroup splitting

Splitting O class projectors P^μ into irreducible projectors $P^\mu_{m_4 m_4}$ for $O \supset C_4$

Development of irreducible projectors $P^\mu_{m_4 m_4}$ and representations $D^\mu_{m_4 m_4}$

Calculating $P^{E_{0404}}$, $P^{E_{2424}}$, $P^{T_{10404}}$, $P^{T_{11414}}$, $P^{T_{22424}}$, $P^{T_{21414}}$,



$O \supset C_4$ induced representation $0_4(C_4) \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples

Elementary induced representation $0_4(C_4) \uparrow O$

Projection reduction of induced representation $0_4(C_4) \uparrow O$

Introduction to ortho-complete eigenvalue-parameter relations

Examples in SF_6 spectroscopy



Calculating $\mathbf{P}^{T_1}_{0_4 0_4}$

$O \supset C_4$	0_4	1_4	2_4	3_4
$A_1 \downarrow C_4$	1	.	.	.
$A_2 \downarrow C_4$.	.	1	.
$E \downarrow C_4$	1	.	1	.
$T_1 \downarrow C_4$	1	1	.	1
$T_2 \downarrow C_4$.	1	1	1

$O: \chi_g^\mu$	$g=1$	r_{1-4}^p	ρ_{xyz}	R_{xyz}^p	i_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$

C_4 characters

$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 e^{\frac{2\pi i m_4 \cdot p}{4}} \mathbf{R}_z^p =$

$\mathbf{p}_{0_4} = (1 + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4$

$\mathbf{p}_{1_4} = (1 + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4$

$\mathbf{p}_{2_4} = (1 - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4$

$\mathbf{p}_{3_4} = (1 - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4$

$\mathbf{P}_{0_4 0_4}^{T_1} = \mathbf{p}_{0_4} \mathbf{P}^{T_1} = \mathbf{P}^{T_1} \mathbf{p}_{0_4}$

$= \sum_g \frac{\ell^{T_1}}{\ell^O} (\chi_g^{T_1}) \cdot \mathbf{g} \cdot (\mathbf{p}_{0_4}) = \sum_g \frac{3}{96} (\chi_g^{T_1}) \cdot \mathbf{g} \cdot (1 \cdot \mathbf{1} + 1 \cdot \rho_z + 1 \cdot \mathbf{R}_z + 1 \cdot \tilde{\mathbf{R}}_z)$

$\mathbf{1}C_4 = \mathbf{1}\{\mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z\}$ $\rho_x C_4 = \{\rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3\}$ $\mathbf{r}_1 C_4 = \{\mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y\}$ $\mathbf{r}_2 C_4 = \{\mathbf{r}_2, \mathbf{r}_3, \mathbf{i}_2, \tilde{\mathbf{R}}_y\}$ $\tilde{\mathbf{r}}_1 C_4 = \{\tilde{\mathbf{r}}_1, \tilde{\mathbf{r}}_3, \tilde{\mathbf{R}}_x, \mathbf{i}_6\}$ $\tilde{\mathbf{r}}_2 C_4 = \{\tilde{\mathbf{r}}_2, \tilde{\mathbf{r}}_4, \mathbf{R}_x, \mathbf{i}_5\}$

$= \frac{1}{32} \chi_{\mathbf{1}}^{T_1}(1, d_{\rho_z}^{0_4}, d_{\mathbf{R}_z}^{0_4}, d_{\tilde{\mathbf{R}}_z}^{0_4}) + \frac{1}{32} \chi_{\rho_x}^{T_1}(1, d_{\rho_z}^{0_4}, d_{\mathbf{R}_z}^{0_4}, d_{\tilde{\mathbf{R}}_z}^{0_4}) + \frac{1}{32} \chi_{\mathbf{r}_1}^{T_1}(1, d_{\rho_z}^{0_4}, d_{\mathbf{R}_z}^{0_4}, d_{\tilde{\mathbf{R}}_z}^{0_4}) + \frac{1}{32} \chi_{\mathbf{r}_2}^{T_1}(1, d_{\rho_z}^{0_4}, d_{\mathbf{R}_z}^{0_4}, d_{\tilde{\mathbf{R}}_z}^{0_4}) + \frac{1}{32} \chi_{\tilde{\mathbf{r}}_1}^{T_1}(1, d_{\rho_z}^{0_4}, d_{\mathbf{R}_z}^{0_4}, d_{\tilde{\mathbf{R}}_z}^{0_4}) + \frac{1}{32} \chi_{\tilde{\mathbf{r}}_2}^{T_1}(1, d_{\rho_z}^{0_4}, d_{\mathbf{R}_z}^{0_4}, d_{\tilde{\mathbf{R}}_z}^{0_4})$

$+ \frac{1}{32} \chi_{\rho_z}^{T_1}(d_{\rho_z}^{0_4}, 1, d_{\tilde{\mathbf{R}}_z}^{0_4}, d_{\mathbf{R}_z}^{0_4}) + \frac{1}{32} \chi_{\rho_y}^{T_1}(d_{\rho_z}^{0_4}, 1, d_{\tilde{\mathbf{R}}_z}^{0_4}, d_{\mathbf{R}_z}^{0_4}) + \frac{1}{32} \chi_{\mathbf{r}_4}^{T_1}(d_{\rho_z}^{0_4}, 1, d_{\tilde{\mathbf{R}}_z}^{0_4}, d_{\mathbf{R}_z}^{0_4}) + \frac{1}{32} \chi_{\mathbf{r}_3}^{T_1}(d_{\rho_z}^{0_4}, 1, d_{\tilde{\mathbf{R}}_z}^{0_4}, d_{\mathbf{R}_z}^{0_4}) + \frac{1}{32} \chi_{\tilde{\mathbf{r}}_3}^{T_1}(d_{\rho_z}^{0_4}, 1, d_{\tilde{\mathbf{R}}_z}^{0_4}, d_{\mathbf{R}_z}^{0_4}) + \frac{1}{32} \chi_{\tilde{\mathbf{r}}_4}^{T_1}(d_{\rho_z}^{0_4}, 1, d_{\tilde{\mathbf{R}}_z}^{0_4}, d_{\mathbf{R}_z}^{0_4})$

$+ \frac{1}{32} \chi_{\mathbf{R}_z}^{T_1}(d_{\tilde{\mathbf{R}}_z}^{0_4}, d_{\mathbf{R}_z}^{0_4}, 1, d_{\rho_z}^{0_4}) + \frac{1}{32} \chi_{\mathbf{i}_4}^{T_1}(d_{\tilde{\mathbf{R}}_z}^{0_4}, d_{\mathbf{R}_z}^{0_4}, 1, d_{\rho_z}^{0_4}) + \frac{1}{32} \chi_{\mathbf{i}_1}^{T_1}(d_{\tilde{\mathbf{R}}_z}^{0_4}, d_{\mathbf{R}_z}^{0_4}, 1, d_{\rho_z}^{0_4}) + \frac{1}{32} \chi_{\mathbf{i}_2}^{T_1}(d_{\tilde{\mathbf{R}}_z}^{0_4}, d_{\mathbf{R}_z}^{0_4}, 1, d_{\rho_z}^{0_4}) + \frac{1}{32} \chi_{\tilde{\mathbf{R}}_x}^{T_1}(d_{\tilde{\mathbf{R}}_z}^{0_4}, d_{\mathbf{R}_z}^{0_4}, 1, d_{\rho_z}^{0_4}) + \frac{1}{32} \chi_{\mathbf{R}_x}^{T_1}(d_{\tilde{\mathbf{R}}_z}^{0_4}, d_{\mathbf{R}_z}^{0_4}, 1, d_{\rho_z}^{0_4})$

$+ \frac{1}{32} \chi_{\tilde{\mathbf{R}}_z}^{T_1}(d_{\mathbf{R}_z}^{0_4}, d_{\tilde{\mathbf{R}}_z}^{0_4}, d_{\rho_z}^{0_4}, 1) + \frac{1}{32} \chi_{\mathbf{i}_3}^{T_1}(d_{\mathbf{R}_z}^{0_4}, d_{\tilde{\mathbf{R}}_z}^{0_4}, d_{\rho_z}^{0_4}, 1) + \frac{1}{32} \chi_{\mathbf{R}_y}^{T_1}(d_{\mathbf{R}_z}^{0_4}, d_{\tilde{\mathbf{R}}_z}^{0_4}, d_{\rho_z}^{0_4}, 1) + \frac{1}{32} \chi_{\tilde{\mathbf{R}}_y}^{T_1}(d_{\mathbf{R}_z}^{0_4}, d_{\tilde{\mathbf{R}}_z}^{0_4}, d_{\rho_z}^{0_4}, 1) + \frac{1}{32} \chi_{\mathbf{i}_6}^{T_1}(d_{\mathbf{R}_z}^{0_4}, d_{\tilde{\mathbf{R}}_z}^{0_4}, d_{\rho_z}^{0_4}, 1) + \frac{1}{32} \chi_{\mathbf{i}_5}^{T_1}(d_{\mathbf{R}_z}^{0_4}, d_{\tilde{\mathbf{R}}_z}^{0_4}, d_{\rho_z}^{0_4}, 1)$

$= \frac{1}{32} (+3)(1, +1, +1, +1) + \frac{1}{32} (-1)(1, +1, +1, +1) + \frac{1}{32} (0)(1, +1, +1, +1) + \frac{1}{32} (0)(1, +1, +1, +1) + \frac{1}{32} (0)(1, +1, +1, +1) + \frac{1}{32} (0)(1, +1, +1, +1)$

$+ \frac{1}{32} (-1)(+1, 1, +1, +1) + \frac{1}{32} (-1)(+1, 1, +1, +1) + \frac{1}{32} (0)(+1, 1, +1, +1) + \frac{1}{32} (0)(+1, 1, +1, +1) + \frac{1}{32} (0)(+1, 1, +1, +1) + \frac{1}{32} (0)(+1, 1, +1, +1)$

$+ \frac{1}{32} (+1)(+1, +1, 1, +1) + \frac{1}{32} (-1)(+1, +1, 1, +1) + \frac{1}{32} (-1)(+1, +1, 1, +1) + \frac{1}{32} (-1)(+1, +1, 1, +1) + \frac{1}{32} (+1)(+1, +1, 1, +1) + \frac{1}{32} (+1)(+1, +1, 1, +1)$

$+ \frac{1}{32} (+1)(+1, +1, +1, 1) + \frac{1}{32} (-1)(+1, +1, +1, 1) + \frac{1}{32} (+1)(+1, +1, +1, 1) + \frac{1}{32} (+1)(+1, +1, +1, 1) + \frac{1}{32} (-1)(+1, +1, +1, 1) + \frac{1}{32} (-1)(+1, +1, +1, 1)$

$\underline{4, 4, 0, 0}, \quad \underline{-4, -4, -4, -4}, \quad \underline{0, 0, 0, 0}, \quad \underline{0, 0, 0, 0}, \quad \underline{0, 0, 0, 0}, \quad \underline{0, 0, 0, 0}$

$\frac{1}{8} (\underline{11+1\rho_z+1\mathbf{R}_z+1\tilde{\mathbf{R}}_z} \quad \underline{-1\rho_x-1\rho_y-1\mathbf{i}_4-1\mathbf{i}_3} \quad + \underline{0\mathbf{r}_1+0\mathbf{r}_4+0\mathbf{i}_1+0\mathbf{R}_y} \quad + \underline{0\mathbf{r}_2+0\mathbf{r}_3+0\mathbf{i}_2+0\tilde{\mathbf{R}}_y} \quad + \underline{0\tilde{\mathbf{r}}_1+0\tilde{\mathbf{r}}_3+0\tilde{\mathbf{R}}_x+0\mathbf{i}_6} \quad + \underline{0\tilde{\mathbf{r}}_2+0\tilde{\mathbf{r}}_4+0\mathbf{R}_x+0\mathbf{i}_5})$

Coset-factored sum:

$\mathbf{P}_{0_4 0_4}^{T_1} = \frac{1}{8} [(1) \cdot \mathbf{1p}_{0_4} + (-1) \cdot \rho_x \mathbf{p}_{0_4} + (0) \cdot \mathbf{r}_1 \mathbf{p}_{0_4} + (0) \cdot \mathbf{r}_2 \mathbf{p}_{0_4} + (0) \cdot \tilde{\mathbf{r}}_1 \mathbf{p}_{0_4} + (0) \cdot \tilde{\mathbf{r}}_2 \mathbf{p}_{0_4}]$

Broken-class-ordered sum:

$\mathbf{P}_{0_4 0_4}^{T_1} = \frac{1}{8} (1 \cdot \mathbf{1} + 0 + 0 + 0 + 0 + 0 + 0 + 0 + 0 + 1\rho_z - 1\rho_x - 1\rho_y + 0 + 0 + 1\mathbf{R}_z + 0 + 0 + 1\tilde{\mathbf{R}}_z + 0 + 0 + 0 + 0 - 1\mathbf{i}_4 - 1\mathbf{i}_3)$

Review Octahedral $O_h \supset O$ group operator structure

Review Octahedral $O_h \supset O \supset D_4 \supset C_4$ subgroup chain correlations

Comparison of $O \supset D_4 \supset C_4$ and $O \supset D_4 \supset D_2$ correlations and level/projector splitting

$O \supset D_4 \supset C_4$ subgroup chain splitting

$O \supset D_4 \supset D_2$ subgroup chain splitting (normal D_2 vs. unnormal D_2)

$O_h \supset O \supset D_4 \supset C_{4v}$ and $O_h \supset O \supset D_4 \supset C_{4v} \supset C_{2v}$ subgroup splitting

Splitting O class projectors P^μ into irreducible projectors $P^\mu_{m_4 m_4}$ for $O \supset C_4$

Development of irreducible projectors $P^\mu_{m_4 m_4}$ and representations $D^\mu_{m_4 m_4}$

Calculating $P^{E_{0404}}$, $P^{E_{2424}}$, $P^{T_{10404}}$, $P^{T_{11414}}$, $P^{T_{22424}}$, $P^{T_{21414}}$,



$O \supset C_4$ induced representation $0_4(C_4) \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples

Elementary induced representation $0_4(C_4) \uparrow O$

Projection reduction of induced representation $0_4(C_4) \uparrow O$

Introduction to ortho-complete eigenvalue-parameter relations

Examples in SF_6 spectroscopy



Calculating $\mathbf{P}_{1_4 1_4}^{T_1}$

$O \supset C_4$	0_4	1_4	2_4	3_4
$A_1 \downarrow C_4$	1	.	.	.
$A_2 \downarrow C_4$.	.	1	.
$E \downarrow C_4$	1	.	1	.
$T_1 \downarrow C_4$	1	1	.	1
$T_2 \downarrow C_4$.	1	1	1

$O: \chi_g^\mu$	$g=1$	\mathbf{r}_{1-4}^p	ρ_{xyz}	\mathbf{R}_{xyz}^p	\mathbf{i}_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

O characters

C_4 characters

$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$

$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 e^{\frac{2\pi i m_4 \cdot p}{4}} \mathbf{R}_z^p =$

$\mathbf{p}_{0_4} = (1 + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4$

$\mathbf{p}_{1_4} = (1 + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4$

$\mathbf{p}_{2_4} = (1 - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4$

$\mathbf{p}_{3_4} = (1 - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4$

$$\mathbf{P}_{1_4 1_4}^{T_1} = \mathbf{p}_{1_4} \mathbf{P}^{T_1} = \mathbf{P}^{T_1} \mathbf{p}_{1_4}$$

$$= \sum_g \frac{\ell^{T_1}}{\circ O} (\chi_g^{T_1}) \cdot \mathbf{g} \cdot (\mathbf{p}_{1_4}) = \sum_g \frac{3}{96} (\chi_g^{T_1}) \cdot \mathbf{g} \cdot (1 \cdot \mathbf{1} - 1 \cdot \rho_z + i \cdot \mathbf{R}_z - i \cdot \tilde{\mathbf{R}}_z)$$

$$1C_4 = \mathbf{1} \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} \quad \rho_x C_4 = \{ \rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3 \} \quad \mathbf{r}_1 C_4 = \{ \mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y \} \quad \mathbf{r}_2 C_4 = \{ \mathbf{r}_2, \mathbf{r}_3, \mathbf{i}_2, \tilde{\mathbf{R}}_y \} \quad \tilde{\mathbf{r}}_1 C_4 = \{ \tilde{\mathbf{r}}_1, \tilde{\mathbf{r}}_3, \tilde{\mathbf{R}}_x, \mathbf{i}_6 \} \quad \tilde{\mathbf{r}}_2 C_4 = \{ \tilde{\mathbf{r}}_2, \tilde{\mathbf{r}}_4, \mathbf{R}_x, \mathbf{i}_5 \}$$

$$= \frac{1}{32} \chi_{\mathbf{1}}^{T_1}(1, d_{\rho_z}^{14}, d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14}) + \frac{1}{32} \chi_{\rho_x}^{T_1}(1, d_{\rho_z}^{14}, d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14}) + \frac{1}{32} \chi_{\mathbf{r}_1}^{T_1}(1, d_{\rho_z}^{14}, d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14}) + \frac{1}{32} \chi_{\mathbf{r}_2}^{T_1}(1, d_{\rho_z}^{14}, d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14}) + \frac{1}{32} \chi_{\tilde{\mathbf{r}}_1}^{T_1}(1, d_{\rho_z}^{14}, d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14}) + \frac{1}{32} \chi_{\tilde{\mathbf{r}}_2}^{T_1}(1, d_{\rho_z}^{14}, d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14})$$

$$+ \frac{1}{32} \chi_{\rho_z}^{T_1}(d_{\rho_z}^{14}, 1, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14}) + \frac{1}{32} \chi_{\rho_y}^{T_1}(d_{\rho_z}^{14}, 1, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14}) + \frac{1}{32} \chi_{\mathbf{r}_4}^{T_1}(d_{\rho_z}^{14}, 1, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14}) + \frac{1}{32} \chi_{\mathbf{r}_3}^{T_1}(d_{\rho_z}^{14}, 1, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14}) + \frac{1}{32} \chi_{\tilde{\mathbf{r}}_3}^{T_1}(d_{\rho_z}^{14}, 1, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14}) + \frac{1}{32} \chi_{\tilde{\mathbf{r}}_4}^{T_1}(d_{\rho_z}^{14}, 1, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14})$$

$$+ \frac{1}{32} \chi_{\mathbf{R}_z}^{T_1}(d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14}, 1, d_{\rho_z}^{14}) + \frac{1}{32} \chi_{\mathbf{i}_4}^{T_1}(d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14}, 1, d_{\rho_z}^{14}) + \frac{1}{32} \chi_{\mathbf{i}_1}^{T_1}(d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14}, 1, d_{\rho_z}^{14}) + \frac{1}{32} \chi_{\mathbf{i}_2}^{T_1}(d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14}, 1, d_{\rho_z}^{14}) + \frac{1}{32} \chi_{\tilde{\mathbf{R}}_x}^{T_1}(d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14}, 1, d_{\rho_z}^{14}) + \frac{1}{32} \chi_{\mathbf{R}_x}^{T_1}(d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14}, 1, d_{\rho_z}^{14})$$

$$+ \frac{1}{32} \chi_{\tilde{\mathbf{R}}_z}^{T_1}(d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\rho_z}^{14}, 1) + \frac{1}{32} \chi_{\mathbf{i}_3}^{T_1}(d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\rho_z}^{14}, 1) + \frac{1}{32} \chi_{\mathbf{R}_y}^{T_1}(d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\rho_z}^{14}, 1) + \frac{1}{32} \chi_{\tilde{\mathbf{R}}_y}^{T_1}(d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\rho_z}^{14}, 1) + \frac{1}{32} \chi_{\mathbf{i}_6}^{T_1}(d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\rho_z}^{14}, 1) + \frac{1}{32} \chi_{\mathbf{i}_5}^{T_1}(d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\rho_z}^{14}, 1)$$

$$= \frac{1}{32} (+3)(1, -1, +i, -i) + \frac{1}{32} (-1)(1, -1, +i, -i) + \frac{1}{32} (0)(1, -1, +i, -i) + \frac{1}{32} (0)(1, -1, +i, -i) + \frac{1}{32} (0)(1, -1, +i, -i) + \frac{1}{32} (0)(1, -1, +i, -i)$$

$$+ \frac{1}{32} (-1)(-1, 1, -i, +i) + \frac{1}{32} (-1)(-1, 1, -i, +i) + \frac{1}{32} (0)(-1, 1, -i, +i) + \frac{1}{32} (0)(-1, 1, -i, +i) + \frac{1}{32} (0)(-1, 1, -i, +i) + \frac{1}{32} (0)(-1, 1, -i, +i)$$

$$+ \frac{1}{32} (+1)(-i, +i, 1, -1) + \frac{1}{32} (-1)(-i, +i, 1, -1) + \frac{1}{32} (-1)(-i, +i, 1, -1) + \frac{1}{32} (-1)(-i, +i, 1, -1) + \frac{1}{32} (+1)(-i, +i, 1, -1) + \frac{1}{32} (+1)(-i, +i, 1, -1)$$

$$+ \frac{1}{32} (+1)(+i, -i, -1, 1) + \frac{1}{32} (-1)(+i, -i, -1, 1) + \frac{1}{32} (+1)(+i, -i, -1, 1) + \frac{1}{32} (+1)(+i, -i, -1, 1) + \frac{1}{32} (-1)(+i, -i, -1, 1) + \frac{1}{32} (-1)(+i, -i, -1, 1)$$

$$+ 4, -4, 4i, -4i, \quad 0, \quad 0, \quad 0, \quad 0, \quad +2i, -2i, -2, +2, \quad +2i, -2i, -2, +2, \quad -2i, +2i, +2, -2, \quad -2i, +2i, +2, -2.$$

$$\frac{1}{8} (\underline{11} - \underline{1}\rho_z + \underline{i}\mathbf{R}_z - \underline{i}\tilde{\mathbf{R}}_z + \underline{0}\rho_x + \underline{0}\rho_y + \underline{0}\mathbf{i}_4 + \underline{0}\mathbf{i}_3 + \frac{i}{2}\mathbf{r}_1 - \frac{i}{2}\mathbf{r}_4 - \frac{1}{2}\mathbf{i}_1 + \frac{1}{2}\mathbf{R}_y + \frac{i}{2}\mathbf{r}_2 - \frac{i}{2}\mathbf{r}_3 - \frac{1}{2}\mathbf{i}_2 + \frac{1}{2}\tilde{\mathbf{R}}_y - \frac{i}{2}\tilde{\mathbf{r}}_1 + \frac{i}{2}\tilde{\mathbf{r}}_3 + \frac{1}{2}\tilde{\mathbf{R}}_x - \frac{1}{2}\mathbf{i}_6 - \frac{i}{2}\tilde{\mathbf{r}}_2 + \frac{i}{2}\tilde{\mathbf{r}}_4 + \frac{1}{2}\mathbf{R}_x - \frac{1}{2}\mathbf{i}_5)$$

Coset-factored sum:

$$\mathbf{P}_{1_4 1_4}^{T_1} = \frac{1}{8} [(1) \cdot \mathbf{1} \mathbf{p}_{1_4} + (0) \cdot \rho_x \mathbf{p}_{1_4} + (\frac{i}{2}) \cdot \mathbf{r}_1 \mathbf{p}_{1_4} + (\frac{i}{2}) \cdot \mathbf{r}_2 \mathbf{p}_{1_4} + (-\frac{i}{2}) \cdot \tilde{\mathbf{r}}_1 \mathbf{p}_{1_4} + (-\frac{i}{2}) \cdot \tilde{\mathbf{r}}_2 \mathbf{p}_{1_4}]$$

Broken-class-ordered sum:

$$\mathbf{P}_{1_4 1_4}^{T_1} = \frac{1}{8} (\mathbf{1} \cdot \mathbf{1} + \frac{i}{2}\mathbf{r}_1 + \frac{i}{2}\mathbf{r}_2 - \frac{i}{2}\mathbf{r}_3 - \frac{i}{2}\mathbf{r}_4 - \frac{i}{2}\tilde{\mathbf{r}}_1 - \frac{i}{2}\tilde{\mathbf{r}}_2 + \frac{i}{2}\tilde{\mathbf{r}}_3 + \frac{i}{2}\tilde{\mathbf{r}}_4 + \mathbf{0}\rho_x + \mathbf{0}\rho_y - \mathbf{1}\rho_z + \frac{1}{2}\mathbf{R}_x + \frac{1}{2}\mathbf{R}_y + \mathbf{i}\mathbf{R}_z + \frac{1}{2}\tilde{\mathbf{R}}_x + \frac{1}{2}\tilde{\mathbf{R}}_y - \mathbf{i}\tilde{\mathbf{R}}_z - \frac{i}{2}\mathbf{i}_1 - \frac{i}{2}\mathbf{i}_2 + \mathbf{0}\mathbf{i}_3 + \mathbf{0}\mathbf{i}_4 - \frac{i}{2}\mathbf{i}_5 - \frac{i}{2}\mathbf{i}_6)$$

Review Octahedral $O_h \supset O$ group operator structure

Review Octahedral $O_h \supset O \supset D_4 \supset C_4$ subgroup chain correlations

Comparison of $O \supset D_4 \supset C_4$ and $O \supset D_4 \supset D_2$ correlations and level/projector splitting

$O \supset D_4 \supset C_4$ subgroup chain splitting

$O \supset D_4 \supset D_2$ subgroup chain splitting (normal D_2 vs. unnormal D_2)

$O_h \supset O \supset D_4 \supset C_{4v}$ and $O_h \supset O \supset D_4 \supset C_{4v} \supset C_{2v}$ subgroup splitting

Splitting O class projectors P^μ into irreducible projectors $P^\mu_{m_4m_4}$ for $O \supset C_4$

Development of irreducible projectors $P^\mu_{m_4m_4}$ and representations $D^\mu_{m_4m_4}$

Calculating $P^{E_{0404}}$, $P^{E_{2424}}$, $P^{T_{10404}}$, $P^{T_{11414}}$, $P^{T_{22424}}$, $P^{T_{21414}}$,



$O \supset C_4$ induced representation $0_4(C_4) \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples

Elementary induced representation $0_4(C_4) \uparrow O$

Projection reduction of induced representation $0_4(C_4) \uparrow O$

Introduction to ortho-complete eigenvalue-parameter relations

Examples in SF_6 spectroscopy



Calculating $\mathbf{P}^{T_2}_{2_4 2_4}$

$O \supset C_4$	0_4	1_4	2_4	3_4
$A_1 \downarrow C_4$	1	.	.	.
$A_2 \downarrow C_4$.	.	1	.
$E \downarrow C_4$	1	.	1	.
$T_1 \downarrow C_4$	1	1	.	1
$T_2 \downarrow C_4$.	1	1	1

$O: \chi_g^\mu$	$g=1$	r_{1-4}^p	ρ_{xyz}	R_{xyz}^p	i_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

O characters

C_4 characters

$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$

$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 e^{\frac{2\pi i m_4 \cdot p}{4}} \mathbf{R}_z^p =$

$$\left\{ \begin{aligned} \mathbf{p}_{0_4} &= (\mathbf{1} + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z) / 4 \\ \mathbf{p}_{1_4} &= (\mathbf{1} + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z) / 4 \\ \mathbf{p}_{2_4} &= (\mathbf{1} - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z) / 4 \\ \mathbf{p}_{3_4} &= (\mathbf{1} - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z) / 4 \end{aligned} \right.$$

$$\mathbf{P}_{2_4 2_4}^{T_2} = \mathbf{p}_{2_4} \mathbf{P}^{T_2} = \mathbf{P}^{T_2} \mathbf{p}_{2_4}$$

$$= \sum_g \frac{\ell^{T_2}}{\circ O} (\chi_g^{T_2}) \cdot \mathbf{g} \cdot (\mathbf{p}_{2_4}) = \sum_g \frac{3}{96} (\chi_g^{T_2}) \cdot \mathbf{g} \cdot (\mathbf{1} \cdot \mathbf{1} + \mathbf{1} \cdot \rho_z - \mathbf{1} \cdot \mathbf{R}_z - \mathbf{1} \cdot \tilde{\mathbf{R}}_z)$$

$$\mathbf{1}C_4 = \mathbf{1}\{\mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z\} \quad \rho_x C_4 = \{\rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3\} \quad \mathbf{r}_1 C_4 = \{\mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y\} \quad \mathbf{r}_2 C_4 = \{\mathbf{r}_2, \mathbf{r}_3, \mathbf{i}_2, \tilde{\mathbf{R}}_y\} \quad \tilde{\mathbf{r}}_1 C_4 = \{\tilde{\mathbf{r}}_1, \tilde{\mathbf{r}}_3, \tilde{\mathbf{R}}_x, \mathbf{i}_6\} \quad \tilde{\mathbf{r}}_2 C_4 = \{\tilde{\mathbf{r}}_2, \tilde{\mathbf{r}}_4, \mathbf{R}_x, \mathbf{i}_5\}$$

$$= \frac{1}{32} \chi_{\mathbf{1}}^{T_2}(1, d_{\rho_z}^{24}, d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24}) + \frac{1}{32} \chi_{\rho_x}^{T_2}(1, d_{\rho_z}^{24}, d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24}) + \frac{1}{32} \chi_{\mathbf{r}_1}^{T_2}(1, d_{\rho_z}^{24}, d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24}) + \frac{1}{32} \chi_{\mathbf{r}_2}^{T_2}(1, d_{\rho_z}^{24}, d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24}) + \frac{1}{32} \chi_{\tilde{\mathbf{r}}_1}^{T_2}(1, d_{\rho_z}^{24}, d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24}) + \frac{1}{32} \chi_{\tilde{\mathbf{r}}_2}^{T_2}(1, d_{\rho_z}^{24}, d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24})$$

$$+ \frac{1}{32} \chi_{\rho_z}^{T_2}(d_{\rho_z}^{24}, 1, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24}) + \frac{1}{32} \chi_{\rho_y}^{T_2}(d_{\rho_z}^{24}, 1, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24}) + \frac{1}{32} \chi_{\mathbf{r}_4}^{T_2}(d_{\rho_z}^{24}, 1, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24}) + \frac{1}{32} \chi_{\mathbf{r}_3}^{T_2}(d_{\rho_z}^{24}, 1, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24}) + \frac{1}{32} \chi_{\tilde{\mathbf{r}}_3}^{T_2}(d_{\rho_z}^{24}, 1, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24}) + \frac{1}{32} \chi_{\tilde{\mathbf{r}}_4}^{T_2}(d_{\rho_z}^{24}, 1, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24})$$

$$+ \frac{1}{32} \chi_{\mathbf{R}_z}^{T_2}(d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24}, 1, d_{\rho_z}^{24}) + \frac{1}{32} \chi_{\mathbf{i}_4}^{T_2}(d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24}, 1, d_{\rho_z}^{24}) + \frac{1}{32} \chi_{\mathbf{i}_1}^{T_2}(d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24}, 1, d_{\rho_z}^{24}) + \frac{1}{32} \chi_{\mathbf{i}_2}^{T_2}(d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24}, 1, d_{\rho_z}^{24}) + \frac{1}{32} \chi_{\tilde{\mathbf{R}}_x}^{T_2}(d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24}, 1, d_{\rho_z}^{24}) + \frac{1}{32} \chi_{\mathbf{R}_x}^{T_2}(d_{\tilde{\mathbf{R}}_z}^{24}, d_{\mathbf{R}_z}^{24}, 1, d_{\rho_z}^{24})$$

$$+ \frac{1}{32} \chi_{\tilde{\mathbf{R}}_z}^{T_2}(d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\rho_z}^{24}, 1) + \frac{1}{32} \chi_{\mathbf{i}_3}^{T_2}(d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\rho_z}^{24}, 1) + \frac{1}{32} \chi_{\mathbf{R}_y}^{T_2}(d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\rho_z}^{24}, 1) + \frac{1}{32} \chi_{\tilde{\mathbf{R}}_y}^{T_2}(d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\rho_z}^{24}, 1) + \frac{1}{32} \chi_{\mathbf{i}_6}^{T_2}(d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\rho_z}^{24}, 1) + \frac{1}{32} \chi_{\mathbf{i}_5}^{T_2}(d_{\mathbf{R}_z}^{24}, d_{\tilde{\mathbf{R}}_z}^{24}, d_{\rho_z}^{24}, 1)$$

$$= \frac{1}{32} (+3)(1, +1, -1, -1) + \frac{1}{32} (-1)(1, +1, -1, -1) + \frac{1}{32} (0)(1, +1, -1, -1) + \frac{1}{32} (0)(1, +1, -1, -1) + \frac{1}{32} (0)(1, +1, -1, -1) + \frac{1}{32} (0)(1, +1, -1, -1)$$

$$+ \frac{1}{32} (-1)(+1, 1, -1, -1) + \frac{1}{32} (-1)(+1, 1, -1, -1) + \frac{1}{32} (0)(+1, 1, -1, -1) + \frac{1}{32} (0)(+1, 1, -1, -1) + \frac{1}{32} (0)(+1, 1, -1, -1) + \frac{1}{32} (0)(+1, 1, -1, -1)$$

$$+ \frac{1}{32} (-1)(-1, -1, 1, +1) + \frac{1}{32} (+1)(-1, -1, 1, +1) + \frac{1}{32} (+1)(-1, -1, 1, +1) + \frac{1}{32} (+1)(-1, -1, 1, +1) + \frac{1}{32} (-1)(-1, -1, 1, +1) + \frac{1}{32} (-1)(-1, -1, 1, +1)$$

$$+ \frac{1}{32} (-1)(-1, -1, +1, 1) + \frac{1}{32} (+1)(-1, -1, +1, 1) + \frac{1}{32} (-1)(-1, -1, +1, 1) + \frac{1}{32} (-1)(-1, -1, +1, 1) + \frac{1}{32} (+1)(-1, -1, +1, 1) + \frac{1}{32} (+1)(-1, -1, +1, 1)$$

$$\underline{\underline{4, 4, -4, -4, -4, -4, 4, 4, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0}}$$

$$\frac{1}{8} (\mathbf{1}\mathbf{1} + \mathbf{1}\rho_z - \mathbf{1}\mathbf{R}_z - \mathbf{1}\tilde{\mathbf{R}}_z - \mathbf{1}\rho_x - \mathbf{1}\rho_y + \mathbf{1}\mathbf{i}_4 + \mathbf{1}\mathbf{i}_3 + \mathbf{0}\mathbf{r}_1 + \mathbf{0}\mathbf{r}_4 + \mathbf{0}\mathbf{i}_1 + \mathbf{0}\mathbf{R}_y + \mathbf{0}\mathbf{r}_2 + \mathbf{0}\mathbf{r}_3 + \mathbf{0}\mathbf{i}_2 + \mathbf{0}\tilde{\mathbf{R}}_y + \mathbf{0}\tilde{\mathbf{r}}_1 + \mathbf{0}\tilde{\mathbf{r}}_3 + \mathbf{0}\tilde{\mathbf{R}}_x + \mathbf{0}\mathbf{i}_6 + \mathbf{0}\tilde{\mathbf{r}}_2 + \mathbf{0}\tilde{\mathbf{r}}_4 + \mathbf{0}\mathbf{R}_x + \mathbf{0}\mathbf{i}_5)$$

Coset-factored sum:

$$\mathbf{P}_{2_4 2_4}^{T_2} = \frac{1}{8} [(\mathbf{1}) \cdot \mathbf{1}\mathbf{p}_{2_4} + (\mathbf{1}) \cdot \rho_x \mathbf{p}_{2_4} + (\mathbf{0}) \cdot \mathbf{r}_1 \mathbf{p}_{2_4} + (\mathbf{0}) \cdot \mathbf{r}_2 \mathbf{p}_{2_4} + (\mathbf{0}) \cdot \tilde{\mathbf{r}}_1 \mathbf{p}_{2_4} + (\mathbf{0}) \cdot \tilde{\mathbf{r}}_2 \mathbf{p}_{2_4}]$$

Broken-class-ordered sum:

$$\mathbf{P}_{2_4 2_4}^{T_2} = \frac{1}{8} (\mathbf{1} \cdot \mathbf{1} + \mathbf{0} + \mathbf{0} + \mathbf{0} + \mathbf{0} + \mathbf{0} + \mathbf{0} + \mathbf{0} + \mathbf{0} + \mathbf{0} - \mathbf{1}\rho_x - \mathbf{1}\rho_y + \mathbf{1}\rho_z + \mathbf{0} + \mathbf{0} - \mathbf{1}\mathbf{R}_z + \mathbf{0} + \mathbf{0} - \mathbf{1}\tilde{\mathbf{R}}_z + \mathbf{0} + \mathbf{0} + \mathbf{0} + \mathbf{0} + \mathbf{1}\mathbf{i}_4 + \mathbf{1}\mathbf{i}_3)$$

Review Octahedral $O_h \supset O$ group operator structure

Review Octahedral $O_h \supset O \supset D_4 \supset C_4$ subgroup chain correlations

Comparison of $O \supset D_4 \supset C_4$ and $O \supset D_4 \supset D_2$ correlations and level/projector splitting

$O \supset D_4 \supset C_4$ subgroup chain splitting

$O \supset D_4 \supset D_2$ subgroup chain splitting (normal D_2 vs. unnormal D_2)

$O_h \supset O \supset D_4 \supset C_{4v}$ and $O_h \supset O \supset D_4 \supset C_{4v} \supset C_{2v}$ subgroup splitting

Splitting O class projectors \mathbf{P}^μ into irreducible projectors $\mathbf{P}^\mu_{m_4m_4}$ for $O \supset C_4$

Development of irreducible projectors $\mathbf{P}^\mu_{m_4m_4}$ and representations $D^\mu_{m_4m_4}$

Calculating $\mathbf{P}^E_{0_40_4}$, $\mathbf{P}^E_{2_42_4}$, $\mathbf{P}^{T_1}_{0_40_4}$, $\mathbf{P}^{T_1}_{1_41_4}$, $\mathbf{P}^{T_2}_{2_42_4}$, $\mathbf{P}^{T_2}_{1_41_4}$,


$O \supset C_4$ induced representation $0_4(C_4) \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples

Elementary induced representation $0_4(C_4) \uparrow O$

Projection reduction of induced representation $0_4(C_4) \uparrow O$

Introduction to ortho-complete eigenvalue-parameter relations

Examples in SF_6 spectroscopy



Calculating $\mathbf{P}^{T_2}_{1414}$

$O \supset C_4$	0_4	1_4	2_4	3_4
$A_1 \downarrow C_4$	1	.	.	.
$A_2 \downarrow C_4$.	.	1	.
$E \downarrow C_4$	1	.	1	.
$T_1 \downarrow C_4$	1	1	.	1
$T_2 \downarrow C_4$.	1	1	1

$O: \chi_g^\mu$	$g=1$	r_{1-4}^p	ρ_{xyz}	R_{xyz}^p	i_{1-6}
$\mu=A_1$	1	1	1	1	1
A_2	1	1	1	-1	-1
E	2	-1	2	0	0
T_1	3	0	-1	1	-1
T_2	3	0	-1	-1	1

O characters

C_4 characters

$$d_{R^p}^{m_4} = e^{\frac{2\pi i m_4 \cdot p}{4}}$$

$$\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 e^{\frac{2\pi i m_4 \cdot p}{4}} \mathbf{R}_z^p =$$

$$\left\{ \begin{aligned} \mathbf{p}_{0_4} &= (\mathbf{1} + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{1_4} &= (\mathbf{1} + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{2_4} &= (\mathbf{1} - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z)/4 \\ \mathbf{p}_{3_4} &= (\mathbf{1} - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z)/4 \end{aligned} \right.$$

$$\mathbf{P}_{1414}^{T_2} = \mathbf{p}_{14} \mathbf{P}^{T_2} = \mathbf{P}^{T_2} \mathbf{p}_{14}$$

$$= \sum_g \frac{\ell^{T_2}}{\circ O} (\chi_g^{T_2}) \cdot \mathbf{g} \cdot (\mathbf{p}_{14}) = \sum_g \frac{3}{96} (\chi_g^{T_2}) \cdot \mathbf{g} \cdot (\mathbf{1} \cdot \mathbf{1} - \mathbf{1} \cdot \rho_z + i \cdot \mathbf{R}_z - i \cdot \tilde{\mathbf{R}}_z)$$

$$\mathbf{1}C_4 = \mathbf{1} \{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} \quad \rho_x C_4 = \{ \rho_x, \rho_y, \mathbf{i}_4, \mathbf{i}_3 \} \quad \mathbf{r}_1 C_4 = \{ \mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y \} \quad \mathbf{r}_2 C_4 = \{ \mathbf{r}_2, \mathbf{r}_3, \mathbf{i}_2, \tilde{\mathbf{R}}_y \} \quad \tilde{\mathbf{r}}_1 C_4 = \{ \tilde{\mathbf{r}}_1, \tilde{\mathbf{r}}_3, \tilde{\mathbf{R}}_x, \mathbf{i}_6 \} \quad \tilde{\mathbf{r}}_2 C_4 = \{ \tilde{\mathbf{r}}_2, \tilde{\mathbf{r}}_4, \mathbf{R}_x, \mathbf{i}_5 \}$$

$$= \frac{1}{32} \chi_{\mathbf{1}}^{T_2} (1, d_{\rho_z}^{14}, d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14}) + \frac{1}{32} \chi_{\rho_x}^{T_2} (1, d_{\rho_z}^{14}, d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14}) + \frac{1}{32} \chi_{\mathbf{r}_1}^{T_2} (1, d_{\rho_z}^{14}, d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14}) + \frac{1}{32} \chi_{\mathbf{r}_2}^{T_2} (1, d_{\rho_z}^{14}, d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14}) + \frac{1}{32} \chi_{\tilde{\mathbf{r}}_1}^{T_2} (1, d_{\rho_z}^{14}, d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14}) + \frac{1}{32} \chi_{\tilde{\mathbf{r}}_2}^{T_2} (1, d_{\rho_z}^{14}, d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14})$$

$$+ \frac{1}{32} \chi_{\rho_z}^{T_2} (d_{\rho_z}^{14}, 1, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14}) + \frac{1}{32} \chi_{\rho_y}^{T_2} (d_{\rho_z}^{14}, 1, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14}) + \frac{1}{32} \chi_{\mathbf{r}_4}^{T_2} (d_{\rho_z}^{14}, 1, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14}) + \frac{1}{32} \chi_{\mathbf{r}_3}^{T_2} (d_{\rho_z}^{14}, 1, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14}) + \frac{1}{32} \chi_{\tilde{\mathbf{r}}_3}^{T_2} (d_{\rho_z}^{14}, 1, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14}) + \frac{1}{32} \chi_{\tilde{\mathbf{r}}_4}^{T_2} (d_{\rho_z}^{14}, 1, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14})$$

$$+ \frac{1}{32} \chi_{\mathbf{R}_z}^{T_2} (d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14}, 1, d_{\rho_z}^{14}) + \frac{1}{32} \chi_{\mathbf{i}_4}^{T_2} (d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14}, 1, d_{\rho_z}^{14}) + \frac{1}{32} \chi_{\mathbf{i}_1}^{T_2} (d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14}, 1, d_{\rho_z}^{14}) + \frac{1}{32} \chi_{\mathbf{i}_2}^{T_2} (d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14}, 1, d_{\rho_z}^{14}) + \frac{1}{32} \chi_{\tilde{\mathbf{R}}_x}^{T_2} (d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14}, 1, d_{\rho_z}^{14}) + \frac{1}{32} \chi_{\mathbf{R}_x}^{T_2} (d_{\tilde{\mathbf{R}}_z}^{14}, d_{\mathbf{R}_z}^{14}, 1, d_{\rho_z}^{14})$$

$$+ \frac{1}{32} \chi_{\tilde{\mathbf{R}}_z}^{T_2} (d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\rho_z}^{14}, 1) + \frac{1}{32} \chi_{\mathbf{i}_3}^{T_2} (d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\rho_z}^{14}, 1) + \frac{1}{32} \chi_{\mathbf{R}_y}^{T_2} (d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\rho_z}^{14}, 1) + \frac{1}{32} \chi_{\tilde{\mathbf{R}}_y}^{T_2} (d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\rho_z}^{14}, 1) + \frac{1}{32} \chi_{\mathbf{i}_6}^{T_2} (d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\rho_z}^{14}, 1) + \frac{1}{32} \chi_{\mathbf{i}_5}^{T_2} (d_{\mathbf{R}_z}^{14}, d_{\tilde{\mathbf{R}}_z}^{14}, d_{\rho_z}^{14}, 1)$$

$$= \frac{1}{32} (+3)(1, -1, +i, -i) + \frac{1}{32} (-1)(1, -1, +i, -i) + \frac{1}{32} (0)(1, -1, +i, -i) + \frac{1}{32} (0)(1, -1, +i, -i) + \frac{1}{32} (0)(1, -1, +i, -i) + \frac{1}{32} (0)(1, -1, +i, -i)$$

$$+ \frac{1}{32} (-1)(-1, 1, -i, +i) + \frac{1}{32} (-1)(-1, 1, -i, +i) + \frac{1}{32} (0)(-1, 1, -i, +i) + \frac{1}{32} (0)(-1, 1, -i, +i) + \frac{1}{32} (0)(-1, 1, -i, +i) + \frac{1}{32} (0)(-1, 1, -i, +i)$$

$$+ \frac{1}{32} (-1)(-i, +i, 1, -1) + \frac{1}{32} (+1)(-i, +i, 1, -1) + \frac{1}{32} (+1)(-i, +i, 1, -1) + \frac{1}{32} (+1)(-i, +i, 1, -1) + \frac{1}{32} (-1)(-i, +i, 1, -1) + \frac{1}{32} (-1)(-i, +i, 1, -1)$$

$$+ \frac{1}{32} (-1)(+i, -i, -1, 1) + \frac{1}{32} (+1)(+i, -i, -1, 1) + \frac{1}{32} (-1)(+i, -i, -1, 1) + \frac{1}{32} (-1)(+i, -i, -1, 1) + \frac{1}{32} (+1)(+i, -i, -1, 1) + \frac{1}{32} (+1)(+i, -i, -1, 1)$$

$$\underline{+4, -4, 4i, -4i} \quad \underline{0, 0, 0, 0} \quad \underline{-2i, 2i, 2, -2} \quad \underline{-2i, 2i, 2, -2} \quad \underline{2i, -2i, -2, 2} \quad \underline{2i, -2i, -2, 2}$$

$$\frac{1}{8} (\underline{11-1\rho_z+i\mathbf{R}_z-i\tilde{\mathbf{R}}_z} \quad + \underline{0\rho_x+0\rho_y+0\mathbf{i}_4+0\mathbf{i}_3} \quad -\frac{i}{2}\mathbf{r}_1+\frac{i}{2}\mathbf{r}_4+\frac{1}{2}\mathbf{i}_1-\frac{1}{2}\mathbf{R}_y \quad -\frac{i}{2}\mathbf{r}_2+\frac{i}{2}\mathbf{r}_3+\frac{1}{2}\mathbf{i}_2-\frac{1}{2}\tilde{\mathbf{R}}_y \quad +\frac{i}{2}\tilde{\mathbf{r}}_1-\frac{i}{2}\tilde{\mathbf{r}}_3-\frac{1}{2}\tilde{\mathbf{R}}_x+\frac{1}{2}\mathbf{i}_6 \quad +\frac{i}{2}\tilde{\mathbf{r}}_2-\frac{i}{2}\tilde{\mathbf{r}}_4-\frac{1}{2}\mathbf{R}_x+\frac{1}{2}\mathbf{i}_5)$$

Coset-factored sum:

$$\mathbf{P}_{1414}^{T_2} = \frac{1}{8} [(1) \cdot \mathbf{1p}_{14} + (0) \cdot \rho_x \mathbf{p}_{14} + (-\frac{i}{2}) \cdot \mathbf{r}_1 \mathbf{p}_{14} + (-\frac{i}{2}) \cdot \mathbf{r}_2 \mathbf{p}_{14} + (-\frac{i}{2}) \cdot \tilde{\mathbf{r}}_1 \mathbf{p}_{14} + (-\frac{i}{2}) \cdot \tilde{\mathbf{r}}_2 \mathbf{p}_{14}]$$

Broken-class-ordered sum:

$$\mathbf{P}_{1414}^{T_2} = \frac{1}{8} (\mathbf{1} \cdot \mathbf{1} \quad -\frac{i}{2}\mathbf{r}_1 - \frac{i}{2}\mathbf{r}_2 \quad +\frac{i}{2}\mathbf{r}_3 + \frac{i}{2}\mathbf{r}_4 \quad +\frac{i}{2}\tilde{\mathbf{r}}_1 + \frac{i}{2}\tilde{\mathbf{r}}_2 \quad -\frac{i}{2}\tilde{\mathbf{r}}_3 - \frac{i}{2}\tilde{\mathbf{r}}_4 \quad +\mathbf{0}\rho_x + \mathbf{0}\rho_y - \mathbf{1}\rho_z \quad -\frac{1}{2}\mathbf{R}_x - \frac{1}{2}\mathbf{R}_y + i\mathbf{R}_z \quad -\frac{1}{2}\tilde{\mathbf{R}}_x - \frac{1}{2}\tilde{\mathbf{R}}_y - i\tilde{\mathbf{R}}_z \quad +\frac{1}{2}\mathbf{i}_1 + \frac{1}{2}\mathbf{i}_2 \quad +\mathbf{0}\mathbf{i}_3 + \mathbf{0}\mathbf{i}_4 \quad +\frac{1}{2}\mathbf{i}_5 + \frac{1}{2}\mathbf{i}_6)$$

Review Octahedral $O_h \supset O$ group operator structure

Review Octahedral $O_h \supset O \supset D_4 \supset C_4$ subgroup chain correlations

Comparison of $O \supset D_4 \supset C_4$ and $O \supset D_4 \supset D_2$ correlations and level/projector splitting

$O \supset D_4 \supset C_4$ subgroup chain splitting

$O \supset D_4 \supset D_2$ subgroup chain splitting (normal D_2 vs. unnormal D_2)

$O_h \supset O \supset D_4 \supset C_{4v}$ and $O_h \supset O \supset D_4 \supset C_{4v} \supset C_{2v}$ subgroup splitting

Splitting O class projectors P^μ into irreducible projectors $P^\mu_{m+m_4}$ for $O \supset C_4$

Development of irreducible projectors $P^\mu_{m+m_4}$ and representations $D^\mu_{m+m_4}$

Calculating $P^{E_{0404}}$, $P^{E_{2424}}$, $P^{T_{10404}}$, $P^{T_{11414}}$, $P^{T_{22424}}$, $P^{T_{21414}}$,

→ *$O \supset C_4$ induced representation $O_4(C_4) \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples* **←**

Elementary induced representation $O_4(C_4) \uparrow O$


Projection reduction of induced representation $O_4(C_4) \uparrow O$

Introduction to ortho-complete eigenvalue-parameter relations

Examples in SF_6 spectroscopy

$O \supset C_4$ induced representation $O_4 \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples

Fig. 4.3.1 PSDS



Solve XY_6 radial vibration $\mathbf{K}=\mathbf{a}$ -matrix


$$\begin{pmatrix} \langle 1 | \mathbf{a} | 1 \rangle & \langle 1 | \mathbf{a} | 2 \rangle & \cdots & \langle 1 | \mathbf{a} | 6 \rangle \\ \langle 2 | \mathbf{a} | 1 \rangle & \langle 2 | \mathbf{a} | 2 \rangle & \cdots & \langle 2 | \mathbf{a} | 6 \rangle \\ \cdot & & & \\ \cdot & h = 2k + t, & & \\ \cdot & s = k/2 & & \\ \langle 6 | \mathbf{a} | 1 \rangle & \langle 6 | \mathbf{a} | 2 \rangle & \cdots & \langle 6 | \mathbf{a} | 6 \rangle \end{pmatrix} = \begin{pmatrix} h & t & s & s & s & s \\ t & h & s & s & s & s \\ s & s & h & t & s & s \\ s & s & t & h & s & s \\ s & s & s & s & h & t \\ s & s & s & s & t & h \end{pmatrix},$$

Solve SF_6 J-tunneling Hamiltonian \mathbf{H}

$$\begin{pmatrix} \langle 1 | \mathbf{H} | 1 \rangle & \langle 1 | \mathbf{H} | 2 \rangle & \cdots & \langle 1 | \mathbf{H} | 6 \rangle \\ \langle 2 | \mathbf{H} | 1 \rangle & \langle 2 | \mathbf{H} | 2 \rangle & \cdots & \langle 2 | \mathbf{H} | 6 \rangle \\ \cdot & \cdot & & \cdot \\ \cdot & \cdot & & \cdot \\ \cdot & \cdot & & \cdot \\ \langle 6 | \mathbf{H} | 1 \rangle & \langle 6 | \mathbf{H} | 2 \rangle & \cdots & \langle 6 | \mathbf{H} | 6 \rangle \end{pmatrix} = \begin{pmatrix} H & T & S & S & S & S \\ T & H & S & S & S & S \\ S & S & H & T & S & S \\ S & S & T & H & S & S \\ S & S & S & S & H & T \\ S & S & S & S & T & H \end{pmatrix}$$

$O \supset C_4$ induced representation $O_4 \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples

Fig. 4.3.1 PSDS



Assuming C_4 -local symmetry conditions for $|1\rangle$ state


$$|1\rangle = \mathbf{1}|1\rangle = R_3|1\rangle = R_3^2|1\rangle = R_3^3|1\rangle$$

O operators (Two notations: Older Princ.of Symm.Dynamics and Spectra. and Newer Int.J.Mol.Sci)

PSDS:	1	\mathbf{r}_1	\mathbf{r}_2	\mathbf{r}_3	\mathbf{r}_4	\mathbf{r}_1^2	\mathbf{r}_2^2	\mathbf{r}_3^2	\mathbf{r}_4^2	\mathbf{R}_1^2	\mathbf{R}_2^2	\mathbf{R}_3^2	\mathbf{R}_1	\mathbf{R}_2	\mathbf{R}_3	\mathbf{R}_1^3	\mathbf{R}_2^3	\mathbf{R}_3^3	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6
IJMS:	1	\mathbf{r}_1	\mathbf{r}_2	\mathbf{r}_3	\mathbf{r}_4	$\tilde{\mathbf{r}}_1$	$\tilde{\mathbf{r}}_2$	$\tilde{\mathbf{r}}_3$	$\tilde{\mathbf{r}}_4$	ρ_x	ρ_y	ρ_z	\mathbf{R}_x	\mathbf{R}_y	\mathbf{R}_z	$\tilde{\mathbf{R}}_x$	$\tilde{\mathbf{R}}_y$	$\tilde{\mathbf{R}}_z$	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6

$O \supset C_4$ induced representation $O_4 \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples

Fig. 4.3.1 PSDS



Assuming C_4 -local symmetry conditions for $|1\rangle$ state

$$|1\rangle = \mathbf{1}|1\rangle = R_3|1\rangle = R_3^2|1\rangle = R_3^3|1\rangle$$

Using C_4 -local symmetry projector equations $P^A \equiv P^{O_4} = (\mathbf{1} + R_3 + R_3^2 + R_3^3)/4$.


$$|1\rangle = P^{O_4}|1\rangle = (\mathbf{1} + R_3 + R_3^2 + R_3^3)|1\rangle/4.$$

O operators (Two notations: Older Princ. of Symm. Dynamics and Spectra. and Newer Int. J. Mol. Sci)

PSDS:	$\mathbf{1}$	\mathbf{r}_1	\mathbf{r}_2	\mathbf{r}_3	\mathbf{r}_4	\mathbf{r}_1^2	\mathbf{r}_2^2	\mathbf{r}_3^2	\mathbf{r}_4^2	\mathbf{R}_1^2	\mathbf{R}_2^2	\mathbf{R}_3^2	\mathbf{R}_1	\mathbf{R}_2	\mathbf{R}_3	\mathbf{R}_1^3	\mathbf{R}_2^3	\mathbf{R}_3^3	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6
IJMS:	$\mathbf{1}$	\mathbf{r}_1	\mathbf{r}_2	\mathbf{r}_3	\mathbf{r}_4	$\tilde{\mathbf{r}}_1$	$\tilde{\mathbf{r}}_2$	$\tilde{\mathbf{r}}_3$	$\tilde{\mathbf{r}}_4$	ρ_x	ρ_y	ρ_z	\mathbf{R}_x	\mathbf{R}_y	\mathbf{R}_z	$\tilde{\mathbf{R}}_x$	$\tilde{\mathbf{R}}_y$	$\tilde{\mathbf{R}}_z$	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6

$O \supset C_4$ induced representation $O_4 \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples

Fig. 4.3.1 PSDS



Assuming C_4 -local symmetry conditions for $|1\rangle$ state

$$|1\rangle = \mathbf{1}|1\rangle = R_3|1\rangle = R_3^2|1\rangle = R_3^3|1\rangle$$

Using C_4 -local symmetry projector equations $P^A \equiv P^{O_4} = (\mathbf{1} + R_3 + R_3^2 + R_3^3)/4$.

$$|1\rangle = P^{O_4}|1\rangle = (\mathbf{1} + R_3 + R_3^2 + R_3^3)|1\rangle/4$$

These apply to all six $|g\rangle = g|1\rangle$ -base states. $|g\rangle = |gR_3\rangle = |gR_3^2\rangle = |gR_3^3\rangle$


$$|g\rangle = g|1\rangle = gR_3|1\rangle = gR_3^2|1\rangle = gR_3^3|1\rangle$$

O operators (Two notations: Older Princ. of Symm. Dynamics and Spectra. and Newer Int. J. Mol. Sci)

PSDS:	$\mathbf{1}$	\mathbf{r}_1	\mathbf{r}_2	\mathbf{r}_3	\mathbf{r}_4	\mathbf{r}_1^2	\mathbf{r}_2^2	\mathbf{r}_3^2	\mathbf{r}_4^2	\mathbf{R}_1^2	\mathbf{R}_2^2	\mathbf{R}_3^2	\mathbf{R}_1	\mathbf{R}_2	\mathbf{R}_3	\mathbf{R}_1^3	\mathbf{R}_2^3	\mathbf{R}_3^3	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6
IJMS:	$\mathbf{1}$	\mathbf{r}_1	\mathbf{r}_2	\mathbf{r}_3	\mathbf{r}_4	$\tilde{\mathbf{r}}_1$	$\tilde{\mathbf{r}}_2$	$\tilde{\mathbf{r}}_3$	$\tilde{\mathbf{r}}_4$	ρ_x	ρ_y	ρ_z	\mathbf{R}_x	\mathbf{R}_y	\mathbf{R}_z	$\tilde{\mathbf{R}}_x$	$\tilde{\mathbf{R}}_y$	$\tilde{\mathbf{R}}_z$	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6

$O \supset C_4$ induced representation $O_4 \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples

Fig. 4.3.1 PSDS



Thus we label states by left cosets $r_e C_4$ of Local C_4

Assuming C_4 -local symmetry conditions for $|1\rangle$ state

$$|1\rangle = \mathbf{1}|1\rangle = R_3|1\rangle = R_3^2|1\rangle = R_3^3|1\rangle$$

Using C_4 -local symmetry projector equations $P^A \equiv P^{O_4} = (\mathbf{1} + R_3 + R_3^2 + R_3^3)/4$.

$$|1\rangle = P^{O_4}|1\rangle = (\mathbf{1} + R_3 + R_3^2 + R_3^3)|1\rangle/4$$

These apply to all six $|g\rangle = g|1\rangle$ -base states. $|g\rangle = |gR_3\rangle = |gR_3^2\rangle = |gR_3^3\rangle$

$$|g\rangle = g|1\rangle = gR_3|1\rangle = gR_3^2|1\rangle = gR_3^3|1\rangle$$

O operators (Two notations: Older Princ.of Symm.Dynamics and Spectra. and Newer Int.J.Mol.Sci)

PSDS:	$\mathbf{1}$	\mathbf{r}_1	\mathbf{r}_2	\mathbf{r}_3	\mathbf{r}_4	\mathbf{r}_1^2	\mathbf{r}_2^2	\mathbf{r}_3^2	\mathbf{r}_4^2	\mathbf{R}_1^2	\mathbf{R}_2^2	\mathbf{R}_3^2	\mathbf{R}_1	\mathbf{R}_2	\mathbf{R}_3	\mathbf{R}_1^3	\mathbf{R}_2^3	\mathbf{R}_3^3	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6
IJMS:	$\mathbf{1}$	\mathbf{r}_1	\mathbf{r}_2	\mathbf{r}_3	\mathbf{r}_4	$\tilde{\mathbf{r}}_1$	$\tilde{\mathbf{r}}_2$	$\tilde{\mathbf{r}}_3$	$\tilde{\mathbf{r}}_4$	ρ_x	ρ_y	ρ_z	\mathbf{R}_x	\mathbf{R}_y	\mathbf{R}_z	$\tilde{\mathbf{R}}_x$	$\tilde{\mathbf{R}}_y$	$\tilde{\mathbf{R}}_z$	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6

$O \supset C_4$ induced representation $O_4 \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples





Fig. 4.3.1 PSDS

Thus we label states by left cosets $\mathbf{r}_e C_4$ of Local C_4

$\langle 5 | = \langle r_1^2 |$
 $\{ |1\rangle = |1\rangle, |2\rangle = |R_1^2\rangle, |3\rangle = |r_1\rangle, |4\rangle = |r_2\rangle, |5\rangle = |r_1^2\rangle, |6\rangle = |r_2^2\rangle \}$



$\mathbf{1}C_4 = C_4 = \{1, R_3, R_3^2, R_3^3\}$

$\mathbf{1}C_4 = C_4 = \{1, R_3, R_3^2, R_3^3\}$
 $R_1^2(\mathbf{1}, R_3, R_3^2, R_3^3) = (R_1^2, i_4, R_2^2, i_3)$
 $r_1(\mathbf{1}, R_3, R_3^2, R_3^3) = (r_1, i_1, r_4, R_2)$
 $r_2(\mathbf{1}, R_3, R_3^2, R_3^3) = (r_2, i_2, r_3, R_2^3)$
 $r_1^2(\mathbf{1}, R_3, R_3^2, R_3^3) = (r_1^2, R_1^3, r_3^2, i_6)$
 $r_2^2(\mathbf{1}, R_3, R_3^2, R_3^3) = (r_2^2, R_1, r_4^2, i_5)$

Compare to IJMS cosets on pages 25 -60:

Assuming C_4 -local symmetry conditions for $|1\rangle$ state

$|1\rangle = \mathbf{1}|1\rangle = R_3|1\rangle = R_3^2|1\rangle = R_3^3|1\rangle$

Using C_4 -local symmetry projector equations $P^A \equiv P^{O_4} = (\mathbf{1} + R_3 + R_3^2 + R_3^3)/4$

$|1\rangle = P^{O_4}|1\rangle = (\mathbf{1} + R_3 + R_3^2 + R_3^3)|1\rangle/4$

These apply to all six $|g\rangle = g|1\rangle$ -base states. $|g\rangle = |gR_3\rangle = |gR_3^2\rangle = |gR_3^3\rangle$

$|g\rangle = g|1\rangle = gR_3|1\rangle = gR_3^2|1\rangle = gR_3^3|1\rangle$

Switch columns 1 with 2

- $\{ \mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z \}$
- $\{ \rho_x, \rho_y, i_4, i_3 \}$
- $\{ r_1, r_4, i_1, \mathbf{R}_y \}$
- $\{ r_2, r_3, i_2, \tilde{\mathbf{R}}_y \}$
- $\{ \tilde{r}_1, \tilde{r}_3, \tilde{\mathbf{R}}_x, i_6 \}$
- $\{ \tilde{r}_2, \tilde{r}_4, \mathbf{R}_x, i_5 \}$

O operators (Two notations: Older Princ. of Symm. Dynamics and Spectra. and Newer Int. J. Mol. Sci)

PSDS:	$\mathbf{1}$	r_1	r_2	r_3	r_4	r_1^2	r_2^2	r_3^2	r_4^2	\mathbf{R}_1^2	\mathbf{R}_2^2	\mathbf{R}_3^2	\mathbf{R}_1	\mathbf{R}_2	\mathbf{R}_3	\mathbf{R}_1^3	\mathbf{R}_2^3	\mathbf{R}_3^3	i_1	i_2	i_3	i_4	i_5	i_6
IJMS:	$\mathbf{1}$	r_1	r_2	r_3	r_4	\tilde{r}_1	\tilde{r}_2	\tilde{r}_3	\tilde{r}_4	ρ_x	ρ_y	ρ_z	\mathbf{R}_x	\mathbf{R}_y	\mathbf{R}_z	$\tilde{\mathbf{R}}_x$	$\tilde{\mathbf{R}}_y$	$\tilde{\mathbf{R}}_z$	i_1	i_2	i_3	i_4	i_5	i_6

Review Octahedral $O_h \supset O$ group operator structure

Review Octahedral $O_h \supset O \supset D_4 \supset C_4$ subgroup chain correlations

Comparison of $O \supset D_4 \supset C_4$ and $O \supset D_4 \supset D_2$ correlations and level/projector splitting

$O \supset D_4 \supset C_4$ subgroup chain splitting

$O \supset D_4 \supset D_2$ subgroup chain splitting (normal D_2 vs. unnormal D_2)

$O_h \supset O \supset D_4 \supset C_{4v}$ and $O_h \supset O \supset D_4 \supset C_{4v} \supset C_{2v}$ subgroup splitting

Splitting O class projectors P^μ into irreducible projectors $P^\mu_{m+m_4}$ for $O \supset C_4$

Development of irreducible projectors $P^\mu_{m+m_4}$ and representations $D^\mu_{m+m_4}$

Calculating $P^{E_{0404}}$, $P^{E_{2424}}$, $P^{T_{10404}}$, $P^{T_{11414}}$, $P^{T_{22424}}$, $P^{T_{21414}}$,

$O \supset C_4$ induced representation $O_4(C_4) \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples

Elementary induced representation $O_4(C_4) \uparrow O$

Projection reduction of induced representation $O_4(C_4) \uparrow O$

Introduction to ortho-complete eigenvalue-parameter relations

Examples in SF_6 spectroscopy

Elementary induced representation $0_4(C_4) \uparrow O$





Fig. 4.3.1 PSDS



Thus we label states by left cosets $r_l C_4$ of Local C_4

$$\{|1\rangle = |1\rangle, |2\rangle = |R_1^2\rangle, |3\rangle = |r_1\rangle, |4\rangle = |r_2\rangle, |5\rangle = |r_1^2\rangle, |6\rangle = |r_2^2\rangle\}$$

$$1C_4 = C_4 = \{1, R_3, R_3^2, R_3^3\}$$

$$1C_4 = C_4 = \{1, R_3, R_3^2, R_3^3\}$$

$$R_1^2(1, R_3, R_3^2, R_3^3) = (R_1^2, i_4, R_2^2, i_3),$$

$$r_1(1, R_3, R_3^2, R_3^3) = (r_1, i_1, r_4, R_2),$$

$$r_2(1, R_3, R_3^2, R_3^3) = (r_2, i_2, r_3, R_2^3),$$

$$r_1^2(1, R_3, R_3^2, R_3^3) = (r_1^2, R_1^3, r_3^2, i_6),$$

$$r_2^2(1, R_3, R_3^2, R_3^3) = (r_2^2, R_1, r_4^2, i_5),$$

This "coset-basis" spans a scalar $0_4(C_4)$ induced representation $0_4(C_4) \uparrow O$

$$\begin{aligned} \mathbf{i}_4|1\rangle &= \mathbf{i}_4|1\rangle, & \mathbf{i}_4|2\rangle &= \mathbf{i}_4 R_1^2|1\rangle, & \mathbf{i}_4|3\rangle &= \mathbf{i}_4 r_1|1\rangle, & \mathbf{i}_4|4\rangle &= \mathbf{i}_4 r_2|1\rangle, & \mathbf{i}_4|5\rangle &= \mathbf{i}_4 r_1^2|1\rangle, & \mathbf{i}_4|6\rangle &= \mathbf{i}_4 r_2^2|1\rangle. \\ &= R_1^2|1\rangle, & &= R_3^3|1\rangle, & &= i_5|1\rangle, & &= i_6|1\rangle, & &= i_2|1\rangle, & &= i_1|1\rangle. \\ &= |2\rangle, & &= |1\rangle, & &= |6\rangle, & &= |5\rangle, & &= |4\rangle, & &= |3\rangle. \end{aligned}$$

Elementary induced representation $O_4(C_4) \uparrow O$





Fig. 4.3.1 PSDS



Thus we label states by left cosets $r_l C_4$ of Local C_4

$$\{|1\rangle = |1\rangle, |2\rangle = |R_1^2\rangle, |3\rangle = |r_1\rangle, |4\rangle = |r_2\rangle, |5\rangle = |r_1^2\rangle, |6\rangle = |r_2^2\rangle\}$$

$$1C_4 = C_4 = \{1, R_3, R_3^2, R_3^3\}$$

$$1C_4 = C_4 = \{1, R_3, R_3^2, R_3^3\}$$

$$R_1^2(1, R_3, R_3^2, R_3^3) = (R_1^2, i_4, R_2^2, i_3),$$

$$r_1(1, R_3, R_3^2, R_3^3) = (r_1, i_1, r_4, R_2),$$

$$r_2(1, R_3, R_3^2, R_3^3) = (r_2, i_2, r_3, R_2^3),$$

$$r_1^2(1, R_3, R_3^2, R_3^3) = (r_1^2, R_1^3, r_3^2, i_6),$$

$$r_2^2(1, R_3, R_3^2, R_3^3) = (r_2^2, R_1, r_4^2, i_5),$$

This “coset-basis” spans a scalar $O_4(C_4)$ induced representation $O_4(C_4) \uparrow O$

$$\begin{aligned} \mathbf{i}_4|1\rangle &= \mathbf{i}_4|1\rangle, & \mathbf{i}_4|2\rangle &= \mathbf{i}_4 R_1^2|1\rangle, & \mathbf{i}_4|3\rangle &= \mathbf{i}_4 r_1|1\rangle, & \mathbf{i}_4|4\rangle &= \mathbf{i}_4 r_2|1\rangle, & \mathbf{i}_4|5\rangle &= \mathbf{i}_4 r_1^2|1\rangle, & \mathbf{i}_4|6\rangle &= \mathbf{i}_4 r_2^2|1\rangle. \\ &= R_1^2|1\rangle, & &= R_3^3|1\rangle, & &= i_5|1\rangle, & &= i_6|1\rangle, & &= i_2|1\rangle, & &= i_1|1\rangle. \\ &= |2\rangle, & &= |1\rangle, & &= |6\rangle, & &= |5\rangle, & &= |4\rangle, & &= |3\rangle. \end{aligned}$$

For example here is $O_4(C_4)$ induced representation $O_4(C_4) \uparrow O(\mathbf{i}_4)$

$$\mathcal{S}^{O_4 \uparrow O}(\mathbf{i}_4) = \begin{pmatrix} \langle 1|\mathbf{i}_4|1\rangle & \langle 1|\mathbf{i}_4|2\rangle & \cdots & \langle 1|\mathbf{i}_4|6\rangle \\ \langle 2|\mathbf{i}_4|1\rangle & \langle 2|\mathbf{i}_4|2\rangle & & \vdots \\ \vdots & \vdots & & \vdots \\ \langle 6|\mathbf{i}_4|1\rangle & \langle 6|\mathbf{i}_4|2\rangle & & \langle 1|\mathbf{i}_4|6\rangle \end{pmatrix} = \begin{pmatrix} |1\rangle & |2\rangle & |3\rangle & |4\rangle & |5\rangle & |6\rangle \\ \langle 1| & 1 & \cdot & \cdot & \cdot & \cdot \\ \langle 2| & 1 & \cdot & \cdot & \cdot & \cdot \\ \langle 3| & \cdot & \cdot & \cdot & \cdot & 1 \\ \langle 4| & \cdot & \cdot & \cdot & 1 & \cdot \\ \langle 5| & \cdot & \cdot & 1 & \cdot & \cdot \\ \langle 6| & \cdot & 1 & \cdot & \cdot & \cdot \end{pmatrix}$$

Elementary induced representation $O_4(C_4) \uparrow O$





Fig. 4.3.1 PSDS



Thus we label states by left cosets $r_l C_4$ of Local C_4

$$\{ |1\rangle = |1\rangle, |2\rangle = |R_1^2\rangle, |3\rangle = |r_1\rangle, |4\rangle = |r_2\rangle, |5\rangle = |r_1^2\rangle, |6\rangle = |r_2^2\rangle \}$$

$$1C_4 = C_4 = \{1, R_3, R_3^2, R_3^3\}$$

$$R_1^2(1, R_3, R_3^2, R_3^3) = (R_1^2, i_4, R_2^2, i_3),$$

$$r_1(1, R_3, R_3^2, R_3^3) = (r_1, i_1, r_4, R_2),$$

$$r_2(1, R_3, R_3^2, R_3^3) = (r_2, i_2, r_3, R_2^3),$$

$$r_1^2(1, R_3, R_3^2, R_3^3) = (r_1^2, R_1^3, r_3^2, i_6),$$

$$r_2^2(1, R_3, R_3^2, R_3^3) = (r_2^2, R_1, r_4^2, i_5),$$

This "coset-basis" spans a scalar $O_4(C_4)$ induced representation $O_4(C_4) \uparrow O$

$$\begin{aligned} \mathbf{i}_4|1\rangle &= \mathbf{i}_4|1\rangle, & \mathbf{i}_4|2\rangle &= \mathbf{i}_4 R_1^2|1\rangle, & \mathbf{i}_4|3\rangle &= \mathbf{i}_4 r_1|1\rangle, & \mathbf{i}_4|4\rangle &= \mathbf{i}_4 r_2|1\rangle, & \mathbf{i}_4|5\rangle &= \mathbf{i}_4 r_1^2|1\rangle, & \mathbf{i}_4|6\rangle &= \mathbf{i}_4 r_2^2|1\rangle. \\ &= R_1^2|1\rangle, & &= R_3^3|1\rangle, & &= i_5|1\rangle, & &= i_6|1\rangle, & &= i_2|1\rangle, & &= i_1|1\rangle. \\ &= |2\rangle, & &= |1\rangle, & &= |6\rangle, & &= |5\rangle, & &= |4\rangle, & &= |3\rangle. \end{aligned}$$

For example here is $O_4(C_4)$ induced representation $O_4(C_4) \uparrow O(\mathbf{i}_4)$ and $O_4(C_4) \uparrow O(\mathbf{i}_3)$

$$\mathcal{F}^{O_4 \uparrow O}(\mathbf{i}_4) = \begin{pmatrix} \langle 1|\mathbf{i}_4|1\rangle & \langle 1|\mathbf{i}_4|2\rangle & \cdots & \langle 1|\mathbf{i}_4|6\rangle \\ \langle 2|\mathbf{i}_4|1\rangle & \langle 2|\mathbf{i}_4|2\rangle & & \vdots \\ \vdots & & & \vdots \\ \langle 6|\mathbf{i}_4|1\rangle & \langle 6|\mathbf{i}_4|2\rangle & & \langle 1|\mathbf{i}_4|6\rangle \end{pmatrix} = \begin{pmatrix} |1\rangle & |2\rangle & |3\rangle & |4\rangle & |5\rangle & |6\rangle \\ \langle 1| & 1 & \cdot & \cdot & \cdot & \cdot \\ \langle 2| & 1 & \cdot & \cdot & \cdot & \cdot \\ \langle 3| & \cdot & \cdot & \cdot & \cdot & 1 \\ \langle 4| & \cdot & \cdot & \cdot & 1 & \cdot \\ \langle 5| & \cdot & \cdot & 1 & \cdot & \cdot \\ \langle 6| & \cdot & 1 & \cdot & \cdot & \cdot \end{pmatrix} \quad \mathcal{F}^{O_4 \uparrow O}(\mathbf{i}_3) = \begin{pmatrix} \cdot & 1 & \cdot & \cdot & \cdot & \cdot \\ 1 & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & 1 & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & 1 \\ \cdot & \cdot & 1 & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & 1 & \cdot & \cdot \end{pmatrix}$$

Elementary induced representation $0_4(C_4) \uparrow O$





Fig. 4.3.1 PSDS



Thus we label states by left cosets $r_l C_4$ of Local C_4

$$\{|1\rangle = |1\rangle, |2\rangle = |R_1^2\rangle, |3\rangle = |r_1\rangle, |4\rangle = |r_2\rangle, |5\rangle = |r_1^2\rangle, |6\rangle = |r_2^2\rangle\}$$

$$1C_4 = C_4 = \{1, R_3, R_3^2, R_3^3\}$$

$$1C_4 = C_4 = \{1, R_3, R_3^2, R_3^3\}$$

$$R_1^2(1, R_3, R_3^2, R_3^3) = (R_1^2, i_4, R_2^2, i_3),$$

$$r_1(1, R_3, R_3^2, R_3^3) = (r_1, i_1, r_4, R_2),$$

$$r_2(1, R_3, R_3^2, R_3^3) = (r_2, i_2, r_3, R_2^3),$$

$$r_1^2(1, R_3, R_3^2, R_3^3) = (r_1^2, R_1^3, r_3^2, i_6),$$

$$r_2^2(1, R_3, R_3^2, R_3^3) = (r_2^2, R_1, r_4^2, i_5),$$

Here is $0_4(C_4)$ induced representation $\mathcal{J}^{0_4 \uparrow O}(\mathbf{I}_i)$ of a linear combination of \mathbf{i} -class rotations

$$\mathbf{I}_i = i_1 \mathbf{i}_1 + i_2 \mathbf{i}_2 + i_3 \mathbf{i}_3 + i_4 \mathbf{i}_4 + i_5 \mathbf{i}_5 + i_6 \mathbf{i}_6$$

$$\mathcal{J}^{0_4 \uparrow O}(\mathbf{I}_i) =$$

	$ 1\rangle$	$ 2\rangle$	$ 3\rangle$	$ 4\rangle$	$ 5\rangle$	$ 6\rangle$
$\langle 1 $	1	$i_3 + i_4$	i_1	i_2	i_6	i_5
$\langle 2 $	$i_3 + i_4$	1	i_2	i_1	i_5	i_6
$\langle 3 $	i_1	i_2	1	$i_5 + i_6$	i_3	i_4
$\langle 4 $	i_2	i_1	$i_5 + i_6$	1	i_4	i_3
$\langle 5 $	i_6	i_5	i_3	i_4	1	$i_1 + i_2$
$\langle 6 $	i_5	i_6	i_4	i_3	$i_1 + i_2$	1

$$\mathcal{J}^{0_4 \uparrow O}(\mathbf{i}_3) = \begin{pmatrix} \cdot & 1 & \cdot & \cdot & \cdot & \cdot \\ 1 & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & 1 & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & 1 \\ \cdot & \cdot & 1 & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & 1 & \cdot & \cdot \end{pmatrix}$$

Review Octahedral $O_h \supset O$ group operator structure

Review Octahedral $O_h \supset O \supset D_4 \supset C_4$ subgroup chain correlations

Comparison of $O \supset D_4 \supset C_4$ and $O \supset D_4 \supset D_2$ correlations and level/projector splitting

$O \supset D_4 \supset C_4$ subgroup chain splitting

$O \supset D_4 \supset D_2$ subgroup chain splitting (normal D_2 vs. unnormal D_2)

$O_h \supset O \supset D_4 \supset C_{4v}$ and $O_h \supset O \supset D_4 \supset C_{4v} \supset C_{2v}$ subgroup splitting

Splitting O class projectors P^μ into irreducible projectors $P^\mu_{m+m_4}$ for $O \supset C_4$

Development of irreducible projectors $P^\mu_{m+m_4}$ and representations $D^\mu_{m+m_4}$

Calculating $P^{E_{0404}}$, $P^{E_{2424}}$, $P^{T_{10404}}$, $P^{T_{11414}}$, $P^{T_{22424}}$, $P^{T_{21414}}$,



$O \supset C_4$ induced representation $O_4(C_4) \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples

Elementary induced representation $O_4(C_4) \uparrow O$

Projection reduction of induced representation $O_4(C_4) \uparrow O$

Introduction to ortho-complete eigenvalue-parameter relations

Examples in SF_6 spectroscopy




Projection reduction of induced representation $O_4(C_4)\uparrow O$


Scalar A_1 eigenket

$$\begin{aligned} |e_{0_4 0_4}^{A_1}\rangle &= \mathbf{P}_{0_4 0_4}^{A_1} |1\rangle / \sqrt{N^{A_1}} \\ &= \frac{1}{24} \sum_{p=1}^{24} D_{0_4 0_4}^{A_1*}(g_p) \mathbf{g}_p |1\rangle / \sqrt{N^{A_1}} \\ &= (|1\rangle + |2\rangle + |3\rangle + |4\rangle + |5\rangle + |6\rangle) / \sqrt{6} \end{aligned}$$

$$|e_{0_4 0_4}^{A_1}\rangle = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \frac{1}{\sqrt{6}}$$



A_1 $H + 4S$
 FREQUENCY OR ENERGY SPECTRUM




$$|e_{0_4 0_4}^{A_1}\rangle = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \frac{1}{\sqrt{6}}$$

Projection reduction of induced representation $O_4(C_4)\uparrow O$

Scalar A_1 eigenket 0_40_4

$$\begin{aligned} |e_{0_40_4}^{A_1}\rangle &= \mathbf{P}_{0_40_4}^{A_1} |1\rangle / \sqrt{N^{A_1}} \\ &= \frac{1}{24} \sum_{p=1}^{24} D_{0_40_4}^{A_1*}(g_p) \mathbf{g}_p |1\rangle / \sqrt{N^{A_1}} \\ &= (|1\rangle + |2\rangle + |3\rangle + |4\rangle + |5\rangle + |6\rangle) / \sqrt{6} \end{aligned}$$




Tensor E -eigenket 0_40_4

Diagonal (idempotent) Projector $\mathbf{P}_{jj}^{A_1}$
From p.49-50:


$$\begin{aligned} |e_{0_40_4}^E\rangle &= \mathbf{P}_{0_40_4}^E |1\rangle / \sqrt{N^E} \\ &= \frac{2}{24} \sum_{p=1}^{24} D_{0_40_4}^{E*}(g_p) \mathbf{g}_p |1\rangle / \sqrt{N^E} \\ &= (|1\rangle + |2\rangle - \frac{1}{2}|3\rangle - \frac{1}{2}|4\rangle - \frac{1}{2}|5\rangle - \frac{1}{2}|6\rangle) / \sqrt{3} \end{aligned}$$

$$\mathbf{P}_{0_40_4}^E = \frac{1}{12} [(\mathbf{1}) \cdot \mathbf{1} \mathbf{p}_{0_4} + (\mathbf{1}) \cdot \rho_x \mathbf{p}_{0_4} + (-\frac{1}{2}) \cdot \mathbf{r}_1 \mathbf{p}_{0_4} + (-\frac{1}{2}) \cdot \mathbf{r}_2 \mathbf{p}_{0_4} + (-\frac{1}{2}) \cdot \tilde{\mathbf{r}}_1 \mathbf{p}_{0_4} + (-\frac{1}{2}) \cdot \tilde{\mathbf{r}}_2 \mathbf{p}_{0_4}]$$


$$\{\mathbf{1}, \rho_z, \mathbf{R}_z, \tilde{\mathbf{R}}_z\} \{\rho_x, \rho_y, \mathbf{i}_3, \mathbf{i}_4\} \{\mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y\} \{\mathbf{r}_2, \mathbf{r}_3, \mathbf{i}_2, \tilde{\mathbf{R}}_y\} \{\tilde{\mathbf{r}}_1, \tilde{\mathbf{r}}_3, \tilde{\mathbf{R}}_x, \mathbf{i}_6\} \{\tilde{\mathbf{r}}_2, \tilde{\mathbf{r}}_4, \mathbf{R}_x, \mathbf{i}_5\}$$



$$|e_{0_40_4}^{A_1}\rangle = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \frac{1}{\sqrt{6}}$$



A_1 $H + 4S$
FREQUENCY OR ENERGY SPECTRUM



$$|e_{0_40_4}^{A_1}\rangle = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \frac{1}{\sqrt{6}}$$

MolVibes Web Simulation - 6 Atom 3D vibrations with O_h symmetry

Projection reduction of induced representation $O_4(C_4)\uparrow O$

Scalar A_1 eigenket 0_40_4

$$\begin{aligned} |e_{0_40_4}^{A_1}\rangle &= \mathbf{P}_{0_40_4}^{A_1} |1\rangle / \sqrt{N^{A_1}} \\ &= \frac{1}{24} \sum_{p=1}^{24} D_{0_40_4}^{A_1*}(g_p) \mathbf{g}_p |1\rangle / \sqrt{N^{A_1}} \\ &= (|1\rangle + |2\rangle + |3\rangle + |4\rangle + |5\rangle + |6\rangle) / \sqrt{6} \end{aligned}$$


Off-Diagonal
(nilpotent)

Projector $\mathbf{P}^{l_{jk}}$

Derived next lectures

Tensor E -eigenket 2_40_4

$$\begin{aligned} |e_{2_40_4}^E\rangle &= \mathbf{P}_{2_40_4}^E |1\rangle / \sqrt{N^E} \\ &= \frac{2}{24} \sum_{p=1}^{24} D_{2_40_4}^{E*}(g_p) \mathbf{g}_p |1\rangle / \sqrt{N^E} \\ &= (|3\rangle + |4\rangle - |5\rangle - |6\rangle) / 2 \end{aligned}$$



Projection reduction of induced representation $O_4(C_4)\uparrow O$

Scalar A_1 eigenket 0_40_4

$$\begin{aligned} |e_{0_40_4}^{A_1}\rangle &= \mathbf{P}_{0_40_4}^{A_1} |1\rangle / \sqrt{N^{A_1}} \\ &= \frac{1}{24} \sum_{p=1}^{24} D_{0_40_4}^{A_1*}(g_p) \mathbf{g}_p |1\rangle / \sqrt{N^{A_1}} \\ &= (|1\rangle + |2\rangle + |3\rangle + |4\rangle + |5\rangle + |6\rangle) / \sqrt{6} \end{aligned}$$

Vector T_1 -eigenket 0_40_4


$$\begin{aligned} |e_{0_40_4}^{T_1}\rangle &= \mathbf{P}_{0_40_4}^{T_1} |1\rangle / \sqrt{N^{T_1}} \\ &= \frac{1}{24} \sum_{p=1}^{24} D_{0_40_4}^{T_1*}(g_p) \mathbf{g}_p |1\rangle / \sqrt{N^{T_1}} \\ &= (|1\rangle - |2\rangle + 0 + 0 + 0 + 0) / \sqrt{2} \end{aligned}$$

Diagonal
(idempotent)
Projector \mathbf{P}_{jj}^{U}
From p.53:

$$\mathbf{P}_{0_40_4}^{T_1} = \frac{1}{8} [(1) \cdot \mathbf{1} \mathbf{p}_{0_4} + (-1) \cdot \rho_x \mathbf{p}_{0_4} + (0) \cdot \mathbf{r}_1 \mathbf{p}_{0_4} + (0) \cdot \mathbf{r}_2 \mathbf{p}_{0_4} + (0) \cdot \tilde{\mathbf{r}}_1 \mathbf{p}_{0_4} + (0) \cdot \tilde{\mathbf{r}}_2 \mathbf{p}_{0_4}]$$



$$\{ \mathbf{1}, \rho_x, \mathbf{R}_z, \tilde{\mathbf{R}}_z \} \{ \rho_x, \rho_y, \mathbf{i}_3, \mathbf{i}_4 \} \{ \mathbf{r}_1, \mathbf{r}_4, \mathbf{i}_1, \mathbf{R}_y \} \{ \mathbf{r}_2, \mathbf{r}_3, \mathbf{i}_2, \tilde{\mathbf{R}}_y \} \{ \tilde{\mathbf{r}}_1, \tilde{\mathbf{r}}_3, \tilde{\mathbf{R}}_x, \mathbf{i}_6 \} \{ \tilde{\mathbf{r}}_2, \tilde{\mathbf{r}}_4, \mathbf{R}_x, \mathbf{i}_5 \}$$

$$|e_{0_40_4}^E\rangle = \begin{pmatrix} 2 \\ 2 \\ -1 \\ -1 \\ -1 \\ -1 \end{pmatrix} \frac{1}{2\sqrt{3}}$$




E

T₁




$$|e_{2_40_4}^E\rangle = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 1 \\ -1 \\ -1 \end{pmatrix} \frac{1}{2}$$




$$|T_1\rangle_{0_40_4} = \begin{pmatrix} 1 \\ -1 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} \frac{1}{\sqrt{2}}$$

$$|e_{0_40_4}^{A_1}\rangle = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \frac{1}{\sqrt{6}}$$



A₁

FREQUENCY OR ENERGY
SPECTRUM



$$|e_{0_40_4}^{A_1}\rangle = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \frac{1}{\sqrt{6}}$$

Projection reduction of induced representation $O_4(C_4)\uparrow O$

Scalar A_1 eigenket 0_40_4

$$\begin{aligned} |e_{0_40_4}^{A_1}\rangle &= \mathbf{P}_{0_40_4}^{A_1} |1\rangle / \sqrt{N^{A_1}} \\ &= \frac{1}{24} \sum_{p=1}^{24} D_{0_40_4}^{A_1*}(g_p) \mathbf{g}_p |1\rangle / \sqrt{N^{A_1}} \\ &= (|1\rangle + |2\rangle + |3\rangle + |4\rangle + |5\rangle + |6\rangle) / \sqrt{6} \end{aligned}$$

Vector T_1 -eigenket $\pm 1_40_4$ and 0_40_4


$$\begin{aligned} |e_{\pm 1_40_4}^{T_1}\rangle &= \mathbf{P}_{1_40_4}^{T_1} |1\rangle / \sqrt{N^{T_1}} \\ &= \frac{1}{24} \sum_{p=1}^{24} D_{\pm 1_40_4}^{T_1*}(g_p) \mathbf{g}_p |1\rangle / \sqrt{N^{T_1}} \\ &= (0 + 0 + |3\rangle + |4\rangle \pm i|5\rangle \pm i|6\rangle) / 2 \end{aligned}$$

Off-Diagonal
(nilpotent)

Projector \mathbf{P}_{jk}^{μ}

Derived next lectures


$$|e_{0_40_4}^E\rangle = \begin{pmatrix} 2 \\ 2 \\ -1 \\ -1 \\ -1 \\ -1 \end{pmatrix} \frac{1}{2\sqrt{3}}$$







E

T₁

$$|T_{1_40_4}^{T_1}\rangle = \begin{pmatrix} 0 \\ 0 \\ -1 \\ 1 \\ -i \\ i \end{pmatrix} \frac{1}{2}$$



$$|e_{2_40_4}^E\rangle = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 1 \\ -1 \\ -1 \end{pmatrix} \frac{1}{2}$$




$\pm 1_4 = x \pm iy$

$0_4 = z$


$$|T_{1_40_4}^{T_1}\rangle = \begin{pmatrix} 1 \\ -1 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} \frac{1}{\sqrt{2}}$$

$$|e_{0_40_4}^{A_1}\rangle = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \frac{1}{\sqrt{6}}$$



A₁

FREQUENCY OR ENERGY SPECTRUM



$$|e_{0_40_4}^{A_1}\rangle = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \frac{1}{\sqrt{6}}$$

Projection reduction of induced representation $O_4(C_4)\uparrow O$

Tunneling T (next-neighbor) is too "Tiny" to contribute to E^a

$$E^{A_1} = H + T + 4S$$

$$E^{T_1} = H - T$$

$$E^E = H + T - 2S$$







Figure 4.3.3 Evidence of an (A_1T_1E) spectral cluster in methane laser spectra. (Courtesy of Dr. Allan Pine, MIT Lincoln Laboratories, from *Journal of Optical Society of America* 66, 97 (1976)). The ordering and approximate spacing of the A_1T_1 and E lines is consistent with that of Figure 4.3.2.

$$\begin{pmatrix} \langle 1|\mathbf{H}|1\rangle & \langle 1|\mathbf{H}|2\rangle & \cdots & \langle 1|\mathbf{H}|6\rangle \\ \langle 2|\mathbf{H}|1\rangle & \langle 2|\mathbf{H}|2\rangle & \cdots & \langle 2|\mathbf{H}|6\rangle \\ \vdots & \vdots & \ddots & \vdots \\ \vdots & \vdots & \vdots & \vdots \\ \langle 6|\mathbf{H}|1\rangle & \langle 6|\mathbf{H}|2\rangle & \cdots & \langle 6|\mathbf{H}|6\rangle \end{pmatrix} = \begin{pmatrix} H & T & S & S & S & S \\ T & H & S & S & S & S \\ S & S & H & T & S & S \\ S & S & T & H & S & S \\ S & S & S & S & H & T \\ S & S & S & S & T & H \end{pmatrix}$$

$$|E_{0_4 0_4}\rangle = \frac{1}{2\sqrt{3}} \begin{pmatrix} 2 \\ 2 \\ -1 \\ -1 \\ -1 \\ -1 \end{pmatrix}$$




$$\underline{\underline{E_g}} \quad H - 2S$$




$$\underline{\underline{T_{1u}}} \quad H$$

$$|T_{1_4 0_4}\rangle = \frac{1}{2} \begin{pmatrix} 0 \\ 0 \\ -1 \\ 1 \\ -i \\ i \end{pmatrix}$$




$$|e_{0_4 0_4}^{A_1}\rangle = \frac{1}{\sqrt{6}} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$$



$$\underline{\underline{A_{1g}}} \quad H + 4S$$

FREQUENCY OR ENERGY SPECTRUM



Projection reduction of induced representation $O_4(C_4)\uparrow O$

Tunneling T (next-neighbor) is too "Tiny" to contribute to E^a

$$E^{A_1} = H + T + 4S$$

$$E^{T_1} = H - T$$

$$E^E = H + T - 2S$$




Figure 4.3.3 Evidence of an (A_1T_1E) spectral cluster in methane laser spectra. (Courtesy of Dr. Allan Pine, MIT Lincoln Laboratories, from *Journal of Optical Society of America* 66, 97 (1976)). The ordering and approximate spacing of the A_1T_1 and E lines is consistent with that of Figure 4.3.2.

$$\begin{pmatrix} \langle 1|\mathbf{H}|1\rangle & \langle 1|\mathbf{H}|2\rangle & \cdots & \langle 1|\mathbf{H}|6\rangle \\ \langle 2|\mathbf{H}|1\rangle & \langle 2|\mathbf{H}|2\rangle & \cdots & \langle 2|\mathbf{H}|6\rangle \\ \vdots & \vdots & \ddots & \vdots \\ \langle 6|\mathbf{H}|1\rangle & \langle 6|\mathbf{H}|2\rangle & \cdots & \langle 6|\mathbf{H}|6\rangle \end{pmatrix} = \begin{pmatrix} H & T & S & S & S & S \\ T & H & S & S & S & S \\ S & S & H & T & S & S \\ S & S & T & H & S & S \\ S & S & S & S & H & T \\ S & S & S & S & T & H \end{pmatrix}$$


$O_h \supset D_{4h} \supset C_{4v} \supset C_{2v}$ subgroup splitting

Labels correct u or g parity!

$O_h \supset C_{4v}$	A'	B'	A''	B''	E
$A_{1g} \downarrow C_{4v}$	1
$A_{2g} \downarrow C_{4v}$.	1	.	.	.
$E_g \downarrow C_{4v}$	1	1	.	.	.
$T_{1g} \downarrow C_{4v}$.	.	1	.	1
$T_{2g} \downarrow C_{4v}$.	.	.	1	1
$A_{1g} \downarrow C_{4v}$.	.	1	.	.
$A_{2u} \downarrow C_{4v}$.	.	.	1	.
$E_u \downarrow C_{4v}$.	.	1	1	.
$T_{1u} \downarrow C_{4v}$	1	.	.	.	1
$T_{2u} \downarrow C_{4v}$.	1	.	.	1

Also at MolVibes - $O_h - C_{4v}$ correlation table (eps)

Other correlation tables



Review Octahedral $O_h \supset O$ group operator structure

Review Octahedral $O_h \supset O \supset D_4 \supset C_4$ subgroup chain correlations

Comparison of $O \supset D_4 \supset C_4$ and $O \supset D_4 \supset D_2$ correlations and level/projector splitting

$O \supset D_4 \supset C_4$ subgroup chain splitting

$O \supset D_4 \supset D_2$ subgroup chain splitting (normal D_2 vs. unnormal D_2)

$O_h \supset O \supset D_4 \supset C_{4v}$ and $O_h \supset O \supset D_4 \supset C_{4v} \supset C_{2v}$ subgroup splitting

Splitting O class projectors P^μ into irreducible projectors $P^\mu_{m+m_4}$ for $O \supset C_4$

Development of irreducible projectors $P^\mu_{m+m_4}$ and representations $D^\mu_{m+m_4}$

Calculating $P^{E_{0404}}$, $P^{E_{2424}}$, $P^{T_{10404}}$, $P^{T_{11414}}$, $P^{T_{22424}}$, $P^{T_{21414}}$,



$O \supset C_4$ induced representation $O_4(C_4) \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples

Elementary induced representation $O_4(C_4) \uparrow O$

Projection reduction of induced representation $O_4(C_4) \uparrow O$

Introduction to ortho-complete eigenvalue-parameter relations


Examples in SF_6 spectroscopy



$$\begin{aligned} \ell^{A_1} &= 1 \\ \ell^{A_2} &= 1 \\ \ell^E &= 2 \\ \ell^{T_1} &= 3 \\ \ell^{T_2} &= 3 \end{aligned}$$

Example: $G=O$ Centrum: $\kappa(O) = \sum_{(\alpha)} (\ell^\alpha)^0 = 1^0 + 1^0 + 2^0 + 3^0 + 3^0 = 5$
 Cubic-Octahedral Rank: $\rho(O) = \sum_{(\alpha)} (\ell^\alpha)^1 = 1^1 + 1^1 + 2^1 + 3^1 + 3^1 = 10$
 Group O Order: $o(O) = \sum_{(\alpha)} (\ell^\alpha)^2 = 1^2 + 1^2 + 2^2 + 3^2 + 3^2 = 24$

O group	$\chi_{\kappa_g}^\alpha$	$g = 1$	r_{1-4}	ρ_{xyz}	R_{xyz}	i_{1-6}
			\tilde{r}_{1-4}		\tilde{R}_{xyz}	
s -orbital r^2	$\alpha = A_1$	1	1	1	1	1
d -orbitals	A_2	1	1	1	-1	-1
$\{x^2+y^2-2z^2, x^2-y^2\}$	E	2	-1	2	0	0
p -orbitals $\{x, y, z\}$	T_1	3	0	-1	1	-1
$\{xz, yz, xy\}$	T_2	3	0	-1	-1	1
d -orbitals						



$O \supset C_4$ $(0)_4 (1)_4 (2)_4 (3)_4 = (-1)_4$ $O \supset C_3$ $(0)_3 (1)_3 (2)_3 = (-1)_3$

A_1	1	•	•	•
A_2	•	•	1	•
E	1	•	1	•
T_1	1	1	•	1
T_2	•	1	1	1

A_1	1	•	•
A_2	1	•	•
E	•	1	1
T_1	1	1	1
T_2	1	1	1

$O \supset C_4$	0_4	1_4	2_4	3_4	$\mathbf{1} \cdot \mathbf{P}^\alpha = (\mathbf{p}_{0_4} + \mathbf{p}_{1_4} + \mathbf{p}_{2_4} + \mathbf{p}_{3_4}) \cdot \mathbf{P}^\alpha$	<i>where:</i> $\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 e^{im \cdot p/4} \mathbf{R}_z^p$ $\mathbf{p}_{m_4} = \begin{cases} \mathbf{p}_{0_4} = (1 + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z) / 4 \\ \mathbf{p}_{1_4} = (1 + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z) / 4 \\ \mathbf{p}_{2_4} = (1 - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z) / 4 \\ \mathbf{p}_{3_4} = (1 - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z) / 4 \end{cases}$
$A_1 \downarrow C_4$	1	.	.	.	$\mathbf{1} \cdot \mathbf{P}^{A_1} = \mathbf{P}_{0_4 0_4}^{A_1} + 0 + 0 + 0$	
$A_2 \downarrow C_4$.	.	1	.	$\mathbf{1} \cdot \mathbf{P}^{A_2} = 0 + 0 + \mathbf{P}_{2_4 2_4}^{A_2} + 0$	
$E \downarrow C_4$	1	.	1	.	$\mathbf{1} \cdot \mathbf{P}^E = \mathbf{P}_{0_4 0_4}^E + 0 + \mathbf{P}_{2_4 2_4}^E + 0$	
$T_1 \downarrow C_4$	1	1	.	1	$\mathbf{1} \cdot \mathbf{P}^{T_1} = \mathbf{P}_{0_4 0_4}^{T_1} + \mathbf{P}_{1_4 1_4}^{T_1} + 0 + \mathbf{P}_{3_4 3_4}^{T_1}$	
$T_2 \downarrow C_4$.	1	1	1	$\mathbf{1} \cdot \mathbf{P}^{T_2} = 0 + \mathbf{P}_{1_4 1_4}^{T_2} + \mathbf{P}_{2_4 2_4}^{T_2} + \mathbf{P}_{3_4 3_4}^{T_2}$	

$\mathbf{P}_{n_4 n_4}^{(\alpha)} (O \supset C_4)$	1	$r_1 r_2 \tilde{r}_3 \tilde{r}_4$	$\tilde{r}_1 \tilde{r}_2 r_3 r_4$	$\rho_x \rho_y$	ρ_z	$R_x \tilde{R}_x R_y \tilde{R}_y$	R_z	\tilde{R}_z	$i_1 i_2 i_5 i_6$	$i_3 i_4$
$24 \cdot \mathbf{P}_{0_4 0_4}^{A_1}$	1	1	1	1	1	1	1	1	1	1
$24 \cdot \mathbf{P}_{2_4 2_4}^{A_2}$	1	1	1	1	1	-1	-1	-1	-1	-1
$12 \cdot \mathbf{P}_{0_4 0_4}^E$	1	$-\frac{1}{2}$	$-\frac{1}{2}$	1	1	$-\frac{1}{2}$	1	1	$-\frac{1}{2}$	1
$12 \cdot \mathbf{P}_{2_4 2_4}^E$	1	$-\frac{1}{2}$	$-\frac{1}{2}$	1	1	$+\frac{1}{2}$	-1	-1	$+\frac{1}{2}$	-1
$8 \cdot \mathbf{P}_{1_4 1_4}^{T_1}$	1	$-\frac{i}{2}$	$+\frac{i}{2}$	0	-1	$+\frac{1}{2}$	$-i$	$+i$	$-\frac{1}{2}$	0
$8 \cdot \mathbf{P}_{3_4 3_4}^{T_1}$	1	$+\frac{i}{2}$	$-\frac{i}{2}$	0	-1	$+\frac{1}{2}$	$+i$	$-i$	$-\frac{1}{2}$	0
$8 \cdot \mathbf{P}_{0_4 0_4}^{T_1}$	1	0	0	-1	1	0	1	1	0	-1
$8 \cdot \mathbf{P}_{1_4 1_4}^{T_2}$	1	$+\frac{i}{2}$	$-\frac{i}{2}$	0	-1	$-\frac{1}{2}$	$-i$	$+i$	$+\frac{1}{2}$	0
$8 \cdot \mathbf{P}_{3_4 3_4}^{T_2}$	1	$-\frac{i}{2}$	$+\frac{i}{2}$	0	-1	$-\frac{1}{2}$	$+i$	$-i$	$+\frac{1}{2}$	0
$8 \cdot \mathbf{P}_{2_4 2_4}^{T_2}$	1	0	0	-1	1	0	-1	-1	0	1

*Summary of
 $O \supset C_4$
diagonal
(idempotent)
projectors
 \mathbf{P}_{jj}^μ*

$O \supset C_4$	0_4	1_4	2_4	3_4	$\mathbf{1} \cdot \mathbf{P}^\alpha = (\mathbf{p}_{0_4} + \mathbf{p}_{1_4} + \mathbf{p}_{2_4} + \mathbf{p}_{3_4}) \cdot \mathbf{P}^\alpha$	<p>where: $\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 e^{im \cdot p/4} \mathbf{R}_z^p$</p> <p>$\mathbf{p}_{0_4} = (1 + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z) / 4$ $\mathbf{p}_{1_4} = (1 + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z) / 4$ $\mathbf{p}_{2_4} = (1 - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z) / 4$ $\mathbf{p}_{3_4} = (1 - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z) / 4$</p>
$A_1 \downarrow C_4$	1	.	.	.	$\mathbf{1} \cdot \mathbf{P}^{A_1} = \mathbf{P}_{0_4 0_4}^{A_1} + 0 + 0 + 0$	
$A_2 \downarrow C_4$.	.	1	.	$\mathbf{1} \cdot \mathbf{P}^{A_2} = 0 + 0 + \mathbf{P}_{2_4 2_4}^{A_2} + 0$	
$E \downarrow C_4$	1	.	1	.	$\mathbf{1} \cdot \mathbf{P}^E = \mathbf{P}_{0_4 0_4}^E + 0 + \mathbf{P}_{2_4 2_4}^E + 0$	
$T_1 \downarrow C_4$	1	1	.	1	$\mathbf{1} \cdot \mathbf{P}^{T_1} = \mathbf{P}_{0_4 0_4}^{T_1} + \mathbf{P}_{1_4 1_4}^{T_1} + 0 + \mathbf{P}_{3_4 3_4}^{T_1}$	
$T_2 \downarrow C_4$.	1	1	1	$\mathbf{1} \cdot \mathbf{P}^{T_2} = 0 + \mathbf{P}_{1_4 1_4}^{T_2} + \mathbf{P}_{2_4 2_4}^{T_2} + \mathbf{P}_{3_4 3_4}^{T_2}$	

Summary of $O \supset C_4$ diagonal (idempotent) projectors

$\mathbf{P}_{n_4 n_4}^{(\alpha)} (O \supset C_4)$	$\mathbf{1}$	$r_1 r_2 \tilde{r}_3 \tilde{r}_4$	$\tilde{r}_1 \tilde{r}_2 r_3 r_4$	$\rho_x \rho_y$	ρ_z	$R_x \tilde{R}_x R_y \tilde{R}_y$	R_z	\tilde{R}_z	$i_1 i_2 i_5 i_6$	$i_3 i_4$
$24 \cdot \mathbf{P}_{0_4 0_4}^{A_1}$	1	1	1	1	1	1	1	1	(+1)	1
$24 \cdot \mathbf{P}_{2_4 2_4}^{A_2}$	1	1	1	1	1	-1	-1	-1	-1	-1
$12 \cdot \mathbf{P}_{0_4 0_4}^E$	1	$-\frac{1}{2}$	$-\frac{1}{2}$	1	1	$-\frac{1}{2}$	1	1	($-\frac{1}{2}$)	1
$12 \cdot \mathbf{P}_{2_4 2_4}^E$	1	$-\frac{1}{2}$	$-\frac{1}{2}$	1	1	$+\frac{1}{2}$	-1	-1	$+\frac{1}{2}$	-1
$8 \cdot \mathbf{P}_{1_4 1_4}^{T_1}$	1	$-\frac{i}{2}$	$+\frac{i}{2}$	0	-1	$+\frac{1}{2}$	-i	+i	$-\frac{1}{2}$	0
$8 \cdot \mathbf{P}_{3_4 3_4}^{T_1}$	1	$+\frac{i}{2}$	$-\frac{i}{2}$	0	-1	$+\frac{1}{2}$	+i	-i	$-\frac{1}{2}$	0
$8 \cdot \mathbf{P}_{0_4 0_4}^{T_1}$	1	0	0	-1	1	0	1	1	(0)	-1
$8 \cdot \mathbf{P}_{1_4 1_4}^{T_2}$	1	$+\frac{i}{2}$	$-\frac{i}{2}$	0	-1	$-\frac{1}{2}$	-i	+i	$+\frac{1}{2}$	0
$8 \cdot \mathbf{P}_{3_4 3_4}^{T_2}$	1	$-\frac{i}{2}$	$+\frac{i}{2}$	0	-1	$-\frac{1}{2}$	+i	-i	$+\frac{1}{2}$	0
$8 \cdot \mathbf{P}_{2_4 2_4}^{T_2}$	1	0	0	-1	1	0	-1	-1	0	1

The $0_4 \uparrow$ cluster
 i_{16} split i_{34} split


$\mathbf{P}_{0_4 0_4}^{A_1} \underline{\underline{+1}}$
 $\mathbf{P}_{0_4 0_4}^{T_1} \underline{\underline{0}}$
 $\mathbf{P}_{0_4 0_4}^E \underline{\underline{-1/2}}$

$O \supset C_4$	0_4	1_4	2_4	3_4	$\mathbf{1} \cdot \mathbf{P}^\alpha = (\mathbf{p}_{0_4} + \mathbf{p}_{1_4} + \mathbf{p}_{2_4} + \mathbf{p}_{3_4}) \cdot \mathbf{P}^\alpha$	<i>where:</i> $\mathbf{p}_{m_4} = \frac{1}{4} \sum_{p=0}^3 e^{im \cdot p/4} \mathbf{R}_z^p$ $\mathbf{p}_{0_4} = (1 + \mathbf{R}_z + \rho_z + \tilde{\mathbf{R}}_z) / 4$ $\mathbf{p}_{1_4} = (1 + i\mathbf{R}_z - \rho_z - i\tilde{\mathbf{R}}_z) / 4$ $\mathbf{p}_{2_4} = (1 - \mathbf{R}_z + \rho_z - \tilde{\mathbf{R}}_z) / 4$ $\mathbf{p}_{3_4} = (1 - i\mathbf{R}_z - \rho_z + i\tilde{\mathbf{R}}_z) / 4$
$A_1 \downarrow C_4$	1	.	.	.	$\mathbf{1} \cdot \mathbf{P}^{A_1} = \mathbf{P}_{0_4 0_4}^{A_1} + 0 + 0 + 0$	
$A_2 \downarrow C_4$.	.	1	.	$\mathbf{1} \cdot \mathbf{P}^{A_2} = 0 + 0 + \mathbf{P}_{2_4 2_4}^{A_2} + 0$	
$E \downarrow C_4$	1	.	1	.	$\mathbf{1} \cdot \mathbf{P}^E = \mathbf{P}_{0_4 0_4}^E + 0 + \mathbf{P}_{2_4 2_4}^E + 0$	
$T_1 \downarrow C_4$	1	1	.	1	$\mathbf{1} \cdot \mathbf{P}^{T_1} = \mathbf{P}_{0_4 0_4}^{T_1} + \mathbf{P}_{1_4 1_4}^{T_1} + 0 + \mathbf{P}_{3_4 3_4}^{T_1}$	
$T_2 \downarrow C_4$.	1	1	1	$\mathbf{1} \cdot \mathbf{P}^{T_2} = 0 + \mathbf{P}_{1_4 1_4}^{T_2} + \mathbf{P}_{2_4 2_4}^{T_2} + \mathbf{P}_{3_4 3_4}^{T_2}$	

Summary of $O \supset C_4$ diagonal (idempotent) projectors

$\mathbf{P}_{n_4 n_4}^{(\alpha)} (O \supset C_4)$	1	$r_1 r_2 \tilde{r}_3 \tilde{r}_4$	$\tilde{r}_1 \tilde{r}_2 r_3 r_4$	$\rho_x \rho_y$	ρ_z	$R_x \tilde{R}_x R_y \tilde{R}_y$	R_z	\tilde{R}_z	\mathbf{P}_{jj}^{μ}	
									$i_1 i_2 i_5 i_6$	$i_3 i_4$
$24 \cdot \mathbf{P}_{0_4 0_4}^{A_1}$	1	1	1	1	1	1	1	1	+1	(+1)
$24 \cdot \mathbf{P}_{2_4 2_4}^{A_2}$	1	1	1	1	1	-1	-1	-1	-1	-1
$12 \cdot \mathbf{P}_{0_4 0_4}^E$	1	-1/2	-1/2	1	1	-1/2	1	1	-1/2	(+1)
$12 \cdot \mathbf{P}_{2_4 2_4}^E$	1	-1/2	-1/2	1	1	+1/2	-1	-1	+1/2	-1
$8 \cdot \mathbf{P}_{1_4 1_4}^{T_1}$	1	-i/2	+i/2	0	-1	+1/2	-i	+i	-1/2	0
$8 \cdot \mathbf{P}_{3_4 3_4}^{T_1}$	1	+i/2	-i/2	0	-1	+1/2	+i	-i	-1/2	0
$8 \cdot \mathbf{P}_{0_4 0_4}^{T_1}$	1	0	0	-1	1	0	1	1	0	(-1)
$8 \cdot \mathbf{P}_{1_4 1_4}^{T_2}$	1	+i/2	-i/2	0	-1	-1/2	-i	+i	+1/2	0
$8 \cdot \mathbf{P}_{3_4 3_4}^{T_2}$	1	-i/2	+i/2	0	-1	-1/2	+i	-i	+1/2	0
$8 \cdot \mathbf{P}_{2_4 2_4}^{T_2}$	1	0	0	-1	1	0	-1	-1	0	1

The $0_4 \uparrow$ cluster
 i_{16} split *i_{34} split*



Elementary induced representation $O_4(C_4) \uparrow O$





Fig. 4.3.1 PSDS

Thus we label states by left cosets $r_l C_4$ of Local C_4

$$\{|1\rangle = |1\rangle, |2\rangle = |R_1^2\rangle, |3\rangle = |r_1\rangle, |4\rangle = |r_2\rangle, |5\rangle = |r_1^2\rangle, |6\rangle = |r_2^2\rangle\}$$

$$1C_4 = C_4 = \{1, R_3, R_3^2, R_3^3\}$$

$$1C_4 = C_4 = \{1, R_3, R_3^2, R_3^3\}$$

$$R_1^2(1, R_3, R_3^2, R_3^3) = (R_1^2, i_4, R_2^2, i_3),$$

$$r_1(1, R_3, R_3^2, R_3^3) = (r_1, i_1, r_4, R_2),$$

$$r_2(1, R_3, R_3^2, R_3^3) = (r_2, i_2, r_3, R_2^3),$$

$$r_1^2(1, R_3, R_3^2, R_3^3) = (r_1^2, R_1^3, r_3^2, i_6),$$

$$r_2^2(1, R_3, R_3^2, R_3^3) = (r_2^2, R_1, r_4^2, i_5),$$

Here is $O_4(C_4)$ induced representation $\mathcal{J}^{O_4 \uparrow O}(\mathbf{I}_i)$ of a linear combination of \mathbf{i} -class rotations

$$\mathbf{I}_i = i_1 \mathbf{i}_1 + i_2 \mathbf{i}_2 + i_3 \mathbf{i}_3 + i_4 \mathbf{i}_4 + i_5 \mathbf{i}_5 + i_6 \mathbf{i}_6 \longrightarrow \mathbf{I}_i = i_{34} (\mathbf{i}_3 + \mathbf{i}_4) + i_{16} (\mathbf{i}_1 + \mathbf{i}_2 + \mathbf{i}_5 + \mathbf{i}_6)$$

$$\mathcal{J}^{O_4 \uparrow O}(\mathbf{I}_i) =$$

	$ 1\rangle$	$ 2\rangle$	$ 3\rangle$	$ 4\rangle$	$ 5\rangle$	$ 6\rangle$
$\langle 1 $	1	$i_3 + i_4$	i_1	i_2	i_6	i_5
$\langle 2 $	$i_3 + i_4$	1	i_2	i_1	i_5	i_6
$\langle 3 $	i_1	i_2	1	$i_5 + i_6$	i_3	i_4
$\langle 4 $	i_2	i_1	$i_5 + i_6$	1	i_4	i_3
$\langle 5 $	i_6	i_5	i_3	i_4	1	$i_1 + i_2$
$\langle 6 $	i_5	i_6	i_4	i_3	$i_1 + i_2$	1

Let:

$$i_3 = i_{34} = i_4$$

and/or:

$$i_{16} = i_1 = i_2 = i_5 = i_6$$

	$ 1\rangle$	$ 2\rangle$	$ 3\rangle$	$ 4\rangle$	$ 5\rangle$	$ 6\rangle$
$\langle 1 $	1	$2i_{34}$	i_{16}	i_{16}	i_{16}	i_{16}
$\langle 2 $	$2i_{34}$	1	i_{16}	i_{16}	i_{16}	i_{16}
$\langle 3 $	i_{16}	i_{16}	1	$2i_{16}$	i_{34}	i_{34}
$\langle 4 $	i_{16}	i_{16}	$2i_{16}$	1	i_{34}	i_{34}
$\langle 5 $	i_{16}	i_{16}	i_{34}	i_{34}	1	$2i_{16}$
$\langle 6 $	i_{16}	i_{16}	i_{34}	i_{34}	$2i_{16}$	1

Review Octahedral $O_h \supset O$ group operator structure

Review Octahedral $O_h \supset O \supset D_4 \supset C_4$ subgroup chain correlations

Comparison of $O \supset D_4 \supset C_4$ and $O \supset D_4 \supset D_2$ correlations and level/projector splitting

$O \supset D_4 \supset C_4$ subgroup chain splitting

$O \supset D_4 \supset D_2$ subgroup chain splitting (normal D_2 vs. unnormal D_2)

$O_h \supset O \supset D_4 \supset C_{4v}$ and $O_h \supset O \supset D_4 \supset C_{4v} \supset C_{2v}$ subgroup splitting

Splitting O class projectors P^μ into irreducible projectors $P^\mu_{m+m_4}$ for $O \supset C_4$

Development of irreducible projectors $P^\mu_{m+m_4}$ and representations $D^\mu_{m+m_4}$

Calculating $P^{E_{0404}}$, $P^{E_{2424}}$, $P^{T_{10404}}$, $P^{T_{11414}}$, $P^{T_{22424}}$, $P^{T_{21414}}$,



$O \supset C_4$ induced representation $0_4(C_4) \uparrow O \sim A_1 \oplus T_1 \oplus E$ and spectral analysis examples

Elementary induced representation $0_4(C_4) \uparrow O$


Projection reduction of induced representation $0_4(C_4) \uparrow O$

Introduction to ortho-complete eigenvalue-parameter relations


Examples in SF_6 spectroscopy



Angular Momentum Cones and Quantum Polar Angles




(a) SF₆ ν₄ Rotational Structure




FT IR and Laser Diode Spectra
K.C. Kim, W.B. Person, D. Seitz, and B.J. Krohn
J. Mol. Spectrosc. 76, 322 (1979).

Primary AET species mixing increases with distance from "separatrix"


(b) P(88) Fine Structure (Rotational anisotropy effects)



SF₆ ν₃ P(88) ~ 16m



(c) Superfine Structure (Rotational axis tunneling)



Observed repeating sequence(s) .. **A₁ T₁ E T₂ T₁ E T₂ A₂ T₂ T₁** A₁ T₁ E T₂ T₁ E T₂ A₂ T₂ T₁ A₁ ..

$O=C_4$ (0)₄ (1)₄ (2)₄ (3)₄ = (-1)₄

A ₁	1	•	•	•
A ₂	•	•	1	•
E	1	•	1	•
T ₁	1	1	•	1
T ₂	•	1	1	1

$O=C_3$ (0)₃ (1)₃ (2)₃ = (-1)₃

A ₁	1	•	•
A ₂	1	•	•
E	•	1	1
T ₁	1	1	1
T ₂	1	1	1

Local correlations explain clustering...
... but what about spacing and ordering?...

...and physical consequences?

Major mixing lowest two LUSTERS

Eigenvalues of $\mathbf{H} = B\mathbf{J}^2 + \cos\phi\mathbf{T}^{[4]} + \sin\phi\mathbf{T}^{[6]}$ vs. mix angle $\phi: 0 < \phi < \pi$

